



HAL
open science

Breeding for wheat-pea mixtures: are the traits of pea varieties in sole crop predictive of their behaviour in mixture?

Nathalie Moutier, Matthieu Floriot, Cécile Le Gall, Christine Fintz, Mathieu Conseil, Laurence Fontaine, Alain Baranger

► To cite this version:

Nathalie Moutier, Matthieu Floriot, Cécile Le Gall, Christine Fintz, Mathieu Conseil, et al.. Breeding for wheat-pea mixtures: are the traits of pea varieties in sole crop predictive of their behaviour in mixture?. EUCARPIA International conference “Breeding and seed sector innovations for organic food systems”, Mar 2021, Cesis, Latvia. hal-03475738

HAL Id: hal-03475738

<https://hal.inrae.fr/hal-03475738v1>

Submitted on 11 Dec 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

BREEDING FOR WHEAT-PEA MIXTURES: ARE THE TRAITS OF PEA VARIETIES IN SOLE CROP PREDICTIVE OF THEIR BEHAVIOR IN MIXTURE?

Nathalie MOUTIER¹, Matthieu FLORIOT², Cécile LE GALL³, Christine FINTZ⁴,
Mathieu CONSEIL⁵, Laurence FONTAINE⁴, Alain BARANGER¹

¹ INRAE, UMR IGEPP, Le Rheu, France,

² Agri-Obtentions, Orsonville, France,

³ Terres Inovia, Thiverval-Grignon, France,

⁴ GEVES, Beaucouzé, France,

⁵ ITAB, Angers, France

nathalie.moutier@inrae.fr

Keywords: intercropping, grain legumes, genotype x management interaction, cultivar selection

In France, the production of grain legumes has declined significantly over the past 20 years. Two of the main technical issues in pea cropping are the instability of yields, linked to the effects of abiotic and biotic stresses, and the high lodging rate at harvesting. An alternative to overcome these limiting factors is to grow peas in mixture with a cereal to better control canopy lodging, diseases and pests, and generate additional income. This is a widespread practice in Organic Farming.

One emerging issue is whether the choice of the pea cultivar that would be best suited to be mixed with the cereal should be based on identified varietal traits, already assessed in sole crop. We therefore explored to what extent the traits and performances evaluated in sole crop are predictive of those observed in mixtures? We compared pea varieties in sole crop and in mixture with bread winter wheat, for traits likely to ensure the success of the mixture, including phenology (such as beginning and end of flowering dates), morphology (such as pea height in the canopy at the beginning, at the end of flowering, and at the end of the cycle), and finally productivity (such as thousand grains weight and grain yield). Six trials were carried out in Organic Farming (*) or very low input (**) systems during the 2015/2016 to 2018/2019 seasons, in the Parisian Basin (Orsonville**), the north-west (Rennes*) and the center (Civray**) of France. Five to nine winter pea cultivars with contrasting phenological, morphological and productivity traits were sown as sole crop and mixed (at 75% of the sole crop doses) on the row with a single bread winter wheat cultivar.

In our environmental conditions:

- The dates of beginning and end of flowering, and the thousand grain weights of pea cultivars were very similar in mixture and in sole crop. The values of these traits in sole crop therefore seem to be predictive of those in mixture.
- All pea varieties were higher in mixture than in sole crop, whatever the developmental stage, confirming the ongoing tutor effect of wheat on pea.
- The pea variety heights at the start and end of flowering in sole crop seem to be predictive of those observed in mixture.
- Both for pea varieties height at the end of the cycle and for pea varieties grain yield, the values in sole crop did not seem to be predictive of those observed in mixture. Consequently, the development of specific breeding schemes and ratings for the evaluation of these traits in mixture appears necessary.

Acknowledgements

This work was supported by funding from the special account "agricultural and rural development" CASDAR managed by the French Ministry of Agriculture, Agri-food and Forests "ECoVAB" project and from the European Union's Horizon 2020 Research and Innovation Programme "Remix" project under grant agreement n°727217.