



**HAL**  
open science

## Increased exopolysaccharide production and microbial activity affect soil water retention and field performance of tomato under water deficit

Samuel Le Gall, Annette Bérard, David Page, L. Lanoe, Nadia Bertin, Claude Doussan

### ► To cite this version:

Samuel Le Gall, Annette Bérard, David Page, L. Lanoe, Nadia Bertin, et al.. Increased exopolysaccharide production and microbial activity affect soil water retention and field performance of tomato under water deficit. *Rhizosphere*, 2021, 19, 10.1016/j.rhisph.2021.100408 . hal-03480484

**HAL Id: hal-03480484**

**<https://hal.inrae.fr/hal-03480484v1>**

Submitted on 2 Aug 2023

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial 4.0 International License

## Increased exopolysaccharide production and microbial activity affect soil water retention and field performance of tomato under water deficit

Le Gall S.<sup>1,5\*</sup>, Bérard A<sup>1.</sup>, Page D.<sup>2.</sup>, Lanoe L.<sup>3.</sup>, Bertin N.<sup>4.</sup>, Doussan C.<sup>1</sup>

<sup>1</sup> INRAE, Avignon Université, EMMAH, F-84000, Avignon, France

<sup>2</sup> INRAE, Avignon Université, SQPOV, F-84000, Avignon, France

<sup>3</sup> SONITO, F-84000, Avignon, Franc

<sup>4</sup> INRAE, Avignon Université, PSH, F-84000, Avignon, France

<sup>5</sup> Present address: Agrosphere IBG-3, Forschungszentrum Jülich, Jülich, NRW, Germany

\* Corresponding author: [s.le.gall@fz-juelich.de](mailto:s.le.gall@fz-juelich.de)

1 Increased exopolysaccharide production and microbial activity affect soil  
2 water retention and field performance of tomato under water deficit

### 3 Summary

4 According to the literature, biological processes in the rhizosphere could play a role in the  
5 adaptation of plants to droughts under a changing climate. A previous study has identified  
6 significantly different productivity level and fruit quality for two tomato varieties under wa-  
7 ter deficit conditions. We conducted a field study, with and without water deficit, with these  
8 two varieties to examine whether microbiological activity and exopolysaccharides concen-  
9 tration could affect soil hydrophysical properties. The rhizosphere soil had indeed distinct  
10 bio-chemical and hydrophysical properties between the two cultivars and between the two  
11 water-related conditions. The quantity of soil exopolysaccharide and/or nitrogenous sub-  
12 stances, and the activity of microorganisms (fungi in particular) explains part of the soil wa-  
13 ter retention measurements. In addition, these mechanisms are significantly accentuated for  
14 the cultivar with the best productive capability under water-limited condition—i.e. with  
15 commercial yield, fruit dry matter and water use efficiency which are respectively 35%, 28%,  
16 and 31% higher for the productive cultivar.

### 17 Key words

18 Rhizosphere, Water deficit, Soil water retention, Exopolysaccharide, Microbial activity

## 19 Introduction

20 The current increase in temperature, evapotranspiration, and in the seasonal variation of  
21 water regimes linked to climate change lead to a higher pressure on the water resources,  
22 which affects the agricultural production (Jia et al., 2019, Boeck et al., 2011). This calls for  
23 the development of more water-efficient agroecological systems, able to withstand more  
24 frequent water deficit periods (Lipiec et al., 2013). The water retention capacity of soils has a  
25 strong influence on their capacity to satisfy crops' water requirements and could thus miti-  
26 gate the effects of extreme climatic events (Łabędzki, 2016).

27 The rhizosphere is the area of soil around the roots influenced by plant roots and where the  
28 plant, the soil, and the associated microorganisms interact (Bargett et al., 2014; Hinsinger et  
29 al., 2009; Angers et al., 1998). The rhizosphere is the portion of soil that is physically in con-  
30 tact with the root system. The rhizosphere is part of the rhizosphere, but rhizosphere may  
31 extend beyond the rhizosphere (Marasco et al., 2018; Pang et al., 2017). The rhizosphere is  
32 both a "hot spot" of biological activity and metabolic processes in the soil (Zhang et al.,  
33 2020), and the zone where root water and nutrient uptake takes place. Moreover, the rhizo-  
34 sphere has specific physicochemical properties: soil aggregates are more stable; bulk densi-  
35 ty, porosity, water and nutrients transfer as well as pH are modified with respect to the bulk  
36 soil (Hinsinger et al., 2009). These properties are linked to the specific biochemical activities  
37 of roots, such as rhizodeposition and exudation of organic compounds (Personeni et al.,  
38 2017), which can then be consumed by the associated microbial communities (van Veelen et al.  
39 2018). In addition, these microbial communities, stimulated by the rhizosphere environ-  
40 ment, can also excrete similar organic molecules, including Exopolysaccharides (EPS),  
41 such as Exopolysaccharides (EPS) (Redmile-Gordon et al., 2014). EPS may also consti-

42 tute the majority of root mucilage (Nazari, 2021). These polymeric substances contribute to  
43 the stability of soil aggregates (Crouzet et al., 2019), improve soil water retention (Czarnes et  
44 al., 2000), and present either hydrophilic or hydrophobic characteristics depending on their  
45 state of hydration (Carminati and Vetterlein, 2013). Until now, few studies have focused on  
46 the influence of both the physicochemical and biological properties of the rhizosphere on  
47 plant water use efficiency (WUE) (Benard et al. 2019). However, since all the water tran-  
48 spired by a plant must cross the rhizosphere soil layer, understanding its functioning and its  
49 properties is critical (Bengough, 2012). Thus, acting on the functioning of the rhizosphere  
50 could be a lever to maintain the productivity of crops suffering from water deficit (Ahmed et  
51 al., 2018, Doussan et al., 2015).

52 Tomato (*Solanum lycopersicum*) is widely grown in the Mediterranean region and its produc-  
53 tion is one of the most important worldwide in terms of market value and quantity (FAO-  
54 STAT, n.d.). Understanding how this crop adapts to droughts is therefore of prime im-  
55 portance. Under glasshouse conditions, water deficit negatively affects plant growth, leaf  
56 water potential and gas exchange intensity during the early stages of tomato development  
57 (Duan et al., 2016) as well as fruit mass and composition at the latter stages (Ripoll et al.  
58 2016). The impact of water deficit was also observed in the field for industrial tomato culti-  
59 vars (Arbex de Castro Vilas Boas et al., 2017). In particular, under water deficit, the cultivar  
60 Terradou, compared to H1015, showed a higher dry yield (on average 76 kg.ha<sup>-1</sup> and 56.kg  
61 ha<sup>-1</sup> respectively) and a higher Water Use Efficiency (WUE) (41 and 33 kg.m<sup>-3</sup> respectively).  
62 Terradou fruits had also a higher sugar content (Glucose, Fructose, higher SSC index) while  
63 H1015 fruits had more acids (Malic, Citric). The fruit puree viscosity increased in general  
64 when the plants underwent water deficit, but more markedly for H1015 than for Terradou.

65 As water deficit is one of the main factors limiting yield in tomatoes (Costa et al., 2007;  
66 Patanè and Consentino, 2010), depending on its intensity, duration, and timing in the crop  
67 cycle (Rinaldi et al., 2007), the response of these cultivars to water deficit deserves further  
68 testing.

69 While several studies have investigated the influence of microbial activity, and in particular  
70 mycorrhizae (Cavagnaro et al., 2006; Hallett et al., 2009; Bowles et al., 2016), on tomato  
71 phenology and its yield components, it is worthwhile to investigate the influence of soil hy-  
72 drophysical properties. Understanding this possible link between soil bio-chemical and hy-  
73 drophysical parameters might broaden our understanding of the plants' responses to  
74 drought. Our objective was therefore to observe in the field the links between different bio-  
75 chemical parameters of the soil (i.e., microbial biomass and activities, EPSac amounts) and  
76 the soil water retention (i.e. Field Capacity and Permanent Wilting Point). We took into ac-  
77 count the influence of (i) the rhizosphere of tomatoes, (ii) the season stage of the plant de-  
78 velopment, (iii) the tomato cultivars, and (iv) the water conditions (well irrigated or deficit).

79 We carried out a field experiment with the two cultivars Terradou and H1015 on the same  
80 Mediterranean experimental site than previously used by Arbex de Castro Vilas Boas et al  
81 (2017). We increased water deficit by limiting irrigation, and we measured the biophysical  
82 parameters mentioned above both in the rhizosphere and in the bulk soil. Our hypothesis  
83 was that the EPS exudation and microbial activities would be higher for Terradou compared  
84 to H1015, and that these processes would allow for a better yield resilience to water deficit  
85 conditions for Terradou. We also assumed that these processes would be relatively more  
86 important in the rhizosphere and that their dynamics would follow the development stages  
87 of the plant.

## 88 Materials and methods

### 89 Experimental set-up and sampling

90 The experimental site is an industrial tomato field of a SONITO (French interprofessional or-  
91 ganisation of the tomato sector). It is located under a Mediterranean climate at Piolenc,  
92 France (44°11'16.9 "N 4°48'10.8 "E), and is used for a varietal trial. The soil is a Stagnosol  
93 (WRB classification; or Redoxisol in French classification) with a loam texture (main soil  
94 properties are shown in Table 1). We studied two widely grown commercial cultivars of to-  
95 mato (*Solanum lycopersicum* L.): Terradou (T) and H1015 (H), previously tested by Arbex de  
96 Castro Vilas Boas et al. (2017). The cultivation practices applied on this field correspond to  
97 current conventional practices (density of 3.3 plants.m<sup>-2</sup>, fertilization and pesticides). During  
98 this experimentation, two irrigation treatments were applied: from planting (May 18, 2018)  
99 to fructification (July 23, 2018), all the plants received the same amount of irrigation water  
100 via daily drip irrigation to compensate 87% of evapotranspiration losses, resulting in nearly  
101 460 mm of total water input before the fructification (135 mm from rainfalls and 325 mm  
102 from irrigation). Then, from fructification to harvest (August 9), half of the plots had water  
103 deficit (WD) conditions i.e., 57% of the water input of the fully irrigated plots (well-watered  
104 "WW"). Each cultivar was grown on two plots (42 m length each) of two rows (33 cm apart)  
105 of tomato plants on cultivation ridges distributed on both sides of the main irrigation tube.  
106 For each cultivar, one plot was WW and one plot was WD (Fig.S1).

107 Soil samples were collected at the beginning of the flowering stage (June 19, 2018) and later  
108 at the fructification stage (August 2, 2018) before harvesting. Five sub-plots were randomly  
109 chosen within the plots. In those sub-plots, we sampled two types of soil: the 'rhizosphere'

110 soil and the corresponding 'bulk' soil. For the rhizosphere soil sampling, three plants per cul-  
111 tivar in each sub-plot were excavated with their proximal root system and surrounding soil  
112 using a spading fork (to a depth of 20 cm maximum). The plants were then shaken vigorously  
113 by hand, but without breaking roots, until no more soil aggregates could be detached from  
114 roots. The soil remaining on root was collected and judged to be the rhizosphere soil. The  
115 rhizosphere soil of the three selected plants was pooled (Göttlein, 2006). We selected three  
116 subsamples of bulk soil located on the outside bare soil of the ridge facing the plants sam-  
117 pled for the rhizosphere soil over 0-20 cm depth (Fig S1). These three subsamples of bulk soil  
118 were also pooled. All soil samples were air-dried, sieved at 2 mm and stored at 4°C until  
119 analysis.

120 Measurements of soil and plant parameters

121 *Soil microbial and fungal catabolic measurements*

122 Substrate Induced Respiration (SIR) of soil was assessed with the FungiResp, adapted from  
123 the MicroResp<sup>TM</sup> colorimetric technique (Ben Sassi et al., 2012; Chapman et al., 2007), con-  
124 sisting in a 96-deep-well microplate filled with soil subsamples. The soil water content was  
125 preliminary adjusted to 40% of the water holding capacity (about 0.22 g/g gravimetric water  
126 content). 25 µl of organic substrates (6.7 mg.g<sup>-1</sup> dw soil) were added in each deep-well. Each  
127 filled soil plate was then sealed face to face with a CO<sub>2</sub> trap microplate including a pH dye  
128 indicator, and an intercalated silicone joint to individualize the plate wells. The systems were  
129 then incubated in the dark at (23°C±1) for six hours. Absorbance of the gel CO<sub>2</sub>-trap was  
130 measured at 570 nm, before and after incubation. A calibration curve of absorbance versus  
131 head space equilibrium CO<sub>2</sub> concentration (measured by gas chromatography) was fitted to a  
132 regression model, which was then used to compute the amounts of released CO<sub>2</sub> in µg C-



133 CO<sub>2</sub>.g<sup>-1</sup> Dry Soil.h<sup>-1</sup>. We measured the microbial basal respiration by adding 25 µL of demin-  
134 eralised water instead of organic substrate (M.Bas). In order to address substrate-induced  
135 respiration according to fungal communities, we applied a bacterial inhibitor (25 µL for a  
136 final concentration of 78 µg Bronopol.g<sup>-1</sup> soil, concentration selected after performing a  
137 dose/response curve on the respiration of this soil, Ben Sassi et al., 2012) to half of the soil  
138 sub-samples distributed in the deep well-plate (the other half of the soil sub-samples re-  
139 ceived 25 µL of demineralised water). Glucose-induced respiration was used as a proxy of  
140 active total microbial biomass (MB) and fungal biomass (FB) in µg C.g<sup>-1</sup> DrySoil (Anderson  
141 and Domsch, 1978). To access the catabolic profiles of fungal communities (F-CLPP) (F.Tre,  
142 F.Cell, F.Ala, F.Gly, F.Mal, F.GlcN) and microbial communities (M-CLPP) (M.Tre, M.Cell, M.Ala,  
143 M.Gly, M.Mal, M.GlcN) we used various organic substrates (trehalose, cellobiose, alanine,  
144 glycine, malate, glucosamine, respectively) distributed in the deep well-plate, and at the  
145 same amounts of glucose, with and without the addition of bacterial inhibitor (Bérard et al.,  
146 2011).

#### 147 *Soil water retention measurements*

148 The measurement of water retention was carried out for matric potentials of 0.3 bars (i.e.,  
149 Field Capacity) and 15 bars (i.e., Permanent Wilting Point) on soil aggregates by using a pres-  
150 sure plate system. The triplicates of initially saturated samples (about 10 g) were placed in  
151 porous plate enclosures into which nitrogen gas was injected at a pressure of 0.3 or 15 bars.  
152 Samples were left to drain excess water until equilibrium was reached, with no outflow. The  
153 gravimetric water content (g water.g<sup>-1</sup> dry soil) was then measured by weighing the soil be-  
154 fore and after drying at 105°C (24h).

#### 155 *Soil exopolysaccharides measurements*

156 The ExoPolySaccharides (EPSac) that are principally involved in soil physical properties are  
157 those tightly bound to the soil particles (Chen et al., 2014; Crouzet et al. 2019). We used the  
158 physico-chemical Cation Exchange Resin (CER) technique (Redmile-Gordon et al., 2014; Bér-  
159 ard et al., 2020) to extract the bound-ExoPolymeric Substance (bound-EPS) from the soil.  
160 Before bound-EPS extraction, loosely bound EPS were extracted (each 0.5 g of soil sample)  
161 using 5 mL  $\text{CaCl}_2$  ( $10^{-2}$  M) under agitation (50 rpm) for one hour at laboratory temperature  
162 (Redmile-Gordon et al., 2014; Bérard et al., 2020). The soil/ $\text{CaCl}_2$  mixtures were then centri-  
163 fuge (8000 xg for 15 min at  $10^\circ\text{C}$ ), and the supernatants were removed. The remaining cen-  
164 trifuged soil pellets were then extracted by the CER extraction technique. The CER (Dowex  
165 Marathon C  $\text{Na}^+$  form, Sigma Aldrich, Steinheim, Germany) was previously washed with a  
166 phosphate buffer (consisting of 2 mM  $\text{Na}_3\text{PO}_4 \cdot 12\text{H}_2\text{O}$ , 4 mM  $\text{NaH}_2\text{PO}_4 \cdot \text{H}_2\text{O}$ , 9 mM NaCl, 1  
167 mM KCl, adjusted to pH 7 with 1M HCl and stored at  $4^\circ\text{C}$ ) until the pH of the solution stabi-  
168 lized to 7. In a centrifuge tube containing the soil pellet samples, the CER was added to ob-  
169 tain a ratio of 70 g dry CER per 1 g of soil organic matter. Five mL of phosphate buffer pH 7  
170 were then added to the soil and CER. After manual stirring, the samples were incubated for  
171 sixteen hours overnight under agitation (50 rpm) at room temperature. Bound-EPSac were  
172 measured as the total carbohydrate content of the bound-EPS extracted from soil, using the  
173 phenol–sulphuric acid Dubois method (Dubois et al., 1956) with glucose as the standard.

#### 174 *Plant and Fresh Fruit Measurements*

175 At harvest, for each cultivar and irrigation modality, five tomato plants were sampled. We  
176 measured the commercial yield (CoY, in  $\text{t}\cdot\text{ha}^{-1}$ ) corresponding to the tomato marketable  
177 fresh yield for the industry (red and yellow ones), and harvest losses due to rotten tomatoes  
178 (RoY, in  $\text{t}\cdot\text{ha}^{-1}$ ). The dry mass of this commercial yield (DMY in  $\text{kg}\cdot\text{ha}^{-1}$ ) was also measured.

179 We calculated the water use efficiency (WUE -  $\text{kg}\cdot\text{m}^{-3}$ ) as the ratio between the commercial  
180 yield (CoY) and the total water volume provided by irrigation and rainfall to the plant, for a  
181 given surface. Some chemical analyses were also performed on the tomato fruits, such as  
182 the soluble solid content (SSC, in °Brix), the titrable acidity and the pH (as in Arbex de Castro  
183 Vilas Boas et al., 2017).

184 Data Analysis:

185 *Statistical analysis*

186 Four modalities were investigated in the experimental design: the type of soil—rhizosheath  
187 or bulk (SOIL), the season—June or August (SEAS), the cultivar—Terradou or H1015 (CULT),  
188 and the plots' water status modalities in August —WD or WW. 76 soil samples were ana-  
189 lysed. Quantitative data were analysed using R (R core Team, 2017) with R Studio IDE (R Stu-  
190 dio Team, 2020). We first observed the statistical differences between the three first modali-  
191 ties (SOIL, SEAS, CULT) for each of the soil parameter studied (microbial, hydro-physical, and  
192 chemical) at a level of significance of  $p < 0.05$  with an ANOVA F-test (the R stat package and  
193 anova() function). Table 2 shows the main effects and their interactions performed over the  
194 whole dataset of soil parameters and on rhizosheath soil dataset. Table 2 also presents spe-  
195 cific analyses for June, August-WD irrigation and August-WW irrigation datasets, allowing us  
196 to compare the WW and WD data. Homoscedasticity of variances (Bartlett test) and normali-  
197 ty (Shapiro test) were checked before data analyses. The mean and standard deviation for  
198 the groups of data analysed are reported in Table S1.

199 *Principal Component Analysis*

200 Each substrate-induced respiration measurement was normalized to the sum of all SIR  
201 measurements, to exclude the microbial biomass effect on the individual respirations (Ben  
202 Sassi et al. 2012). Principal Component Analysis (PCA) was performed on these normalized  
203 measurements of microbial and fungal catabolic profiles carried out on the seven organic  
204 substrates: glucose, trehalose, cellobiose, alanine, glycine, malate, glucosamine. The *Fac-*  
205 *toMineR* R package with the `fviz_pca_ind()` function (Lê et al., 2008) was used to perform  
206 PCA analysis. A first global PCA analysis was performed on dataset including the 7 microbial  
207 or fungal variables and the whole 76 observations (2 cultivars × 2 seasons × 2 soil types × 2  
208 irrigation levels × 5 replicates, 4 data missing). Then, we performed PCA analyses on the June  
209 data exclusively, as well as for the August-WW and the August-WD.

#### 210 *Linear regression models*

211 Multiple linear regression models (Schielzeth, 2010; Whittingham et al., 2006) with the `lm()`  
212 function in *stats* package were established to relate 0.3 bars and 15 bars water contents to  
213 soil candidate predictor variables: EPSac, microbiological and fungal activity and biomass  
214 measurements. A stepwise regression (Murtaugh, 2009) was used to select the best predic-  
215 tor variables using the AIC information criterion (Hegyi and Gramszegi, 2011) with the func-  
216 tion `ols_step_both_aic()` from the *o/ssa* R-package. The Variable Inflation Factor (VIF) was  
217 computed with the function `ols_vif_tol()` from the same package. The VIF must be lower  
218 than 10 for the variable to be selected (Dormann, 2013). This way, we obtained between  
219 one and four predictor variables in the linear regression models.

## 220 **Results**

221 How soil type, season and cultivar influence soil indicators

222 Significant differences are observed between most of the bio-physicochemical properties of  
223 the rhizosphere soil of tomato plants and the corresponding "bulk" soil far from roots (Tab.2  
224 and S1). Microbial and fungal biomass and activity are almost constantly significantly higher  
225 in rhizosphere. The same holds for water retention values at the Permanent Wilting Point  
226 (W15b) ( $p$ -value =  $1.26e-04$ ), the field capacity (W0.3b) showing the same trend ( $p$ -value =  
227  $6e-2$ ). A higher EPSac amount is also denoted in the rhizosphere on this global dataset (mean  
228 respectively equal to 780 and 760  $\mu\text{g.g}^{-1}$ ;  $p$ -value= $7e-2$ ) (Tab.2 and S1).

229 Microbiological activities, fungal activities, and the amount of EPSac are higher in August  
230 compared to June (highly significantly) (Tab.2 and S1). Concerning hydraulic parameters,  
231 W15b was also higher in August, but W0.3b is higher in June ( $p$ -value= $5.4e-2$  for both). This  
232 temporal difference in hydraulic parameters increases when considering only the rhi-  
233 zosphere soil (respectively  $p$ -value= $4.5e-3$  and  $1.2e-7$ ). In June, some physicochemical and  
234 microbial parameters show significantly higher values in the rhizosphere soil than in bulk  
235 soil, but for August-WW irrigation, most of the microbial indicators exhibit significant higher  
236 values in the rhizosphere soil compared to bulk soil (Tab.2 and S1).

237 Concerning the tomato cultivars, a number of microbial soil indicators are significantly high-  
238 er for the T compared to the H cultivar (Tab.2 and S1). In June, we observe differences in  
239 microbial indicators between H and T cultivars ( $T>H$ ) only for the rhizosphere soils. The soil  
240 indicators for these two varieties are then differentiated into two distinct behaviours accord-  
241 ing to the type of irrigation. Moreover, the analyses of variance show interactions between  
242 the factors season and soil, for the water retention variable W0.3b and for the microbial  
243 variables F.GlcN, F.Ala, M.Cell. The season and cultivar factors also show interactions, for the

244 variables W15b, EPS, M.Mal and in the rhizosphere for the variables W15b, F.Ala, M.Gly,  
245 M.Mal (Tab.2).

246 Differences between soil indicators exacerbated under water deficit

247 Moisture content of soil samples from August campaign sampling differed between WD and  
248 WW, with higher water content in WW plots compared to less irrigated WD plots (respective  
249 averages of 13.7 and 12.4 g.g<sup>-1</sup>, p-value=0.031). In August-WD, we observe differences in  
250 both W15b and W0.3b between rhizosphere and bulk soils, while only W15b values present  
251 significant differences in August-WW (Tab.2). However, the trend is the same for the two  
252 irrigation treatments: W15b values are higher in rhizosphere soils than in bulk soils while it is  
253 the contrary for W0.3b. For August-WD, the soil of T cultivar exhibits parameters (W0.3b and  
254 a number of microbial indicators) that are significantly higher than for H cultivar (for rhi-  
255 zosphere soils alone and for rhizosphere and bulk soils together). In August-WW, the differ-  
256 ences between cultivars are less pronounced than in August-WD and the ranking of cultivars  
257 is inverted: Some microbial and fungal catabolic activities present higher values in H soils  
258 compared to T soils (Tab.2 and S1).

259 Microbial and fungal catabolic profiles

260 The microbial and fungal catabolic profiles of the 76 soil samples were analysed with Princi-  
261 pal Component Analysis (PCA). Considering the catabolic profiles from the global dataset, no  
262 discrimination appears between cultivars (T/H) and soil types (R/B) for microbial (Supp. Fig.  
263 2-b, 2-c) and fungal (Fig. 1-b, 1-c) catabolic profiles. However, PCA shows differences be-  
264 tween August and June sampling seasons. In August catabolic activity of microbial communi-  
265 ties show more homogeneity between samples, with points concentrated in the center of  
266 the graph, compared to June (Supp. Fig.2, % total of variance=79.5). The fungal communities

267 show a discrimination between June and August catabolism, the later showing a higher dif-  
268 ference in catabolism of nitrogen-containing organic substrates (alanine and glycine, glu-  
269 cosamine) (Fig.1-a, % total of variance=73).

270 When considering only June data, no discrimination in the catabolism of both the microbial  
271 and the fungal communities is denoted, either between the cultivars nor between rhi-  
272 zosheath and bulk soils (Fig.2-a; Fig.3-a; Supp. Fig. 2-a; Supp. Fig. 3-a).

273 On the contrary, in August there are differences in catabolic profiles. Microbial catabolic  
274 profiles separate H and T cultivars, with higher catabolism of trehalose and glucose in T  
275 plots, whereas glycine and malate are preferentially catabolized in H plots (Fig. 2-b, 2-c). The  
276 differences in Fungal catabolic profiles observed in August are marked between rhizosheath  
277 and bulk soils (Fig.3 b, c), especially at WD irrigation, where fungal communities preferential-  
278 ly catabolize nitrogen-containing organic substrates (glucosamine, glycine and alanine)  
279 (Fig.3-c).

280 Relation between hydro-physicals, chemical and microbiological indicators

281 Figure 4 presents the relationship between EPSac amounts and the microbial or fungal bio-  
282 masses. EPSac amounts increase with and correlate with microbial biomass ( $R^2= 0.20$ ; p-  
283 value =  $4.09e-5$ ), as well as with fungal biomass ( $R^2= 0.32$ ; p-value =  $1.28e-7$ ).

284 Both for microbial and fungal biomasses, these correlations with EPSac amounts are strong-  
285 er when considering bulk soil and weaker for rhizosheath soil solely (Tab. 3). When consider-  
286 ing the data for June or August, no significant relationship appears (on global, rhizosheath or  
287 bulk soils data), with the exception of a weak relationship for fungal biomass in June ( $R^2=$   
288  $0.22$ ; p-value =  $1.6e-3$ ) (Tab. 3).

289 The best linear models explaining water retention variables (W0.3b and W15b) with the  
290 measured EPSac and microbial/fungal data variables as predictors are shown Table 4 (for  
291 global dataset and for June or August data). Whatever the dataset considered, a significant  
292 correlation emerges between water retention and EPSac/microbial variables ( $R^2$  from 0.16 to  
293 0.74). Most often, the fungal indicators are the significant predictors and EPSac is linked to  
294 W0.3b (except in August-WW irrigation). Fungal catabolism of alanine is also linked to W0.3b  
295 (except in June). The link between water retention (W0.3b and W15b) and EPSac/microbial  
296 variations is stronger in August ( $R^2=0.3$  to 0.74) (and particularly in August-WD irrigation)  
297 compared to June ( $R^2=0.16$  to 0.23).

298 Aboveground productive characteristics and fruit quality measurement

299 There is no significant difference between the two cultivars productive characteristics when  
300 well irrigated (WW) (Fig.5-a). However, under WD these characteristics show systematic  
301 higher values for the T cultivar compared to H (Fig.5-b), with significant differences for CoY,  
302 DMY and WUE (p-value=  $1.9e-2$ ;  $4.8e-2$  and  $1.9e-2$  respectively) except for RoY, for which it  
303 is the opposite (H>T). Titrable acidity and pH of fruits show no significant differences. The  
304 soluble solid content (SSC) is slightly higher for H cultivar compared to T cultivar in context of  
305 deficit irrigation (mean respectively equal to 6.40 and 6.07, in °Brix ; p-value= $3.4e-2$ ).

## 306 Discussion

### 307 Influence of the rhizosheath on soil parameters

308 We highlighted a quantitative effect of the rhizosheath on soil parameters, with higher wa-  
309 ter retention values and higher amounts of exopolysaccharides in the rhizosheath compared  
310 to the bulk soil. We also observed that total microbial and fungal biomass, as well as micro-



311 bial and fungal catabolic activities were significantly higher in the rhizosphere. This differen-  
312 tiation was more pronounced at the fruit maturation stage in August (especially at WD)  
313 compared to June (rhizosphere data: Tab. 2 and S1, interactions between season and soil  
314 factors and also between season and variety factors). A phenology effect seems to occur  
315 here and increases during the season within the rhizosphere (Bardgett et al., 2005). Between  
316 June and August, the rhizosphere soil would have received a fresh carbon supply from the  
317 roots (especially EPSac) of the developed tomato plants, stimulating microbial biomass and  
318 activities and promoting water retention.

319 Qualitatively, these differences were more pronounced for the catabolic profile of fungal  
320 communities at the fruit maturation stage and in water-deficit context (August-WD). Fungal  
321 communities are linked to a higher catabolism of nitrogen containing organic substrates in  
322 rhizosphere soils (Fig.1, Fig.2). We suggest the hypothesis that carbon exudation by roots  
323 boosted by water stress (Henry et al., 2007), could induce a “priming effect” in relation with  
324 the stimulation of microbial and fungal exo-enzymes involved in the degradation of soil or-  
325 ganic nitrogen (Bardgett et al., 2014). Another possible cause would be that the develop-  
326 ment of nitrogen-fixing bacteria was favoured by EPSac secreted by the roots (Nazari, 2021)  
327 in case of water deficit. The contribution of organic nitrogen by diazotrophic bacteria could  
328 then have favoured a higher catabolism of nitrogen containing organic substrates in rhi-  
329 zosphere soils (Zuluaga et al., 2020). To confirm these hypotheses, additional measurements  
330 on the N cycle ( $N_2$  fixing activities, enzymatic activities, nitrogenous forms...) would be nec-  
331 essary.

332 Biochemical activity of roots and their microbial communities. Relationship with  
333 soil water retention

334 Several below ground processes, possibly interacting, can allow the plant to compensate for  
335 water deficit. The particular hydro-physical properties of the rhizosheath are partly linked to  
336 the physical presence of the roots themselves (Aravena et al. 2011, De León-González et al.  
337 2007), and to the production of EPS by the roots and microorganisms (Chenu 1993; Czarnes  
338 et al, 2000; Sher et al., 2020), which are also influenced by changes in moisture and nutrient  
339 conditions (Alami et al. 2000, Henry et al. 2007, Lynch and Whipps 1990, Redmile-Gordon et  
340 al. 2014, Roberson and Firestone 1992).

341 We observe that the amount of soil EPSac follows the same seasonal dynamics as soil micro-  
342 bial and fungal biomass and their catabolic activity, increasing from spring to summer, espe-  
343 cially in rhizosheath soils (Tab. 2, Tab.S1). Indeed, although low, the correlations were posi-  
344 tive between EPSac and microbial and fungal biomasses (the correlation with fungal bio-  
345 masses was higher - Tab. 3 and Fig. 4). In August, at the fruit maturation stage, water reten-  
346 tion at 0.3 and 15 bars was mostly linked to physico-chemical and microbial variables, and  
347 more significantly in case of water-deficit than with full irrigation (Tab. 4). In particular, the  
348 EPSac was partially predictive of the water retention at field capacity (W0.3b) (Tab. 4).

349 The increase in root and microbial biological stimulation in the rhizosheath, notably by a  
350 higher EPS production than their consumption (Redmile-Gordon et al. 2014) could explain  
351 these results. This stimulation in the rhizosheath could have increased with the root density  
352 and the intensification of exudation per unit of root length linked to a phenology effect and  
353 a triggered by water deficit in case of deficit irrigation condition (Henry et al., 2007; Nazari et  
354 al., 2020).

355 Furthermore, fungal biomass and catabolic parameters also seem to be important variables  
356 explaining water retention: the fungal catabolism of alanine, an amino-acid, was almost sys-

357 tematically found as a predictor variable for water retention at field capacity (W0.3b). In  
358 general, fungi are more tolerant to water stress than bacteria (Bérard et al., 2015). Rhizo-  
359 sphere fungi (and mycorrhizae in particular) are known to act on soil aggregation (and water  
360 retention) (Freschet et al., 2021) through the physical action of their hyphae and their pro-  
361 duction of proteins (e.g. glomalin) and polysaccharides (e.g. scleroglucan), resulting in parti-  
362 cle sticking actions (Poirier et al., 2018). Studies involving mycorrhizae and tomatoes show  
363 that mycorrhizae promote plant resistance to drought by stimulating nutrient cycles (includ-  
364 ing nitrogen) and root exudation (especially Amino Acids, which can play an osmoregulatory  
365 role) (Bowles et al. 2016). Moreover, Cavagnaro et al. (2006) showed that mycorrhizae pro-  
366 mote soil aggregation in tomatoes rhizosphere, especially in the presence of mineral N. One  
367 hypothesis would be that tomatoes exude more organic compounds under water deficit  
368 conditions, which would promote nitrogen catabolism (directly and/or through a “priming  
369 effect” or diazotrophic bacteria stimulation as suggested above), fungal development (my-  
370 corrhization) and its direct or indirect effects on water retention. Thus, these different chem-  
371 ical and biological phenomena probably interact with the hydro-physical properties of the  
372 soil (Hallett et al. 2009). However, our hypothesis is based on proxy measurements of bio-  
373 mass and fungal catabolism (FungiResp method). It would have been interesting to measure  
374 the intensity of root colonization by mycorrhizae to confirm this hypothesis, which we were  
375 unable to do during this study and could thus be done in future investigations.

376 Nevertheless, it seems that the fungal catabolism of alanine is related to the hydraulic prop-  
377 erties of the soil without water deficit, while in conditions of water deficit, there could be a  
378 shift to other mechanisms explaining water retention, which would be related to EPSs,   
379 among others. Indeed, in August, we found that the fungal catabolism of alanine explained  
380 the water retention of the soils with higher R<sup>2</sup> values in the WW irrigated plots than in the

381 irrigation-limited plots (WD): in particular for the W0.3b retention at WW irrigation the in-  
382 crease in R2 for F-Alanine was 0.33, while at WD irrigation it dropped to 0.2, with the ap-  
383 pearance of the EPSac as an explanatory variable (increase in R2=0.18; Tab.5).

384 Possible relationships between aboveground cultivars characteristics and their  
385 belowground bio-physico-chemical soil parameters

386 The T cultivar presents a higher production potential compared to the H cultivar in the water  
387 deficit situation, and irrigation seemed to mitigate these differences (Fig. 5). Arbex de Casto  
388 Vilas Boas et al. (2017) showed in a previous similar experiment that WUE was better for T  
389 than H cultivar especially in WD conditions, ranging from 25 kg.m<sup>-3</sup> for WW irrigation to 41  
390 kg.m<sup>-3</sup> for a WD situation in the field. Despite a much higher total water input in this 2018  
391 field experiment and, thus, with lower mean WUE values (between 11 and 18 kg.m<sup>-3</sup>), the  
392 WUE was still significantly higher for T than for H cultivar in WD situation (Fig. 5), indicating  
393 that H seems more sensitive to WD.

394 It seems that our biochemical observations on the soils (microbial activities, especially fungal  
395 activities and amounts of EPSac, Tab.2) leading to a local modification of soil water retention  
396 (Tab.4) are in agreement with the production potential of tomato cultivars T and H, as a  
397 function of their water efficiency (Fig. 5). In fact, cultivar H, which seems to have better ex-  
398 pressed its productive potential under optimal irrigation conditions, presents in its rhi-  
399 zosheath in August-WW a higher F.Ala catabolism, which contributes to explain the water  
400 retentions W0.3b and W15b. On the contrary, cultivar T seems to have better expressed its  
401 production potential under water deficit conditions, in correspondence with higher water  
402 retention W0.3b, together with higher EPSac amounts and higher indicators of fungal activi-  
403 ty and biomass in the T soil plots. Of course, if resistance/tolerance is also related to root

404 system characteristics (root density, root depth) (Fang et Xiong, 2015), our results point to  
405 the fact that biochemical properties of the rhizosheath may also play a role.

406 The water retention properties of soils have a strong influence on crop development and  
407 could factor in the crop's capacity to withstand droughts. A better understanding of the pos-  
408 sible links between soil bio-physicochemical parameters and hydrophysical properties, par-  
409 ticularly in the rhizosheath (Bengough, 2012), could help improve crop resistance to water  
410 deficit. While creating water deficit conditions by limiting irrigation, we measured biophysi-  
411 cal indicators of soil quality and rhizosheath functioning (in comparison to bulk soil). This  
412 field study brings new information regarding the links between soil parameters and produc-  
413 tivity of two tomato cultivars and two types of irrigation.

## 414 Conclusion

415 Our study suggests that rhizosheath affects the soil biophysical parameters measured and  
416 this influence increased between the end of the vegetative phase and the fruit maturation  
417 stage. Secondly, it seems that hydro-physical properties of the soil (e.g., water retention  
418 capacities) are partly linked to two interacting phenomena: the presence in the rhizosheath  
419 environment of specific substances such as exopolysaccharides and/or nitrogenous sub-  
420 stances from roots and microorganisms, and the presence and activity of fungi (possibly my-  
421 corrhizae). The importance of fungal activity stood out during the study and it would be in-  
422 teresting to investigate their effects further by using tools dedicated to fungi and mycorrhi-  
423 za.

424 In addition, these two aforementioned relationships could be a part of the adaptation  
425 mechanisms of the Terradou cultivar to water deficit, enabling higher agricultural productivi-  
426 ty via a higher plant WUE. This study confirms the observations made by Arbex de Castro

427 Vilas Boas et al (2017) and provides hypotheses related to the influence of the rhizosheath  
428 on the responses of these cultivars to water deficit.

429 Our observations—i.e., correlations between the microbial biochemical and hydrophysical  
430 parameters of the soil, which are different for the two cultivars—confirm the importance of  
431 taking into account soil (and particularly rhizosheath soil) parameters when selecting culti-  
432 vars for specific agroclimatic contexts (Nazari et al., 2020, Cattivelli et al., 2008). Finally, this  
433 study, suggests some links between aboveground and soil parameters in relation with water  
434 deficit and plant water use efficiency. This underlines the need of getting a more integrated  
435 and quantitative view of the interactions between soil microbial-biochemical parameters  
436 and soil hydrophysical properties when investigating agro-ecological solutions to water re-  
437 lated yield gap.

## 438 **Acknowledgments**

439 This study was supported by the Structure Fédérative de Recherche Tersys (Avignon,  
440 France). We thank our INRAE colleagues of the unities EMMAH and SQPOV and PSH for their  
441 scientific and technical advices and supports (A.L. Franciullino, T. Clavel, S. Bureau, L.  
442 Capowiez, A. Chapelet, B. Doublet, C. Le Bourvellec, L. Touloumet), as well as the trainees (A.  
443 Davoine, J. Thorel) and members of the UMR EMMAH for their help in sampling campaigns  
444 and the physico-chemical and microbiological characterization of soils. We wish to thank the  
445 Société Nationale Interprofessionnelle de la Tomate (SONITO) (R. Giovinazzo, F. Avril) for  
446 providing their experimental agricultural site and for their support for the operational set up  
447 of the experiment, the monitoring and plant and yield measurements.

## 448 **References**

- 449 Ahmed, M.A., Passioura, J., Carminati, A., 2018. Hydraulic processes in roots and the  
450 rhizosphere pertinent to increasing yield of water-limited grain crops: a critical  
451 review. *Journal of Experimental Botany* 69, 3255–3265.  
452 <https://doi.org/10.1093/jxb/ery183>
- 453 Alami, Y., Achouak, W., Marol, C., Heulin, T., 2000. Rhizosphere Soil Aggregation and Plant  
454 Growth Promotion of Sunflowers by an Exopolysaccharide-Producing Rhizobium sp.  
455 Strain Isolated from Sunflower Roots. *Appl. Environ. Microbiol.* 66, 3393–3398.  
456 <https://doi.org/10.1128/AEM.66.8.3393-3398.2000>
- 457 Anderson, J. P. E., and K. H. Domsch. 1978. "A Physiological Method for the Quantitative  
458 Measurement of Microbial Biomass in Soils." *Soil Biology and Biochemistry* 10 (3):  
459 215–21. [https://doi.org/10.1016/0038-0717\(78\)90099-8](https://doi.org/10.1016/0038-0717(78)90099-8).
- 460 Angers, D.A., Caron, J., 1998. Plant-induced changes in soil structure : Processes and  
461 feedbacks. *Biogeochemistry* 42, 55–72. <https://doi.org/10.1023/A:1005944025343>
- 462 Aravena, J.E., Berli, M., Ghezzehei, T.A., Tyler, S.W., 2011. Effects of Root-Induced  
463 Compaction on Rhizosphere Hydraulic Properties - X-ray Microtomography Imaging  
464 and Numerical Simulations. *Environ. Sci. Technol.* 45, 425–431.  
465 <https://doi.org/10.1021/es102566j>
- 466 Arbex de Castro Vilas Boas, A., Page, D., Giovinazzo, R., Bertin, N., Fanciullino, A.-L., 2017.  
467 Combined Effects of Irrigation Regime, Genotype, and Harvest Stage Determine  
468 Tomato Fruit Quality and Aptitude for Processing into Puree. *Front. Plant Sci.* 8, 1725.  
469 <https://doi.org/10.3389/fpls.2017.01725>
- 470 Bardgett, R.D., Bowman W.D., Kaufmann R. and Schmidt S.K., 2005. A temporal approach to  
471 linking aboveground and belowground ecology. *Trends in Ecology & Evolution* 20,  
472 634-641. <https://doi.org/10.1016/j.tree.2005.08.005>
- 473 Bardgett, R.D., Mommer, L., De Vries, F.T., 2014. Going underground: root traits as drivers of  
474 ecosystem processes. *Trends in Ecology & Evolution* 29, 692–699.  
475 <https://doi.org/10.1016/j.tree.2014.10.006>
- 476 Bengough, A.G., 2012. Water Dynamics of the Root Zone: Rhizosphere Biophysics and Its

477 Control on Soil Hydrology. *Vadose Zone Journal* 11, vzj2011.0111.  
478 <https://doi.org/10.2136/vzj2011.0111>

479 Ben Sassi, M., Dollinger, J., Renault, P., Tlili, A., Bérard, A., 2012. The FungiResp method: An  
480 application of the MicroResp<sup>TM</sup> method to assess fungi in microbial communities as  
481 soil biological indicators. *Ecological Indicators* 23, 482–490.  
482 <https://doi.org/10.1016/j.ecolind.2012.05.002>

483 Benard, Pascal, Mohsen Zarebanadkouki, Mathilde Brax, Robin Kaltenbach, Iwan Jerjen,  
484 Federica Marone, Estelle Couradeau, Vincent J. M. N. L. Felde, Anders Kaestner, and  
485 Andrea Carminati. 2019. “Microhydrological Niches in Soils: How Mucilage and EPS  
486 Alter the Biophysical Properties of the Rhizosphere and Other Biological Hotspots.”  
487 *Vadose Zone Journal* 18 (1): 180211. <https://doi.org/10.3929/ethz-b-000393395>

488 Bérard, A., Bouchet, T., Sévenier, G., Pablo, A.L., Gros, R., 2011. Resilience of soil microbial  
489 communities impacted by severe drought and high temperature in the context of  
490 Mediterranean heat waves. *European Journal of Soil Biology* 47, 333–342.  
491 <https://doi.org/10.1016/j.ejsobi.2011.08.004>

492 Bérard, A., Ben Sassi, M., Kaisermann, A., Renault, P., 2015. Soil microbial community  
493 responses to heat wave components: drought and high temperature. *Climate*  
494 *Research* 66, 243–264. <https://doi.org/10.3354/cr01343>

495 Boeck, H.J.D., Dreesen, F.E., Janssens, I.A., Nijs, I., 2011. Whole-system responses of  
496 experimental plant communities to climate extremes imposed in different seasons.  
497 *New Phytologist* 189, 806–817. <https://doi.org/10.1111/j.1469-8137.2010.03515.x>

498 Bowles, T.M., Barrios-Masias, F.H., Carlisle, E.A., Cavagnaro, T.R., Jackson, L.E., 2016. Effects  
499 of arbuscular mycorrhizae on tomato yield, nutrient uptake, water relations, and soil  
500 carbon dynamics under deficit irrigation in field conditions. *Science of The Total*  
501 *Environment* 566–567, 1223–1234. <https://doi.org/10.1016/j.scitotenv.2016.05.178>

502 Carminati, A., Vetterlein, D., 2013. Plasticity of rhizosphere hydraulic properties as a key for  
503 efficient utilization of scarce resources. *Annals of Botany* 112, 277–290.  
504 <https://doi.org/10.1093/aob/mcs262>

505 Cattivelli, L., Rizza, F., Badeck, F.-W., Mazzucotelli, E., Mastrangelo, A.M., Francia, E., Marè,  
506 C., Tondelli, A., Stanca, A.M., 2008. Drought tolerance improvement in crop plants:



507 An integrated view from breeding to genomics. *Field Crops Research* 105, 1–14.  
508 <https://doi.org/10.1016/j.fcr.2007.07.004>

509 Cavagnaro T.R., Jackson L.E., Six J., Ferris H., Goyal S., Asami D., Scow K.M., 2006. Arbuscular  
510 mycorrhizas, microbial communities, nutrient availability, and soil aggregates in  
511 organic tomato production. *Plant Soil*. 282(1):209–225. doi: 10.1007/s11104-005-  
512 5847-7

513 Chapman, Stephen J., Colin D. Campbell, and Rebekka R. E. Artz. 2007. “Assessing CLPPs  
514 Using MicroResp™.” *Journal of Soils and Sediments* 7 (6): 406–10.  
515 <https://doi.org/10.1065/jss2007.10.259>.

516 Chen, L., Rossi, F., Deng, S., Liu, Y., Wang, G., Adessi, A., De Philippis, R., 2014.  
517 Macromolecular and chemical features of the excreted extracellular polysaccharides  
518 in induced biological soil crusts of different ages. *Soil Biology and Biochemistry* 78, 1–  
519 9. <https://doi.org/10.1016/j.soilbio.2014.07.004>

520 Chenu, C., Roberson, E.B., 1996. Diffusion of glucose in microbial extracellular polysaccharide  
521 as affected by water potential. *Soil Biology and Biochemistry* 28, 877–884.  
522 [https://doi.org/10.1016/0038-0717\(96\)00070-3](https://doi.org/10.1016/0038-0717(96)00070-3)

523 Cornu, P., Valceschini Egizio, Mght-Bournay, O., 2018. Histoire de l’INRA, entre science et po-  
524 litique, Quae. ed. 78026 Versailles Cedex, France

525 Costa, J. Miguel, Maria F. Ortuño, and M. Manuela Chaves. 2007. “Deficit Irrigation as a  
526 Strategy to Save Water: Physiology and Potential Application to Horticulture.” *Journal*  
527 *of Integrative Plant Biology* 49 (10): 1421–34. [https://doi.org/10.1111/j.1672-](https://doi.org/10.1111/j.1672-9072.2007.00556.x)  
528 [9072.2007.00556.x](https://doi.org/10.1111/j.1672-9072.2007.00556.x)

529 Crouzet, O., Consentino, L., Pétraud, J.-P., Marraud, C., Aguer, J.-P., Bureau, S., Le  
530 Bourvellec, C., Touloumet, L., Bérard, A., 2019. Soil Photosynthetic Microbial  
531 Communities Mediate Aggregate Stability: Influence of Cropping Systems and  
532 Herbicide Use in an Agricultural Soil. *Front. Microbiol.* 10, 1319.  
533 <https://doi.org/10.3389/fmicb.2019.01319>

534 Czarnes S., Hallett P. D., Bengough A. G., Young I.M., 2000. Root- and microbial-derived  
535 mucilages affect soil structure and water transport. *European Journal of Soil Science*  
536 51: 435-443.

537 De León-González, F., Gutiérrez-Castorena, M.C., González-Chávez, M.C.A., Castillo-Juárez,  
538 H., 2007. Root-aggregation in a pumiceous sandy soil. *Geoderma* 142, 308–317.  
539 <https://doi.org/10.1016/j.geoderma.2007.08.023>

540 Dormann, C.F., Elith, J., Bacher, S., Buchmann, C., Carl, G., Carré, G., Marquéz, J.R.G., Gruber,  
541 B., Lafourcade, B., Leitão, P.J., Münkemüller, T., McClean, C., Osborne, P.E.,  
542 Reineking, B., Schröder, B., Skidmore, A.K., Zurell, D., Lautenbach, S., 2013.  
543 Collinearity: a review of methods to deal with it and a simulation study evaluating  
544 their performance. *Ecography* 36, 27–46. [https://doi.org/10.1111/j.1600-](https://doi.org/10.1111/j.1600-0587.2012.07348.x)  
545 [0587.2012.07348.x](https://doi.org/10.1111/j.1600-0587.2012.07348.x)

546 Doussan, C., Cousin, I., Berard, A., Chabbi, A., Legendre, L., Czarnes, S., Toussaint, B., Ruy, S.,  
547 2015. Crop systems and plant roots can modify the soil water holding capacity 17,  
548 9285.

549 Duan, H., Wu, J., Huang, G., Zhou, S., Liu, W., Liao, Y., Yang, X., Xiao, Z., Fan, H., 2016.  
550 Individual and interactive effects of drought and heat on leaf physiology of seedlings  
551 in an economically important crop. *AoB PLANTS* plw090.  
552 <https://doi.org/10.1093/aobpla/plw090>

553 Dubois, M., Gilles, K.A., Hamilton, J.K., Rebers, P.A., Smith, F., 1956. Colorimetric Method for  
554 Determination of Sugars and Related Substances | *Analytical Chemistry* [WWW  
555 Document]. URL <https://pubs.acs.org/doi/pdf/10.1021/ac60111a017> (accessed  
556 8.18.20).

557 FAOSTAT [WWW Document], n.d. URL <http://www.fao.org/faostat/en/#data> (accessed  
558 5.11.20).

559 Fang, Y., Xiong, L. General mechanisms of drought response and their application in drought  
560 resistance improvement in plants. *Cell. Mol. Life Sci.* 72, 673–689 (2015).  
561 <https://doi.org/10.1007/s00018-014-1767-0>

562 Freschet, G. T., Roumet, C., Comas, L. H., Weemstra, M., Bengough, A. G., Rewald, B.,  
563 Bardgett, R. D., De Deyn, G., Johnson, D., Klimešová, J., Lukac, M., McCormack, M. L.,  
564 Meier, I. C., Pagès, L., Poorter, H., Prieto, I., Wurzbürger, N., Zadworny, M.,  
565 Bagniewska-Zadworna, A., ... Stokes, A., 2021. Root traits as drivers of plant and  
566 ecosystem function-ing: Current understanding, pitfalls and future research needs.

- 567 New Phytologist. In press
- 568 Göttlein, A., 2006. Sampling of rhizosphere soil and collection of rhizosphere soil solution. In:  
569 Luster, J., Finlay, R. (Eds.), Handbook of Methods Used in Rhizosphere Research.  
570 Swiss Federal Research Institute WSL, Birmensdorf, pp. 25–29.
- 571 Hallett, P.D., Feeney, D.S., Bengough, A.G., Rillig, M.C., Scrimgeour, C.M., Young, I.M., 2009.  
572 Disentangling the impact of AM fungi versus roots on soil structure and water  
573 transport. *Plant Soil* 314, 183–196. <https://doi.org/10.1007/s11104-008-9717-y>
- 574 Hegyi, G., Garamszegi, L.Z., 2011. Using information theory as a substitute for stepwise  
575 regression in ecology and behavior. *Behav Ecol Sociobiol* 65, 69–76.  
576 <https://doi.org/10.1007/s00265-010-1036-7>
- 577 Henry A., Doucette W., Norton J., Bugbee B., 2007. Changes in Crested Wheatgrass Root  
578 Exudation Caused by Flood, Drought, and Nutrient Stress. *Journal of Environmental*  
579 *Quality* 36, 904-912.
- 580 Hinsinger, P., Bengough, A.G., Vetterlein, D. et al. Rhizosphere: biophysics, biogeochemistry  
581 and ecological relevance. *Plant Soil* 321, 117–152 (2009).  
582 <https://doi.org/10.1007/s11104-008-9885-9>
- 583 Jia, G., E. Shevliakova, P. Artaxo, N. De Noblet-Ducoudré, R. Houghton, J. House, K. Kitajima,  
584 C. Lennard, A. Popp, A. Sirin, R. Sukumar, L. Verchot (2019). Land–climate  
585 interactions. In: *Climate Change and Land: an IPCC special report on climate change,*  
586 *desertification, land degradation, sustainable land management, food security, and*  
587 *greenhouse gas fluxes in terrestrial ecosystems* [P.R. Shukla, J. Skea, E. Calvo Buendia,  
588 V. Masson-Delmotte, H.-O. Pörtner, D.C. Roberts, P. Zhai, R. Slade, S. Connors, R. van  
589 Diemen, M. Ferrat, E. Haughey, S. Luz, S. Neogi, M. Pathak, J. Petzold, J. Portugal  
590 Pereira, P. Vyas, E. Huntley, K. Kissick, M, Belkacemi, J. Malley, (eds.)]. In press.
- 591 Łabędzki L., 2016, Actions and measures for mitigation drought and water scarcity in  
592 agriculture. *Journal of Water and Land Development* 29, 3-10.
- 593 Lê, S., Josse, J. & Husson, F. (2008). FactoMineR: An R Package for Multivariate Analysis.  
594 *Journal of Statistical Software*. 25(1). pp. 1-18.
- 595 Lipiec, J., Doussan, C., Nosalewicz, A., Kondracka, K., 2013. Effect of drought and heat

596 stresses on plant growth and yield: a review. *International Agrophysics* 27, 463–477.  
597 <https://doi.org/10.2478/intag-2013-0017>Lynch, J.M., Whipps, J.M., 1990. Substrate  
598 flow in the rhizosphere. *Plant Soil* 129, 1–10. <https://doi.org/10.1007/BF00011685>

599 Marasco, R., Mosqueira, M.J., Fusi, M. et al., 2018. Rhizosphere microbial community  
600 assembly of sympatric desert speargrasses is independent of the plant host.  
601 *Microbiome* 6, 215. <https://doi.org/10.1186/s40168-018-0597-y>

602 Murtaugh, P.A., 2009. Performance of several variable-selection methods applied to real  
603 ecological data. *Ecology Letters* 12, 1061–1068. <https://doi.org/10.1111/j.1461-0248.2009.01361.x>

605 Nazari M., 2021. Plant mucilage components and their functions in the rhizosphere.  
606 *Rhizosphere* 18, 100344. <https://doi.org/10.1016/j.rhisph.2021.100344>

607 Nazari M, Riebeling S, Banfield CC, Akale A, Crosta M, Mason-Jones K, Dippold MA and  
608 Ahmed MA, 2020. Mucilage Polysaccharide Composition and Exudation in Maize  
609 From Contrasting Climatic Regions. *Front. Plant Sci.* 11:587610. [https://doi:](https://doi:10.3389/fpls.2020.587610)  
610 [10.3389/fpls.2020.587610](https://doi:10.3389/fpls.2020.587610)

611 Pang, J., Ryan, M.H., Siddique, K.H.M. et al., 2017. Unwrapping the rhizosphere. *Plant Soil*  
612 418, 129–139. <https://doi.org/10.1007/s11104-017-3358-y>

613 Patanè, C., Cosentino, S.L., 2010. Effects of soil water deficit on yield and quality of  
614 processing tomato under a Mediterranean climate. *Agricultural Water Management*  
615 97, 131–138. <https://doi.org/10.1016/j.agwat.2009.08.021>

616 Personeni E., Nguyen C., Marchal P., Pagès L., 2007. Experimental evaluation of an efflux–  
617 influx model of C exudation by individual apical root segments. *Journal of*  
618 *Experimental Botany*, 58, 8: 2091–2099. doi:10.1093/jxb/erm065

619 Poirier V, Roumet C, Munson AD. 2018. The root of the matter: linking root traits and soil  
620 organic matter stabilization processes. *Soil Biology and Biochemistry* 120: 246–259.

621 Rasse, D.P., Smucker, A.J.M., Santos, D., 2000. Alfalfa Root and Shoot Mulching Effects on  
622 Soil Hydraulic Properties and Aggregation. *Soil Sci. Soc. Am. J.* 64, 725–731.  
623 <https://doi.org/10.2136/sssaj2000.642725x>

624 R Core Team (2017). R: A language and environment for statistical computing. R Foundation

625 for Statistical Computing, Vienna, Austria. URL <https://www.R-project.org/>.

626 Redmile-Gordon, M.A., Brookes, P.C., Evershed, R.P., Goulding, K.W.T., Hirsch, P.R., 2014.  
627 Measuring the soil-microbial interface: Extraction of extracellular polymeric  
628 substances (EPS) from soil biofilms. *Soil Biology and Biochemistry* 72, 163–171.  
629 <https://doi.org/10.1016/j.soilbio.2014.01.025>

630 Rinaldi, M., Ventrella, D., Gagliano, C., 2007. Comparison of nitrogen and irrigation strategies  
631 in tomato using CROPGRO model. A case study from Southern Italy. *Agricultural*  
632 *Water Management* 87, 91–105. <https://doi.org/10.1016/j.agwat.2006.06.006>

633 Ripoll, J., Urban, L., Brunel, B., Bertin, N., 2015. Water deficit effects on tomato quality  
634 depend on fruit developmental stage and genotype. *Journal of plant physiology* 190,  
635 26-35. <https://doi.org/10.1016/j.jplph.2015.10.006>

636 Roberson, E.B., Firestone, M.K., 1992. Relationship between Desiccation and  
637 Exopolysaccharide Production in a Soil *Pseudomonas* sp. *Appl. Environ. Microbiol.* 58,  
638 1284–1291.

639 RStudio Team (2020). RStudio: Integrated Development for R. RStudio, PBC, Boston, MA URL  
640 <http://www.rstudio.com/>.

641 Schielzeth, H., 2010. Simple means to improve the interpretability of regression coefficients.  
642 *Methods in Ecology and Evolution* 1, 103–113. [https://doi.org/10.1111/j.2041-](https://doi.org/10.1111/j.2041-210X.2010.00012.x)  
643 [210X.2010.00012.x](https://doi.org/10.1111/j.2041-210X.2010.00012.x)

644 Sher, Y., Baker, N.R., Herman, D., Fossum, C., Hale, L., Zhang, X.-X., Nuccio, E., Saha, M.,  
645 Zhou, J., Pett-Ridge, J., Firestone, M., 2019. Microbial extracellular polysaccharide  
646 production and aggregate stability controlled by Switchgrass (*Panicum virgatum*) root  
647 biomass and soil water potential. *bioRxiv* 724195. <https://doi.org/10.1101/724195>

648 van Veelen, A., Tourell, M.C., Koebernick, N., Pileio, G., Roose, T., 2018. Correlative  
649 Visualization of Root Mucilage Degradation Using X-ray CT and MRI. *Front. Environ.*  
650 *Sci.* 6. <https://doi.org/10.3389/fenvs.2018.00032>

651 Whittingham, M.J., Stephens, P.A., Bradbury, R.B., Freckleton, R.P., 2006. Why do we still use  
652 stepwise modelling in ecology and behaviour? *Journal of Animal Ecology* 75, 1182–  
653 1189. <https://doi.org/10.1111/j.1365-2656.2006.01141.x>

- 654 Zhang, Xuechen, Yakov Kuzyakov, Huadong Zang, Michaela A. Dippold, Lingling Shi, Sandra  
655 Spielvogel, and Bahar S. Razavi. 2020. "Rhizosphere Hotspots: Root Hairs and  
656 Warming Control Microbial Efficiency, Carbon Utilization and Energy Production." *Soil  
657 Biology and Biochemistry* 148 (September): 107872.  
658 <https://doi.org/10.1016/j.soilbio.2020.107872>
- 659 Zuluaga M.Y.A., Lima Milani K.M., Azeredo Gonçalves L.S., Martinez de Oliveira A.L., 2020.  
660 Diversity and plant growth-promoting functions of diazotrophic/N-scavenging  
661 bacteria isolated from the soils and rhizospheres of two species of *Solanum*. *PLoS  
662 ONE* 15(1): e0227422. <https://doi.org/10.1371/journal.pone.0227422>

Figures and table

## Table of abbreviations

Abbreviation	Description	Unit
<b>A</b>	Soil sampling date at the beginning of fructification stage (August 2. 2018).	-
<b>AIC</b>	Akaike Information Criterion	-
<b>B (Bulk)</b>	Soil sampling from bulk soil (without roots)	-
<b>CoY</b>	Fresh Commercial Tomato Yield (including red and yellow tomato)	kg/ha
<b>DMY</b>	Commercial Tomato Yield in Dry Matter (including red and yellow tomato)	kg/ha
<b>EPSac</b>	ExoPolySaccharides	mg eq glc / g DrySoil
<b>F.Ala</b>	Alanine induced respiration for fungi communities in soil	CO <sub>2</sub> /gDrySoil/h
<b>F.Bas</b>	Basal respiration for fungi communities in soil	CO <sub>2</sub> /gDrySoil/h
<b>F.Cell</b>	Cellulobiose induced respiration for fungi communities in soil	CO <sub>2</sub> /gDrySoil/h
<b>F.Gly</b>	Glycine induced respiration for fungi communities in soil	CO <sub>2</sub> /gDrySoil/h
<b>F.GlcN</b>	Glucosamine induced respiration for fungi communities in soil	CO <sub>2</sub> /gDrySoil/h
<b>F.Mal</b>	Malate induced respiration for fungi communities in soil	CO <sub>2</sub> /gDrySoil/h
<b>F.Tre</b>	Trehalose induced respiration for fungi communities in soil	CO <sub>2</sub> /gDrySoil/h
<b>H</b>	Soil sampling from Heinz 1015 tomato cultivar	-
<b>J</b>	Soil sampling date at the beginning of flowering stage (June 19. 2018).	-
<b>M.Ala</b>	Alanine induced respiration for fungi communities in soil	CO <sub>2</sub> /gDrySoil/h
<b>M.Bas</b>	Basal respiration for fungi communities in soil	CO <sub>2</sub> /gDrySoil/h
<b>M.Cell</b>	Cellulobiose induced respiration for fungi communities in soil	CO <sub>2</sub> /gDrySoil/h
<b>M.Gly</b>	Glycine induced respiration for fungi communities in soil	CO <sub>2</sub> /gDrySoil/h
<b>M.GlcN</b>	Glucosamine induced respiration for fungi communities in soil	CO <sub>2</sub> /gDrySoil/h
<b>M.Mal</b>	Malate induced respiration for fungi communities in soil	CO <sub>2</sub> /gDrySoil/h
<b>M.Tre</b>	Trehalose induced respiration for fungi communities in soil	CO <sub>2</sub> /gDrySoil/h
<b>R (Rhizo)</b>	Soil sampling from rhizosheath soil (root adhering)	-
<b>RoY</b>	Rotten Tomato Yield	t/ha
<b>T</b>	Soil sampling from Terradou tomato cultivar	-
<b>VIF</b>	Variable Inflation Factor	-
<b>W0.3b</b>	Water Retention Capacity at 0.3b (at Field Capacity)	g/g
<b>W15b</b>	Water Retention Capacity at 15b (at Permanent Wilting Point)	g/g
<b>WD</b>	Water deficit irrigation (57% of ETP losses)	-
<b>WUE</b>	Plant Water Use Efficiency	kg/m <sup>3</sup>
<b>WW</b>	Well watered irrigation (87% of ETP losses)	-



Localisation	44°11'N/4°48'E
Texture	loam
Clay (g kg <sup>-1</sup> )	290
Loam (g kg <sup>-1</sup> )	331
Sand (g kg <sup>-1</sup> )	379
SOC (g kg <sup>-1</sup> )	12.5
TN (g kg <sup>-1</sup> )	1.1
C/N	11
CaCO <sub>3</sub> (g kg <sup>-1</sup> )	484
Ca <sup>2+</sup> (cmol+ kg <sup>-1</sup> )	14.6
Mg <sup>2+</sup> (cmol+ kg <sup>-1</sup> )	0.595
CEC (cmol+ kg <sup>-1</sup> )	12.7
pH (water)	8.5
WSA (%)	36

Table 1: Main physical, chemical and microbial characteristics of the soils used for this study on the Piolenc experimental site. SOC: Soil Organic Carbon, TN: Total Nitrogen, C/N: carbon over nitrogen ratio CEC: Cation Exchange Capacity, WSA: Water Stable Aggregates

	W0.3b	W15b	EPSac	F.Glc	F.Tre	F.Cell	F.GlcN	F.Ala	F.Gly	F.Mal	FB	M.Bas	M.Glc	M.Tre	M.Cell	M.GlcN	M.Ala	M.Gly	M.Mal	MB
Global																				
SOIL	R>B	R>B***	R>B	R>B***	R>B**	R>B***	R>B***	R>B***	R>B***	R>B***	R>B***	R>B**	R>B***	R>B***	R>B***	R>B***	R>B***	R>B***	R>B***	R>B***
CULT	T>H	H>T	T>H	T>H	T>H	T>H	T>H	H>T	T>H	H>T	T>H**	T>H	T>H*	T>H**	T>H*	T>H	T>H	T>H	H>T	T>H*
SEAS	J>A	A>J	A>J***	A>J***	A>J***	A>J***	A>J***	A>J***	A>J***	A>J***	A>J***	A>J***	A>J***	A>J***	A>J***	A>J***	A>J***	A>J***	A>J***	A>J***
SOIL:CULT	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
SOIL:SEAS	***	-	-	-	-	-	***	*	-	-	-	-	-	-	*	-	-	-	-	-
CULT:SEAS	-	*	**	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	*	-
Global Rhizo																				
CULT	T>H	H>T	T>H	T>H***	T>H	T>H	T>H	H>T	H>T	T>H	T>H**	T>H	T>H	T>H*	T>H	T>H	T>H	H>T	H>T	T>H
SEAS	J>A***	A>J**	A>J***	A>J***	A>J***	A>J***	A>J***	A>J***	A>J***	A>J***	A>J***	A>J*	A>J***	A>J***	A>J***	A>J***	A>J***	A>J***	A>J***	A>J***
CULT:SEAS	-	*	-	-	-	-	-	*	-	-	-	-	-	-	-	-	-	*	**	-
June																				
SOIL	R>B***	R>B	R>B**	R>B***	R>B	R>B*	R>B	R>B	R>B	R>B*	R>B***	R>B	R>B	R>B**	R>B*	R>B	R>B*	R>B*	R>B**	R>B
CULT	H>T	T>H	H>T	T>H	T>H	T>H	T>H	T>H	T>H	T>H	T>H	T>H	T>H	T>H	T>H	T>H	T>H	T>H	T>H	T>H
SOIL:CULT	-	-	-	*	-	-	-	-	-	-	**	-	-	-	-	-	-	-	-	-
June Rhizo																				
CULT	H>T	T>H	T>H	T>H*	T>H	T>H	T>H	T>H	T>H	T>H	T>H*	T>H	T>H	T>H	T>H	T>H	T>H	T>H	T>H	T>H
August WW																				
SOIL	B>R	R>B*	R>B	R>B**	R>B*	R>B*	R>B***	R>B**	R>B**	R>B*	R>B**	R>B*	R>B***	R>B***	R>B***	R>B***	R>B**	R>B***	R>B***	R>B***
CULT	T>H	H>T*	T>H*	T>H	T>H	H>T	T>H	H>T	H>T	H>T	T>H	T>H	T>H	T>H	T>H	T>H	H>T	H>T	H>T	T>H
SOIL:CULT	-	-	-	-	-	-	-	*	-	-	-	-	-	-	*	-	**	*	-	-
August WW Rhizo																				
CULT	T>H	H>T	T>H	T>H	H>T	H>T	T>H	H>T*	H>T	H>T	H>T	H>T	H>T	H>T	T>H	H>T	H>T	H>T	H>T**	H>T**
August WD																				
SOIL	B>R**	R>B***	R>B	R>B	R>B	R>B	R>B***	R>B***	R>B***	R>B*	R>B	R>B*	R>B***	R>B***	R>B***	R>B***	R>B***	R>B***	R>B***	R>B***
CULT	T>H***	T>H	T>H	T>H*	T>H	T>H	T>H*	H>T	H>T	T>H	T>H*	T>H	T>H	T>H	T>H	T>H	T>H	T>H	H>T	T>H
SOIL:CULT	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
August WD Rhizo																				
CULT	T>H*	H>T	T>H	T>H*	T>H	T>H	T>H	T>H	H>T	T>H	T>H*	T>H	T>H	T>H	T>H	T>H	T>H	H>T	H>T	H>T

Table 2: Analyses of variance (ANOVAs) results for the soil parameters for the three factors : the type of soil rhizosphere or bulk (SOIL), the season (SEAS) and the cultivar (CULT). Main effect and their interactions are calculated on global dataset (whatever the rhizosphere or bulk soil, the season, the cultivar) and on rhizosphere soil dataset. Below are analysis for June, August-WD irrigation and August-WW irrigation dataset. X>Y signifies that the parameter X is greater than Y. Level of significance: \*p<0.05; \*\*p<0.01; \*\*\*p<0.001. See table of abbreviations for more information.


<b>Dataset</b>	<b>Factor</b>	<b>R<sup>2</sup></b>	<b>p-value</b>	<b>Sign. code</b>
<b>Global</b>	MB	0.20	4.09E-05	***
n=76	FB	0.32	1.28E-07	***
<b>Rhizo</b>	MB	0.16	7.39E-03	**
n=38	FB	0.19	3.53E-03	**
<b>Bulk</b>	MB	0.21	2.38E-03	**
n=38	FB	0.39	1.71E-05	***
<b>June</b>	MB	-0.02	6.12E-01	
n=38	FB	0.22	1.60E-03	**
<b>June rhizo</b>	MB	-0.06	9.63E-01	
n=19	FB	0.11	9.07E-02	.
<b>June bulk</b>	MB	-0.03	4.79E-01	
n=19	FB	5E-4	3.29E-01	
<b>August</b>	MB	5E-4	3.19E-01	
n=38	FB	-0.03	7.83E-01	
<b>August rhizo</b>	MB	-0.06	8.48E-01	
n=19	FB	-0.03	5.05E-01	
<b>August bulk</b>	MB	0.07	1.50E-01	
n=19	FB	-0.005	3.54E-01	

Table 3: Results of linear regression between EPSac and the microbial or fungal biomasses (MB, FB). Considering global soil data or subsets of the soil data. a: slope ; b: intercept of linear regression. Level of significance: \*p<0,05; \*\*p<0,01;\*\*\*p<0,001. See table of abbreviations for more information. See table of abbreviations for more information.


Variables selected	AIC	VIF	Final AIC	Final p-value	Signif. Code	Final R <sup>2</sup>
<b>W0.3b global</b>			-445.06	3.69E-04	***	0.21
F.Ala	-433.78	5				
M.Ala	-441.62	3.06				
F.Cell	-444.49	5.43				
EPSac	-445.06	1.73				
<b>W15b global</b>			-562.14	6.14E-05	***	0.21
F.Tré	-548.36	3.67				
M.Tré	-550.61	3.67				
<b>W0.3b June</b>			-215.13	7.91E-03	**	0.23
EPSac	-212.62	1.13				
F.Cell	-212.71	5.09				
F.Tré	-215.13	4.87				
<b>W15b June</b>			-284.35	8.24E-03	**	0.16
M.Cell	-284.35	nn				
<b>W0.3b August-WW</b>			-122.85	4.15E-03	**	0.51
FB	-112.45	1.66				
M.Cell	-114.28	2.73				
F.Ala	-122.85	1.86				
<b>W15b August-WW</b>			-120.62	1.04E-02	*	0.3
F.Ala	-120.62	nn				
<b>W0.3b August-WD</b>			-129.78	3.94E-04	***	0.65
F.Ala	-115.33	2.61				
EPSac	-119.84	1.07				
M.Tré	-123.31	1.98				
F.Gly	-129.78	2.99				
<b>W15b August-WS</b>			-164.75	1.50E-05	***	0.74
F.D.Glc	-157.97	5.2				
M.Gly	-161.08	2.94				
M.D.Glc	-164.75	7.1				

Table 4: Multiple linear regression model between water retention (W0.3b and W15b) and both physico-chemical (EPSac) and microbial parameters (catabolic profile, biomass) for the global dataset and subsets. Best model issued from stepwise regression using AIC selection criteria. The final model include variable with VIF<10. restricted to a maximum 4 explanatory variables. Level of significance: \*p<0.05; \*\*p<0.01;\*\*\*p<0.001. See table of abbreviations for more information.


1-a: Season



1-c: Soil type



1-b: Cultivar



1-d: Correlation graph substrates




Figure 1: Principal Component Analysis (PCA) of fungal catabolic profile of the rhizosphere and bulk soil data. Projection on axes 1 and 2 (73% of variance). Distribution of individuals and correlations shown for cultivar (H and T), soil type (R and B) and sampling season (J and A).




Figure 2: Principal Component Analysis (PCA) of microbial catabolic profile of the rhizosphere and bulk soil data. Projection on axes 1 and 2 (> 64% variance). Microbial catabolic profile for June, August-WW and August-WD data. Distribution of individuals and correlations shown for cultivar (H and T).




Figure 3: Principal Component Analysis (PCA) of fungal catabolic profile for June, August-WW and August-WD data. Projection on axes 1 and 2 (> 65% variance). Distribution of individuals and correlations shown for soil types (R and B).




Fig. 4: Linear regression between EPSac amount and the microbial or fungal biomasses (MB, FB) for the global soil data set.






Figure 5: Box plots comparing H and T varieties for the WW irrigation dataset (Fig.5-a), and the WD irrigation dataset (Fig.5-b), with Rotten tomato Yield (RoY, in t ha-1), fresh Commercial Yield (CoY, in t ha-1) and Dry Matter Yield (DMY, in 10 kg ha-1) and Water Use Efficiency (WUE, in kg m-3). Level of significance: \* $p < 0.05$ . See table of abbreviations for more information.