



**HAL**  
open science

## Effects of Lacaune ewes selection for somatic cells score and milk persistency on rumen bacteria

Guillermo Martinez-Boggio, Annabelle Meynadier, Rachel Rupp, H el ene Larroque, Charlotte Allain, Christel Marie-Etancelin

### ► To cite this version:

Guillermo Martinez-Boggio, Annabelle Meynadier, Rachel Rupp, H el ene Larroque, Charlotte Allain, et al.. Effects of Lacaune ewes selection for somatic cells score and milk persistency on rumen bacteria. EAAP 2020, Aug 2020, porto, Portugal. hal-03480856

**HAL Id: hal-03480856**

**<https://hal.inrae.fr/hal-03480856>**

Submitted on 14 Dec 2021

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destin ee au d ep ot et  a la diffusion de documents scientifiques de niveau recherche, publi es ou non,  emanant des  tablissements d'enseignement et de recherche fran ais ou  trangers, des laboratoires publics ou priv es.

## Effects of Lacaune ewes selection for somatic cells score and milk persistency on rumen bacteria

Guillermo Martinez-Boggio<sup>1\*</sup>, Annabelle Meynadier<sup>1</sup>, Rachel Rupp<sup>1</sup>, H el ene Larroque<sup>1</sup>, Charlotte Allain<sup>2</sup>, Christel Marie-Etancelin<sup>1</sup>

\*[guillermo.martinez-boggio@inrae.fr](mailto:guillermo.martinez-boggio@inrae.fr)

<sup>1</sup>GenPhySE, Universit e de Toulouse, INRAE, ENVT, F-31326 Castanet Tolosan, France

<sup>2</sup>INRAE, Experimental Unit of La Fage, F-12250 Saint-Jean et Saint-Paul, France

The ruminal microbiota plays a central role in the nutrition of its host, directly affecting milk production and animal health. There are no such studies in sheep that describes this link, thus the aim of our study was to determine differences in ruminal bacteria between divergent lines for somatic cells score and milk production persistency of Lacaune ewes.

We had rumen fluid samples of 700 dairy Lacaune ewes raised at the INRAE Experimental Farm of La Fage sampling between 2015 and 2019. These ewes were adult animals, mostly in their second lactation, and belonged to two divergent lines selected for somatic cells score (SCS) or milk production persistency during lactation (PERS). The phenotypes of ewes obtained from milk recording controls over the five years were somatic cells count, milk production, milk fat content, milk protein content and milk dry matter content. Ruminal metagenome was sequenced using 16s rRNA gene with Illumina Miseq technology. We analyzed microbiota sequences with FROGS pipeline to obtain relative abundances of OTUs (in proportion of total sequences per sample) and R Phyloseq package to estimate biodiversity indices. Thereafter we grouped OTUs according to phylum, family and genus levels.

Fixed effects would be included in the variance analyses were the lines (SCC+, SCC-, PERS+, PERS-), run sequencing effect, date of rumen fluid sampling, lactation number (2 up to 7 lactations), stage of lactation (45 to 133 DIM) and litter size (1, 2 or more lambs), moreover we tested also suckling method of the ewe (maternal or artificial). ANOVA of  $\alpha$ -diversity and relative abundance of OTUs or taxa at different taxonomic levels were estimated by including the significant effects. Further, a discriminant analysis (with sPLS-DA) was applied on OTUs with divergent lines as discriminant factor, and some links were researched between phenotypes and OTUs (with sPLS), using MixOmics R-package.

With these analyses, we found several links between ewes' performances and rumen bacteria and we characterized the ruminal microbiota of divergent lines.