



HAL
open science

**Phenotypic and genotypic data of a European beech
(*Fagus sylvatica* L.) progeny trial issued from three plots
along an elevation gradient in Mont Ventoux,
South-Eastern France**

Sylvie Oddou-Muratorio, Julie Gauzere, Nicolas Angeli, Patrice Brahic, Oliver
Brendel, Marie de Castro, Olivier Gilg, Christian Hossann, Frédéric Jean,
Matthieu Lingrand, et al.

► **To cite this version:**

Sylvie Oddou-Muratorio, Julie Gauzere, Nicolas Angeli, Patrice Brahic, Oliver Brendel, et al.. Phenotypic and genotypic data of a European beech (*Fagus sylvatica* L.) progeny trial issued from three plots along an elevation gradient in Mont Ventoux, South-Eastern France. *Annals of Forest Science*, 2021, 78, pp.88. 10.1007/s13595-021-01105-9 . hal-03483916

HAL Id: hal-03483916

<https://hal.inrae.fr/hal-03483916v1>

Submitted on 16 Dec 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License



Phenotypic and genotypic data of a European beech (*Fagus sylvatica* L.) progeny trial issued from three plots along an elevation gradient in Mont Ventoux, South-Eastern France

Sylvie Oddou-Muratorio^{1,2} · Julie Gauzere^{1,3} · Nicolas Angeli⁴ · Patrice Brahic⁵ · Oliver Brendel⁶ · Marie De Castro⁵ · Olivier Gilg⁷ · Christian Hossann⁴ · Frédéric Jean^{1,7} · Matthieu Lingrand¹ · Mehdi Pringarbe⁷ · Frank Rei⁷ · Anne Roig¹ · Jean Thevenet⁷ · Norbert Turion⁷

Received: 8 March 2021 / Accepted: 13 September 2021
© The Author(s) 2021

Key Message

We provide phenotypic and genotypic data for a progeny trial of 5813 European beech seedlings, originating from 60 open-pollinated families collected at three altitudes (1020 m; 1140 m, 1340 m) on Mont Ventoux (44° 11' N; 17° 5' E).

Keywords Growth traits · Functional traits · Seedlings · Half-sib trial · Heritability · Paternity analyses

1 Background

Considering the patterns of adaptive traits' genetic divergence and local adaptation displayed by many tree species at large spatial scale, forest tree populations are usually assumed to have a high evolutionary potential (Alberto et al.

2013). However, there is still limited evidence of the level of genetic variation available within population at key functional traits involved in response to climate. Moreover, we also need to investigate the abilities of tree populations to adapt to local variation of their environment (i.e., microgeographic adaptation, Richardson et al. 2014).

This data paper extensively describes a valuable quantitative genetic experiment designed to address these issues in the European beech (*Fagus sylvatica* L.), a major tree species in Europe. Sixty beech maternal progenies were collected in three plots along an elevation gradient and grown in a common garden under two contrasted experimental conditions (water stress/no water stress), to assess how the variation at twelve adaptive traits partitioned within and among families, plots, and experimental contrasts. Moreover, we genotyped a subset of offspring and all the potentially reproductive adults in the three plots at 13 microsatellite markers to infer paternal relationships and to estimate average relatedness within and between maternal families and genetic divergence among plots.

Handling Editor: Marianne Peiffer

Contribution of all co-authors S. Oddou-Muratorio: Conceptualization, investigation, formal analysis, writing-original draft, supervision, funding acquisition. J. Gauzere: Conceptualization, investigation, formal analysis, data curation, writing-review and editing. N. Angeli: Investigation, resources. P. Brahic: Investigation, resources. O. Brendel: Investigation, formal analysis, validation, writing-review and editing. M. De Castro: Investigation. O. Gilg: Methodology, investigation. C. Hossann: Investigation. F. Jean: Methodology, investigation, writing-review and editing. M. Lingrand: Investigation, validation. M. Pringarbe: Investigation. F. Rei: Investigation. A. Roig: Investigation, validation. J. Thevenet: Methodology, investigation. N. Turion: Investigation

Dataset access is at: <https://doi.org/10.15454/6HETQP>. Associated metadata are available at: <https://metadata-afs.nancy.inra.fr/geonetwork/srv/fre/catalog.search#/metadata/55aa3d92-3866-48e9-b000-f9cbf2d13d03>

✉ Sylvie Oddou-Muratorio
sylvie.muratorio@inrae.fr

Extended author information available on the last page of the article

2 Methods

2.1 Species and field sampling

The European beech is a shade-tolerant species requiring well-drained, moderately deep soils, and relatively high humidity. Its distribution ranges from the northern

Mediterranean regions to the south of Scandinavia. On Mont-Ventoux, Southeast of France (44° 11' N; 17° 5' E), beech forests are at the climatic limit of their ecological range (Fig. 1). On the North slope of Mont-Ventoux, beech forest ranges almost continuously from 750 to 1700 m in elevation. This steep elevation gradient provides almost linear variation in mean temperature and humidity with elevation (Fig. 1). Along this elevation gradient, we selected three plots at elevations N1: 1020 m (dimension: 1.30 ha), N2: 1140 m (2.20 ha), and N4: 1340 m (0.80 ha). These plots extend over 1.5 km (Fig. 2).

Within each plot, we achieved the exhaustive mapping and genotyping of the reproductive beeches (Fig. 2; Bontemps et al. 2013; Gauzere et al. 2013a; Oddou-Muratorio et al. 2018). This sampling design is relevant to perform paternity analysis, as we know that most mating events occur within the maternal neighborhood (average pollen dispersal distance $d = [35; 63]$ m; Gauzere et al. 2013a).

The progeny trial described in this study includes 60 open-pollinated families (numbered with a Family_ID ranging from 1 to 60). In August 2009, 20 highly fertile and randomly distributed trees were chosen as mother-trees in each plot (Oddou-Muratorio and Gauzere 2021). We collected ~ 1164 seeds per mother-tree (min = 725, max = 1655) directly from the canopy (either by climbing or using scissors mounted on a rod). A sample of 344 seeds per mother-tree on average (min = 202, max = 733) was randomly chosen to measure the average seed weight (g), and the proportions of empty, infested, and healthy seeds based on Faxitron numerical X-ray radiography (Faxitron Bioptics, Tucson, AZ; 15–20 kV, 0.3–3 mA). All the seeds were dried to a humidity rate of 8%.

2.2 Half-sib trial establishment

In October 2009, seeds were rehydrated and conserved at + 4 °C during 10 weeks to break dormancy and initiate germination. In April 2010, 8976 seedlings successfully germinated, among which 5854 were kept as “focal” seedlings (91 seedlings per family on average). All the 8976 seedlings were transferred in a common garden at the nursery of Aix-Les-Milles (43° 30' N; 5° 24' E) where they were planted in individual pots of 1.2 l with sand substrate and fertilizer. Seedlings were grouped in boxes of 17 seedlings and stored over 6 shelves (Fig. 3). The 5854 focal seedlings were arranged in 50 complete blocks, each block including ~ 2 seedlings per family (Fig. 3). To ensure that the 5854 focal seedlings had neighboring seedlings (i.e., to avoid border effects), we used 3122 additional seedlings as “border” (i.e., on external sides of boxes) or “neutral” (i.e., within boxes) seedlings (Fig. 3; see also the “Trial Design” spreadsheet in the database). Among the 5854 focal

seedlings, 41 failed to survive to plantation, leading to 5813 effectively surveyed seedlings.

The seedlings grew 3 years (from April 2010 to September 2013) in the common garden. From the 7th of August 2010, we divided the trial in two contrasted experimental conditions: “watered” (from block 1 to 25, 2953 seedlings) versus “water-stressed” (from block 26 to 50, 2860 seedlings). In the “watered” condition, seedlings boxes were watered daily until saturation. The weight of three control boxes was measured when water percolation stopped and allowed to define the average box weight at field capacity (W_{PF1}). The properties of the sandy substrate, measured in the laboratory, made it possible to define the expected box weight at the wilting point ($W_{PF2.5}$) corresponding to 30% of the permanent wilting point. In the “water-stressed” condition, from August to October each year, 3 control boxes were weighed every day ($W_{control}$), and seedlings were not watered as long as $W_{control}$ was above $W_{PF2.5}$. As soon as $W_{control} < W_{PF2.5}$, seedlings were watered again.

2.3 Measurement of phenotypic traits on seedlings in the trial

Fifteen “raw” traits, related to growth, leaf phenology, leaf morphology, leaf physiology, and competition, were directly measured on all or part of the seedlings (Table 1; “2_Phenotype_Table” spreadsheet in the seedlings database). Other “estimated” traits were computed from these raw traits and are available in the “2_Phenotype_Table” spreadsheet in the seedlings database. This data paper gathers the details of the protocols and provides the raw data. It also briefly describes the measured and estimated traits (more details can be found in Gauzere et al. 2016a, b, 2020).


2.3.1 Growth traits

Collar diameter and total height were measured in April 2010 (D0, H0), September 2010 (D1, H1), May 2011 (D2, H2), and November 2011 (D3, H3). The increment rate between September 2010 and November 2011 was computed as: $Diff_D = (D3 - D1)/D1$ and $Diff_H = (H3 - H1)/H1$; $Diff_D$ and $Diff_H$ were analyzed in Gauzere et al. (2016a, b), (2020).

2.3.2 Phenological traits

The timing of bud burst was monitored weekly from April to May in 2011 and 2012. Five stages were used to follow the bud burst dynamics. Moreover, we attempted to bridge the gap between these five stages and the most widespread methodology used to describe phenological stages in plants (i.e., BBCH, Meier et al. 2009), to allow

Fig. 1 Climatic space explored by the three beech plots studied in comparison to the whole species range. This climatic space is represented as the variation of temperature and summer precipitation over beech distribution area, extracted for the SAFRAN database for the period 1958–2015 (collected on an 8 km² grid represented by grey dots). Each black triangle represents the average climate recorded since 2007 with HOBO weather stations for the plot N1 (1020 m), N2 (1140 m), and N4 (1340 m a.s.l.). Reproduced from Gauzere et al. (2020)



the possible posterior analysis and comparison of these data. To that aim, we matched each of our five stages to the closest BBCH reference stage, considering the BBCH scale adapted to trees and shrubs (<https://tempo.pheno.fr/Presentation/Variables-mesures>). The five stages were: (A) buds are dormant (equivalent to the stage 00 in the BBCH scale) or swelling (equivalent to the stage 01 in the BBCH scale); (B) bud scales are broken (BBCH 07); (C) at least 15% of the leaves are emerging (BBCH 08); (D) at least 50% of the leaves are emerging (BBCH 09); and (E) 90% of leaves are spread out (BBCH 19). We also noted dormant buds as stage “0”. The instructions for these measurements are available in the file “SOP_Phenology.pdf” at <https://doi.org/10.15454/6HETQP/WF3JYB>. Then, the sum of budburst scores (Sum_BSS) was computed as in Bontemps et al. (2017): after converting the letters A, B, C, D, and E into marks (from 1 to 5), we summed the marks over all of the dates for each individual. The higher Sum_BSS, the earlier and quicker was leaf unfolding.

The timing of leaf senescence was monitored weekly from October to November in autumn 2011. Three stages were used to follow senescence dynamics (SOP file “Phenology”): (0) leaves have not fallen and are not colored; (1) at least 10% of the leaves are colored or have fallen; and (2) at least 50% of the leaves are colored or have fallen. The instructions for these measurements are available in the file “SOP_Phenology.pdf” at <https://doi.org/10.15454/6HETQP/WF3JYB>. We also computed the sum of senescence scores over all of the dates for each individual (Sum_SenescScores). For both budburst and senescence surveys, phenology was always monitored by the same two groups of observers.

These raw phenological data were transformed to estimate the dates of passage from stage A to B or from B to C for bud burst phenology and the date of passage from stage 0 to 1 or from stage 1 to 2 for leaf senescence (see details in Gauzere et al. 2016a, b).

In November 2011, some seedlings started flushing because of exceptionally warm early autumn temperatures. This unusual event did not lead to the development of fully functioning leaves, but it increased the sensitivity of buds to frost during the winter of 2011/2012. The damaged buds did not restart their development in spring 2012. In May 2012, after all individuals had finished their spring development, we visually quantified the percentage of damaged buds (A: no damages; B: less than 25%; C: between 25 and 50%; D: more than 50%).


2.3.3 Leaf morphological traits

We first measured the fresh leaf area (LA in cm²) with a planimeter. Three leaves were collected on each seedling. The leaves were then dried at 60 °C during about 3 days to finally record the leaf dry mass (LM in mg). The leaf mass area was calculated as $LMA = LM/LA$ (in mg/cm²). The average values of the three estimates of LA, LM, and LMA were kept in the database.

We also measured three other morphological qualitative traits based on scanned images of fresh leaves: the pilosity, presence of lace, and emboss.

The instructions for these measurements are available in the file “SOP_LeafMassArea&Morphology.pdf” at <https://doi.org/10.15454/6HETQP/P8OPPF>.

Fig. 2 Location of the three experimental sites on Mont Ventoux (A), and mapping of the *F. sylvatica* trees in each population (B: N1, C: N2, D: N4). The mother-trees are indicated with black points, and all of the other adult trees are indicated with grey points. Reproduced from Gauzere et al. (2013a, 2013b)



2.3.4 Physiological traits

We measured leaf carbon content (in %), leaf nitrogen content (in %), and leaf carbon isotope composition ($\delta^{13}\text{C}$) as a surrogate of intrinsic water use efficiency (Farquhar and Richards 1984; see detail in Gauzere et al. 2016a, b) for a subset of 1594 individuals, representative of plots, families, and blocks (1039 for the “watered” condition, 555 for the “water-stressed” condition). For each of them, we mixed three collected leaves and dried and ground them in a ball mill. A subsample of 1 ± 0.1 mg was weighed into tin capsules. Leaf nitrogen content was measured with a continuous flow elemental analyzer (Carlo Erba NA 1500; CE Instruments, Rodano, Italy) and the carbon isotope composition with a coupled isotope ratio mass spectrometer (ThermoFinnigan; Delta S, Bremen, Germany). $\delta^{13}\text{C}$ was calculated according to the international standard (Vienna Pee Dee Belemnite, VPDB) using the following equation:

$$\delta^{13}\text{C} = \frac{R_{sa} - R_{sd}}{R_{sd}} \times 1000 \quad (1)$$

where R_{sa} and R_{sd} are the isotopic ratios $^{13}\text{C}/^{12}\text{C}$ of the sample and the standard, respectively. The precision of spectrometric analysis (standard deviation of $\delta^{13}\text{C}$) was assessed with internal laboratory reference material with a matrix close to the measured samples (oak leaves, $n = 16$, $\text{SD} = 0.05 \text{‰}$) and precision among the different runs ranged from 0.08 to 0.13 ‰. These analyses were performed at SIVATECH facilities (SILVATECH, INRAE, 2018, Structural and Functional Analysis of Tree and Wood Facility <https://doi.org/10.15454/1.5572400113627854E12>).

2.3.5 Competition

To estimate the competition acting on each focal seedling, we computed the mean height (i.e., the mean of H3 values, see section “Growth traits” above) of the closest neighbors, i.e., of all the seedlings directly adjacent to a focal individual (8 neighbors in theory, but H3 was not available for border and neutral seedlings). This index was called Neighbor-Height in the database.




Fig. 3 Design of the seedlings trial. (a) Scheme of one typical seedlings box on shelf X, line Y, side A. Each box contains 17 pots, with typically 3 border seedlings (not measured) and 14 focal seedlings. The identifier (ID) of one seedlings is shown in blue, and combines the shelf number (PX), the line number (LY), and the coordinates in the box. (b) Picture of the trial. (c) Spatial arrangement of

the “Watered” condition on shelves 1, 2, 3. The condition includes 25 blocks (blocks 1 to 25). Each block includes 10 boxes (in red) arranged on two sides of the same shelf. (d) Spatial arrangement of the “Water-stress” condition on shelves 4, 5, and 6. The condition includes 25 blocks (blocks 26 to 50)

2.4 Microsatellite genotyping, paternity, and mating system analyses

We genotyped 2088 seedlings (35 offspring per family on average), and all the potentially reproductive adults within each plot (690 in total, including the 60 mother-trees) were genotyped at 13 microsatellite markers (see Gauzere et al.

2013a for genotyping details). These markers were FS1-15, and FS3-04 (Pastorelli et al. 2003); sfc0007-2, sfc-1143, and sfc-0161 (Asuka et al. 2004); mfc7 (Vornam et al. 2004); Csolfagus_19, Csolfagus_6, Csolfagus_29, and Csolfagus_31 (Lefèvre et al. 2012); and Csolfagus_7, Csolfagus_25, and Fi05 (G.G. Vendramin, personal communication). Total DNA was extracted from 50 mg wet-weight frozen leaf material

Table 1 Number of phenotyped seedlings for each trait and water condition

Trait category	Measured/computed trait	n	Watered condition	Water-stressed condition	Total
Growth	Height	4	2534	2454	4988
	Diameter	4	2534	2455	4989
Phenology	Budburst 2011	4	2753	2671	5424
	Senescence 2011	5	2909	2860	5769
	Budburst 2012	6	2522	2581	5103
Leaf morphology	Frost damage	1	1900	1170	3070
	Leaf area	1	2662	2345	5007
	Leaf mass	1	2662	2346	5008
	Leaf pilosity	1	2662	2346	5008
Leaf physiology	Leaf emboss	1	2662	2346	5008
	$\Delta^{13}C$	1	1039	555	1594
	C content	1	1039	555	1594
Competition	N content	1	1039	555	1594
	Mean neighbor height	1	2523	2537	4960

This table shows only “raw” traits, i.e., those directly measured (other traits were computed from these raw traits, and are available in the 2_Phenotype_Table). Traits were grouped in five main categories. The number of times an observation is repeated through time is given by *n*

using the protocol for the DNeasy 96 plant kit (Qiagen). DNA concentration and purity were estimated by measuring the absorbance at 260 and 280 nm in a spectrophotometer and by using pulse-field gel electrophoresis on agarose gel. Extracted DNA was stored at -20°C . Samples were amplified using the Type-it Microsatellite PCR kit (Qiagen). The PCR program was 94°C for 15 min, 30 cycles of 94°C for 30 s, 60°C for 90 s, 72°C for 60 s, and final extension 72°C for 30 min. PCR products were analyzed on a MegaBACE 1000 sequencer and scored with the MegaBACE Genetic Profiler (Amersham Biosciences 2003 version 2.2) against an internal size standard (ET400 DNA size markers). Automatic allele assignment was checked and revised manually twice to ensure consistency of genotyping. Among the 2088 seedlings, 2068 genotypes were successfully read (1382 seedlings from the “watered” condition, and 686 seedlings from the “water-stress” condition). The genotypes of all adults were successfully read.

The genotype dataset was used to search for the father of the genotyped seedlings among all the adult trees of the same plot (including the mother-tree in this self-compatible species). We used the likelihood-based software CERVUS version 3.0 (Marshall et al. 1998), with parameters described in details in Gauzere et al. (2013a). The father was retrieved for 1000 among the 2068 genotyped seedlings. This pedigree information can be used to refine quantitative genetic analyses and, in particular, to account for departure from half-sib assumption in progeny test (Gauzere et al. 2016a, b, 2013b). Independently of paternity analyses, the genotype dataset was also used to estimate the probability for each seedlings to originate from migrant pollen (LogLikMigGL), using the mixed effect mating model (MEMM) and its associated program (Gauzere et al. 2013a; Klein et al. 2011).

Part of the genotype dataset was also used to estimate mating system parameters at family level (Gauzere et al. 2013a): the effective number of pollen donors, the migration rate (estimated from MEMM), and the selfing rate (estimated from MEMM).

2.5 Measurement of phenotypic traits on adult trees in situ

All potentially reproductive beech trees were measured in situ. Although these measurements are not directly related to those made in the seedlings trial experiment described here, we still detailed the different variables measured on adult trees, in an attempt to gather together data as exhaustively as possible. First, the diameter at breast height (dbh) of each adult tree was measured. As beech sometimes produces stump shoots resulting in multiple clonal stems (copice), we measured the number of stems (Nstem) per tree. If a tree displayed multiple stems, the diameter of all stems was measured, and the maximum, mean, and sum of diameter of the clonal copies were computed (respectively MaxDbh,

MeanDbh and SumDbh). For single stem trees, MaxDbh = MeanDbh = SumDbh = Dbh. Additionally, we characterized the stature of each tree through a class variable with 3 levels (dominant, codominant, and suppressed). These levels aimed at accounting for differences in light accessibility among trees with their crown, respectively, above, within or below the surrounding canopy.

Using the spatial coordinates, the conspecific local density was estimated based on the number of reproductive beech neighbors found in disks with a radius of 5 and 20 m around each tree (ConDens5 and ConDens20, respectively). The total competitor density (TotDens5 and TotDens20) was estimated similarly considering trees of all species within these disks.

We also used the Martin-Ek index (Martin & Ek 1984) to quantify the intensity of competition on a focal individual i . This index accounts simultaneously for the diameter and the distance of each competitor j to the competed individual i :

$$\text{MartinTot}_{d_{\max}} = \frac{1}{\text{dbh}_i} \sum_{j=1}^{n_{d_{\max}}} \text{dbh}_j \exp \left[\frac{-16d_{ij}}{\text{dbh}_i + \text{dbh}_j} \right] \quad (2)$$

where dbh_i and dbh_j are the diameter at breast height (in cm) of the competed individual i and of competitor j (any adult tree of any species with $\text{dbh}_j > \text{dbh}_i$), $n_{d_{\max}}$ the total number of competitors in a given radius d_{\max} (in m) around each individual i , and d_{ij} the distance between individuals i and j . We computed this index within radius of 5 and 20 m (MartinTot5 and MartinTot20, respectively). We also used the Martin-Ek index to characterize the intensity of intra-specific competition, by considering only beech as competitor (MartinCon5 and MartinCon20, respectively).

3 Access to the data and metadata description

The seedlings and adult databases are available at Portail Data INRAE: <https://doi.org/10.15454/6HETQP>. Associated metadata access is at <https://metadata-afs.nancy.inra.fr/geonetwork/srv/fre/catalog.search#/metadata/55aa3d92-3866-48e9-b000-f9cbf2d13d03>.

The seedlings database is made of three tables, available as three spreadsheets in one Excel file (entitled: Oddou-Muratorio_et_al_dataBeechHSTrial.xlsx). The **first table** (1_Seeds_Table) contains information on the 60 mother-trees and on the quality of seed lots collected on them (the spreadsheet Seeds_Table_Legend contains variables' description). It includes 16 variables, in particular the number of healthy, empty, and infested seeds, and values of mating system parameters at mother-tree level (effective number of pollen donors, selfing rate, migration rate).

The **second table** (2_Phenotype_Table) contains the phenotypes of the 5813 focal seedlings, as well as general information on all the 8976 focal, border or neutral seedlings raised in the half-sib trial (the spreadsheet Phenotype_Table_Legend contains variables' description). It includes 77 variables in total, and the main raw phenotypic data are summarized in Table 1.

The **third table** (3_Genotype_Table) provides genotypes at 13 microsatellite markers of a subset of 2068 seedlings, their 60 mother-trees, and the 630 candidate fathers (the spreadsheet Genotype_Table_Legend contains variables' description).

Detailed information on the trial design is given in a last dedicated spreadsheet ("Trial Design"), with a schematic representation of the trial and seedling boxes.

One additional excel file (entitled: Oddou-Muratorio_etal_dataAdult_inSitu.xlsx) contains the phenotypes of adult trees (mother-trees and candidate fathers) measured in situ on the three plots (the spreadsheet Legend contains variables' description). These data were originally published in Oddou-Muratorio et al. (2018).

4 Technical validation

Phenotypic measurements were first validated by careful cross-reading of the tabled values, complemented by numerical and graphical analyses. Genotypic data were validated independently by two different operators; moreover, the genotype of the seedlings was validated by comparison with the genotype of their mother-tree. Every record was revised in relation to the normal range of values for each variable. Related variables were confronted and tested for inconsistencies through correlation of time series analyses, and corrected when necessary.

Laboratory equipment was regularly calibrated, and standards were used on each analysis.

5 Reuse potential and limits

Part of this database has already been used to estimate mating system parameters and the variation in adult fecundity (Gauzere et al. 2013a; Oddou-Muratorio et al. 2018), the heritability of adaptive traits (Gauzere et al. 2016a, b), and microgeographic adaptation (Gauzere et al. 2020). However, the different datasets corresponding to these articles are available on different data portals, with no link between them, and sometimes without the raw data. Hence, a main value of the database published here is to gather all of the raw or derived variables that were measured or computed, for all of the individuals. Moreover, some of these data were never published so far (e.g., the 1_Seeds_Table data, or the leaf morphological variables).

This database can be reused for other quantitative genetics studies, for instance, to test new methods of estimation of quantitative genetic parameters based on open-pollinated progeny design (Gauzere et al. 2016a, b, 2013b). It can also be used for ecophysiological studies, for instance, to investigate the relationships between the different studied ecophysiological traits (Bontemps et al. 2017), or to calibrate ecophysiological models accounting for intra-specific variability (Berzaghi et al. 2019; Oddou-Muratorio et al. 2020). Finally, it could provide new entries in existing ecophysiological databases (e.g., Kattge et al. 2011). There are classical limitations for assembling this dataset with other quantitative genetic datasets (lack of a common reference material), but some material (seed, DNA sample) has been secured and conserved at INRAE URFM, and can be made available for future research projects.

Acknowledgements We are indebted to the numerous colleagues from URFM who contributed to field measurement, in particular to François Lefèvre for advices on the conception of the experimental design, to Hendrik Davi for advices on physiological traits measurements (and for Fig. 1), and to Guillaume Simioni for advices on leaf area measurements.

Code availability Not applicable

Funding This study was funded by EC-supported Network of Excellence Evoltree (GOCE-016322, BEECH initiative), ERA-Net BiodivERsA LINKTREE (ANR-08-Biodiversa-006-06), and ERA-Net BiodivERsA TipTree (ANR-12-EBID-0003) projects

Data availability The dataset of the current study is available at: <https://doi.org/10.15454/6HETQP>.

Declarations

Conflict of interest The authors declare no competing interests.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

References

- Alberto FJ, Aitken SN, Alfá R, González-Martínez SC, Hänninen H, Kremer A et al (2013) Potential for evolutionary responses to climate change - evidence from tree populations. *Global Change Biol* 19:1645–1661. <https://doi.org/10.1111/gcb.12181>

- Asuka Y, Tani N, Tsumura Y, Tomaru N (2004) Development and characterization of microsatellite markers for *Fagus crenata* Blume. *Mol Ecol Notes* 4:101–103. <https://doi.org/10.1046/j.1471-8286.2003.00583.x>
- Berzaghi F, Wright JJ, Kramer K, Oddou-Muratorio S, Bohn FJ, Rey CPO et al (2019) Towards a new generation of trait-flexible vegetation models. *Trends Ecol Evol*:1–15. <https://doi.org/10.1016/j.tree.2019.11.006>
- Bontemps A, Davi H, Lefèvre F, Rozenberg P, Oddou-Muratorio S (2017) How do functional traits syndromes covary with growth and reproductive performance in a water-stressed population of *Fagus sylvatica*? *Oikos* 126:1472–1483. <https://doi.org/10.1111/oik.04156>
- Bontemps A, Klein EK, Oddou-Muratorio S (2013) Shift of spatial patterns during early recruitment in *Fagus sylvatica*: Evidence from seed dispersal estimates based on genotypic data. *Forest Ecol Manag* 305:67–76. <https://doi.org/10.1016/j.foreco.2013.05.033>
- Farquhar GD, Richards RA (1984) Isotopic composition of plant carbon correlates with water-use efficiency of wheat genotypes. *Functional Plant Biol* 11:539–552. Retrieved from. <https://doi.org/10.1071/PP9840539>
- Gauzere J, Klein EK, Brendel O, Davi H, Oddou-Muratorio S (2016a) Using partial genotyping to estimate the genetic and maternal determinants of adaptive traits in a progeny trial of *Fagus sylvatica*. *Tree Genet Genomes* 12:1–13. <https://doi.org/10.1007/s11295-016-1062-3>
- Gauzere J, Oddou-Muratorio S, Gay L, Klein EK (2016b) Partial genotyping at polymorphic markers can improve heritability estimates in sibling groups. *Mol Ecol Resources* 16:1340–1352. <https://doi.org/10.1111/1755-0998.12536>
- Gauzere J, Klein EK, Brendel O, Davi H, Oddou-Muratorio S (2020) Microgeographic adaptation and the effect of pollen flow on the adaptive potential of a temperate tree species. *New Phytologist* 227:641–653. <https://doi.org/10.1111/nph.16537>
- Gauzere J, Klein EK, Oddou-Muratorio S (2013a) Ecological determinants of mating system within and between three *Fagus sylvatica* populations along an elevational gradient. *Mol Ecol* 22:5001–5015. <https://doi.org/10.1111/mec.12435>
- Gauzere J, Oddou-Muratorio S, Pichot C, Lefèvre F, Klein E (2013b) Biases in quantitative genetic analyses using open-pollinated progeny tests from natural tree populations. *Acta Botanica Gallica* 160:225–236. <https://doi.org/10.1080/12538078.2013.822827>
- Kattge J, Díaz S, Lavorel S, Prentice IC, Leadley P, Bönisch G et al (2011) TRY - a global database of plant traits. *Global Change Biol* 17:2905–2935. <https://doi.org/10.1111/j.1365-2486.2011.02451.x>
- Klein EK, Carpentier FH, Oddou-Muratorio S (2011) Estimating the variance of male fecundity from genotypes of progeny arrays: evaluation of the Bayesian forward approach. *Methods Ecol Evol* 2:349–361. <https://doi.org/10.1111/j.2041-210X.2010.00085.x>
- Lefèvre S, Wagner S, Petit RJ, De Lafontaine G (2012) Multiplexed microsatellite markers for genetic studies of beech. *Mol Ecol Resources* 12:484–491. <https://doi.org/10.1111/j.1755-0998.2011.03094.x>
- Marshall TC, Slate J, Kruuk LEB, Pemberton JM (1998) Statistical confidence for likelihood-based paternity inference in natural populations. *Mol Ecol* 7:639–655. <https://doi.org/10.1046/j.1365-294x.1998.00374.x>
- Meier U, Bleiholder H, Buhr L, Feller C, Hack H, Heß M et al (2009) The BBCH system to coding the phenological growth stages of plants – history and publications. *J Für Kulturpflanzen* 61:41–52
- Oddou-Muratorio S, Gauzere J, Bontemps A, Rey JF, Klein EK (2018) Tree, sex and size: Ecological determinants of male vs. female fecundity in three *Fagus sylvatica* stands. *Mol Ecol* 27:3131–3145. <https://doi.org/10.1111/mec.14770>
- Oddou-Muratorio S, Hendrik D, Lefèvre F (2020) Integrating evolutionary, demographic and ecophysiological processes to predict the adaptive dynamics of forest tree populations under global change. *Tree Genet Genomes* 16: 67. <https://doi.org/10.1007/s11295-020-01451-1>
- Oddou-Muratorio S, Gauzere J, Angeli N, Brahic P, Brendel O, De Castro M, Gilg O, Hossan C, Jean F, Lingrand M, Pringarbe M, Rei F, Roig A, Thevenet J, Turion N (2021) Phenotypic and genotypic data of a European beech (*Fagus sylvatica* L.) progeny trial issued from three plots along an elevation gradient in Mont Ventoux, South-Eastern France, <https://doi.org/10.15454/6HETQP>, Portail Data INRAE, V5
- Pastorelli R, Smulders MJM, Van't Westende WPC, Vosman B, Gianini R, Vettori C, Vendramin GG (2003) Characterization of microsatellite markers in *Fagus sylvatica* L. and *Fagus orientalis* Lipsky. *Mol Ecol Notes* 3:76–78. <https://doi.org/10.1046/j.1471-8286.2003.00355.x>
- Richardson JL, Urban MC, Bolnick DI, Skelly DK (2014) Microgeographic adaptation and the spatial scale of evolution. *Trends Ecol Evol* 29:165–176. <https://doi.org/10.1016/j.tree.2014.01.002>
- Vornam B, Decarli N, Gailing O (2004) Spatial distribution of genetic variation in a natural beech stand (*Fagus sylvatica* L.) based on microsatellite markers. *Conserv Genet* 5:561–570. <https://doi.org/10.1023/B:COGE.0000041025.82917.ac>

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Authors and Affiliations

Sylvie Oddou-Muratorio^{1,2} · Julie Gauzere^{1,3} · Nicolas Angeli⁴ · Patrice Brahic⁵ · Oliver Brendel⁶ · Marie De Castro⁷ · Olivier Gilg¹ · Christian Hossann⁴ · Frédéric Jean¹ · Matthieu Lingrand¹ · Mehdi Pringarbe¹ · Frank Rei¹ · Anne Roig¹ · Jean Thevenet¹ · Norbert Turion¹

Julie Gauzere
Julie.Gauzere@ed.ac.uk

Nicolas Angeli
nicolas.angeli@inrae.fr

Patrice Brahic
patrice.brahic@onf.fr

Oliver Brendel
oliver.brendel@inrae.fr

Marie De Castro
marie.de-castro@onf.fr

Olivier Gilg
olivier.gilg@inrae.fr

Christian Hossann
christian.hossann@inrae.fr

Frédéric Jean
frederic.jean@inrae.fr

Matthieu Lingrand
matthieu.lingrand@inrae.fr

Mehdi Pringarbe
mehdi.bellahcene-pringarbe@inrae.fr

Frank Rei
franck.rei@inrae.fr

Anne Roig
anne.roig@inrae.fr

Jean Thevenet
jean.thevenet@inrae.fr

Norbert Turion
norbert.turion@inrae.fr

¹ INRAE, URFM, 84 000 Avignon, France

² Present Address: ECOBIOP Université de Pau et des Pays de l'Adour, E2S UPPA, INRAE, ECOBIOP, Saint-Pée-sur-Nivelle, France

³ Institute of Evolutionary Biology, School of Biological Sciences, University of Edinburgh, Edinburgh EH9 3JT, UK

⁴ Université de Lorraine, AgroParisTech, INRAE, SILVA, Silvatech, F-54000 Nancy, France

⁵ ONF, PNRGF, 13 115 Saint-Paul-lès-Durance, France

⁶ Université de Lorraine, AgroParisTech, INRAE, UMR Silva, 54000 Nancy, France

⁷ INRAE, UEFM, 84000 Avignon, France