



FI: The Fecobiome Initiative

Panagiotis Sapountzis, Serafino Teseo, Saria Otani, Frank Møller Aarestrup, Evelyne Forano, Garrett Suen, George Tsiamis, Bradd Haley, Jo Ann van Kessel, Sharon A Huws

► To cite this version:

Panagiotis Sapountzis, Serafino Teseo, Saria Otani, Frank Møller Aarestrup, Evelyne Forano, et al.. FI: The Fecobiome Initiative. *Foodborne Pathogens and Disease*, 2021, 10.1089/fpd.2021.0082 . hal-03510859

HAL Id: hal-03510859

<https://hal.inrae.fr/hal-03510859>

Submitted on 4 Jan 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License



Open camera or QR reader and
scan code to access this article
and other resources online.

FI: The Fecobiome Initiative

Panagiotis Sapountzis,^{1,i} Serafino Teseo,² Saria Otani,³ Frank Møller Aarestrup,³ Evelyne Forano,¹ Garrett Suen,⁴ George Tsiamis,⁵ Bradd Haley,⁶ Jo Ann Van Kessel,⁶ and Sharon A. Huws⁷

Abstract

Animal husbandry has been key to the sustainability of human societies for millennia. Livestock animals, such as cattle, convert plants to protein biomass due to a compartmentalized gastrointestinal tract (GIT) and the complementary contributions of a diverse GIT microbiota, thereby providing humans with meat and dairy products. Research on cattle gut microbial symbionts has mainly focused on the rumen (which is the primary fermentation compartment) and there is a paucity of functional insight on the intestinal (distal end) microbiota, where most foodborne zoonotic bacteria reside. Here, we present the Fecobiome Initiative (or FI), an international effort that aims at facilitating collaboration on research projects related to the intestinal microbiota, disseminating research results, and increasing public availability of resources. By doing so, the FI can help mitigate foodborne and animal pathogens that threaten livestock and human health, reduce the emergence and spread of antimicrobial resistance in cattle and their proximate environment, and potentially improve the welfare and nutrition of animals. We invite all researchers interested in this type of research to join the FI through our website: www.fecobiome.com

Keywords: research initiative, gastrointestinal tract, microbiota, zoonoses

Background

CATTLE HAVE BEEN our domesticated associates since the dawn of agricultural societies, ca 10,000 years ago. With the complementary contributions of a highly specialized gut microbiota, they efficiently convert plant material into protein mass, utilizing dietary components that cannot be digested by monogastrics. This ability has made them our valuable partners since their domestication, as they provide us with a stable supply of protein in the form of meat and dairy products.

As expected, the microbial community residing in the gastrointestinal tract (GIT) of cattle has been the focus of research efforts for more than half a century. The rumen is the primary

fermentation site (Fig. 1) where a complex microbial community, composed of archaea, bacteria, protozoa, and fungi, is responsible for the degradation and fermentation of dietary components and the provision of digested nutrients to the host (Koike and Kobayashi, 2009; Dodd *et al.*, 2011; Comtet-Marre *et al.*, 2017; Huws *et al.*, 2018; Seshadri *et al.*, 2018; Stewart *et al.*, 2018; Kameshwar *et al.*, 2019; Mizrahi *et al.*, 2021).

When partially digested feed and microbial biomass pass through the rumen, digestion continues in the other GIT compartments, until the contents reach the large intestine (Fig. 1), the last section of the GIT. Here, water and nutrients are absorbed (Moran, 2005; Bergmann, 2017), whereas a prominent microbial biomass (second only to that of the foregut) further ferments the remaining dietary components.

¹Université Clermont Auvergne, INRAE, UMR 0454 MEDIS, Clermont-Ferrand, France.

²School of Biological Sciences, Nanyang Technological University, Singapore, Singapore.

³National Food Institute, Technical University of Denmark, Kongens Lyngby, Denmark.

⁴Department of Bacteriology, University of Wisconsin-Madison, Madison, Wisconsin, USA.

⁵Lab of Systems Microbiology and Applied Genomics, University of Patras, Agrinio, Greece.

⁶Environmental Microbial and Food Safety Laboratory, Beltsville Agricultural Research Center, Agricultural Research Service, United States Department of Agriculture, Beltsville, Maryland, USA.

⁷School of Biological Sciences, Institute for Global Food Security, Queen's University Belfast (QUB), Belfast, United Kingdom.

ⁱORCID ID (<https://orcid.org/0000-0001-6286-3918>).




FIG. 1. Overview of the FI vision and goals, the relevant research areas, and how successful implementation could improve animal health and nutrition and reduce the zoonotic and AMR exposure risk to humans. Black arrows show benefits. On top (highlighted with a light blue background), the two main microbial groups of interest (separated based on whether they are beneficial mutualists or animal/zoonotic pathogens) and their implications to potential human exposure and animal husbandry practices are presented. AMR, antimicrobial resistance; FI, Fecobiome Initiative.

The large intestine is of crucial interest not only because of the metabolic processes in which its resident microbiota engage, but also because it is the primary colonization site of zoonotic and animal pathogens. Cattle, for example, are the primary reservoir of the Shiga toxin-producing *Escherichia coli* (STEC). Because STEC pose no threat to an adult host, they do not trigger any host defenses; they exploit the commensal microbiota to transit through the rumen and subsequently colonize the rectal mucosa of the large intestine (Naylor *et al.*, 2003; Sapountzis *et al.*, 2020). Several 16S rRNA amplicon sequencing studies have suggested that members of the cattle rectal microbiota may be able to exclude zoonotic STEC based on mutual exclusion patterns between the rectal microbial community and STEC (Xu *et al.*, 2014; Zaheer *et al.*, 2017; Stenkamp-Strahm *et al.*, 2018; Wang *et al.*, 2018). Other *E. coli* zoonotic pathotypes include the enterotoxigenic *E. coli*, which colonize the small intestine (Fig. 1) and can be detected in animal feces (Nagy and Fekete, 1999).

Similar to *Escherichia*, *Salmonella* pathotypes infect cattle through the fecal-oral route (Holschbach and Peek, 2018) and colonize the rectal mucosa (Palmer *et al.*, 1985); the establishment of several pathotypes has been suggested

to be prevented by a healthy cattle GIT microbiota (Holschbach and Peek, 2018), although experimental evidence is lacking. Other known zoonoses of ruminants include *Clostridium perfringens* (Fohler *et al.*, 2016), *Listeria monocytogenes*, and *Campylobacter* species (Hannon *et al.*, 2009).

In addition to zoonotic pathogens, which are often but not always, harmless hitchhikers of the bovine host, the large intestine is also an Achilles heel for young bovines, which are susceptible to bacterial infections. Early in life, the epimural microbial communities of the intestine are critical for the development and homeostasis of the GIT immune system (Malmuthuge and Guan, 2017; Huws *et al.*, 2018), whereas certain members, such as lactic acid bacteria, which are dominant in very young calves (Alipour *et al.*, 2018), appear to protect against enteric infections and gut dysbiosis (Malmuthuge and Guan, 2017; Morgavi *et al.*, 2020).

The rapidly developing field of genomics has played an important role in deciphering functions of the symbiotic GIT microbes of animals, including humans, as well as the monitoring of antimicrobial resistance (AMR) and virulence genes. Characterization of the genetic machinery of gut microbes has

improved our understanding of: (i) mutualistic (typically nutritional) services gut microbes provide to their hosts; (ii) microbial interactions; and (iii) proximate and evolutionary mechanisms shaping genomes, including factors that facilitate the development of pathogenic traits that can exploit the host and/or associated microbial communities.

Besides the nutritional aspects that may apply to almost all animals (Bourtzis and Miller, 2003; Matthews *et al.*, 2019; Gardiner *et al.*, 2020; Jahnes and Sabree, 2020), the interplay between microbiota and host has also been proven to impact all aspects of the host's health, development, hormone regulation, and behavior, as almost every physiological process is potentially affected by the microbes residing in the GIT (Hara *et al.*, 2013; Kamada *et al.*, 2013; Kanji *et al.*, 2017; Malmuthuge and Guan, 2017; Butt and Volkoff, 2019; Teseo *et al.*, 2019; Liberti and Engel, 2020; Sapountzis *et al.*, 2020).

For the past 10 years, the use of metagenomics and next-generation sequencing studies has led to important insights in all the earlier mentioned research areas, allowing researchers to: (i) take an integrated approach to the characterization of microbial gene inventories (shotgun metagenomics) and the reconstruction of metagenome-assembled genomes of uncultured symbionts, also known as "dark matter" (Bernard *et al.*, 2018); (ii) perform quantitative comparisons of the diversity and structure of microbial communities (biomarkers such as 16S, 18S, and ITS).

In cattle, genomic approaches (both culture and nonculture-based) have mostly focused on the rumen (Pitta *et al.*, 2016; Seshadri *et al.*, 2018; Stewart *et al.*, 2018; Bickhart *et al.*, 2019). Despite the clear role of the large intestine in many aspects of health, nutrition, and the spread of zoonotic pathogens and AMR, the knowledge of its microbiota is mainly limited to:

- (i) descriptive 16S/18S rRNA targeted sequencing (Holman and Gzyl, 2019),
- (ii) single bacterial species (i.e., *Escherichia* and *Salmonella*) and their antagonists that pose a threat to the cattle host or human consumers (Wang *et al.*, 2017; Holschbach and Peek, 2018), and
- (iii) the occurrence and distribution of AMR genes in the feces (Noelle Noyes *et al.*, 2016a, b; Doster *et al.*, 2018; Zaheer *et al.*, 2018, 2019; Keijser *et al.*, 2019; Liu *et al.*, 2019; Rovira *et al.*, 2019; Weinroth *et al.*, 2019; Gaeta *et al.*, 2020; Haley *et al.*, 2020; Lim *et al.*, 2020; Salaheen *et al.*, 2020; Duarte *et al.*, 2021) (see also <https://www.fecobiome.com/resources/data/>).

As of now, very few functional studies or shotgun metagenomic studies have taken an integrated approach to explore the genomic potential of the beneficial microbial residents, the characterization of animal pathotypes or zoonotic hitchhikers and their interactions with the host or each other, the ecological factors impacting their biology, or the potential relationship between microbiota characteristics and the spread of AMR and virulence factors.

We believe it is timely to establish an initiative that can identify common ground between these types of projects, help exploit the full potential of existing sequencing data and other resources, and provide a platform for the emergence of ideas that can generate novel functional interdisciplinary projects. Examples of such projects include the genomic and functional deciphering of the microbial di-

versity of the large intestine, which can provide the cornerstone for all kinds of follow-up transcriptional, chemical, or metabolomic studies and a global effort to collect microbial isolates representing members from all genera that can be found in cattle GIT.

By helping the emergence of such projects (via experimental and intellectual collaboration), which, in turn, can lead to a better understanding of the global microbial diversity of the GIT microbiome, we are confident that the Fecobiome Initiative (FI) can ultimately limit zoonotic pathogens from entering our food chain, better control animal pathogens that hinder the animal development and productivity, reduce the emergence of AMR, optimize the use of antibiotics in animal husbandry, and potentially maximize the nutritional benefits provided by the microbial community to the cattle host.

FI Vision

We hereby propose the formation of the FI (similar to the Greek letter Φ; also known as the golden ratio in mathematics), an international effort that will promote scientific cooperation, dissemination of research findings, and data transparency to facilitate the advancement of research related to microbial communities primarily of the large intestine of cattle, but also related to the rest of the GIT. The vision of the FI is to offer viable solutions in cattle husbandry that will help reduce the zoonotic and AMR risk, animal pathogens, and reduce or optimize the use of antibiotics, with the goal of improving animal welfare while enhancing meat and dairy production.

FI Mission

Through a dedicated website (www.fecobiome.com) that will serve as a coordination and dissemination agent, the FI will:

- (i) bring together and coordinate collaboration among members of the scientific community (e.g., by organizing workshops to align strategies for funding applications and joint proposals) who are interested in the study of microbes residing in the GIT (primarily the large intestine) of cattle,
- (ii) disseminate knowledge by using language and material targeting not only scientists (working on cattle microbiota-related research) but also nonspecialists (e.g., again by organizing events like webinars or workshops for training purposes) and stakeholders involved in animal husbandry, and thus help bridge the gap between the scientific community and non-subject matter specialists,
- (iii) increase public availability of protocols, relevant material (e.g., primers, probes, antibodies), and isolated microbial strains; and by making data (genomic, metagenomic or other) more accessible to the public (i.e., by contributing whenever possible with supplementary metadata and a built-in interface that will allow easier search and recovery of genomic data).

Why Feces?

The organizers of the FI are interested in the study of gut microbes residing primarily in the last compartment of the

cattle GIT, the large intestine. To achieve this, the FI will focus on samples originating from fresh feces as: (i) they present a reliable representation of the microbial community at the distal end of the GIT, and (ii) they do not need special permits for handling of animals in commercial or other farms. In addition, feces can also contain microbial isolates from representatives of other parts of the GIT such as the small intestine and the rumen, which will be valuable for research related to these parts of the GIT (see also the RMG network; <https://rmgnetwork.org/>).

Fresh feces is an excellent sample for: (i) the isolation of cultivable microbes; (ii) isolation and characterization of nucleotide or protein material for -omics approaches, diagnostic polymerase chain reaction, or other molecular and biochemical methods; (iii) chemical analyses for determining nondigested feed components; and (iv) *in vivo* and *in vitro* experimentation. We would note that, despite our interest in the microbial communities of fecal samples, FI does not argue against the use of material from the GIT of animals that has been collected via direct rectal sampling (e.g., rectal swabs) or necropsies (i.e., GIT parts from slaughtered calves) as this material will remain invaluable for other research questions.

FI Goals

The FI has the potential to benefit research in the areas of ecology, evolution, epidemiology, microbiology, human health, and medicine and veterinary medicine, and it may lead to solutions for the improvement of animal husbandry practices. Some examples include:

- Characterization of the distal end microbiome will allow us to formulate strong hypotheses about the nature of the relationship between members of the microbial community and the cattle host, the intertwined dynamics among the community members (cooperative or competitive interactions), and the functional complementarity or redundancy across compartments (e.g., when examined in comparison with similar available datasets from the rumen; Hungate dataset). Insights gleaned from such studies can significantly improve our understanding of the homeostasis and functioning of the GIT microbial community.
- Characterizing the microbial variability on a global scale will allow us to link the variability to host and environmental factors (through correlations of big data sets) such as climate, geographic location, diets, and farm management practices that may affect microbial establishment and resilience. This will allow us to formulate hypotheses about the key factors that influence the GIT microbial community.
- The Whole Genome Sequencing approaches may reveal patterns of genomic evolution, since comparisons of the isolated bacterial strains with their close relatives can reveal traits that have evolved as a result of the unique ecosystem in the large intestine.

The FI can directly contribute to applied solutions in animal husbandry, including the discovery of novel direct-fed microbials to potentially protect against pathogens during early development, to prevent zoonotic pathogens from entering and establishing themselves in the GIT, to limit the

spread of AMR bacteria, and potentially to improve animal nutrition and productivity.

In addition, the FI could lead to improvements in the assessment of zoonotic risks, since currently zoonotic risk and the presence of pathogens have been evaluated by targeting only a handful of known pathogens (mainly *Escherichia* and *Salmonella* genera) and with culture-based approaches. By promoting culture-independent approaches that examine the entire fecal community in parallel with culture-based approaches, we will enhance our ability to identify emerging pathogens or other threats.

Short-Term and Long-Term Objectives and Benefits to the Community

The FI will identify knowledge gaps and focus on research that will complement existing knowledge regarding the microbial inhabitants of the large intestine.

To achieve this, the FI will prioritize research areas such as:

- (i) Characterization of the microbial community of cattle feces by using high-quality/in-depth shotgun metagenomic sequencing to make detailed gene catalogs of the microbial genomic inventory present in the large intestine and identify potential pathogenic and AMR-associated genes.
- (ii) Isolation of microbial strains from fecal samples using both taxa-specific and nonspecific approaches and create a representative library of native aerobic and anaerobic isolates.
- (iii) Using complementary -omics approaches to assemble evidence-based data as these approaches can link, among other factors, DNA sequences, proteins, and metabolites.
- (iv) Large-scale correlations between (meta) genomic data to identify potential factors responsible for the variations between microbial communities (e.g., ecological and geographic datasets, diets, husbandry practices, breed, age, sex, etc.). These correlations require data and isolates from cattle of every breed and age around the globe. We recognize this is an ambitious goal, but we are hopeful that, once the FI is launched, the FI can facilitate data collection, leading to novel results that fuel future projects that allow us to achieve our goals.

We note that the dissemination of data and knowledge is one of the core missions of the FI and toward this end we plan to create a wiki portal (as part of the fecobiome.com website) that will present relevant scientific articles in a simplified and concise view to allow nonsubject matter experts to better connect with the scientific community. Bridging the gap between animal husbandry stakeholders and researchers has the potential to significantly increase the availability of fecal samples from farms, as currently the communication gap between these two parties is often an obstacle.

As (meta)genomic data accumulate, the map of fecal microbial diversity will become more complete, and accomplishing the FI short- and long-term objectives will aid all projects related to the GIT distal end microbiome:

- Microbial isolations and genomic sequencing will provide reference genomes that will serve as templates

for meta-genomic, transcriptomic, and proteomic studies and the identification of novel key genes and taxa (e.g., to be used as markers). This includes pathogenic taxa and virulence genes that can be used as biomarkers, as global comparisons will allow for the evaluation of the most relevant threats.

- Complementary *in vitro* and *in vivo* methods applied to microbial isolates can evaluate their primary and secondary metabolic potential and growth dynamics; insight that, when coupled with metagenomic data, can provide a reconstruction of the community metabolic network and potential ecological interactions.
- Characterizing the large intestine microbes will foster synergies with existing networks related to the cattle GIT microbiome (i.e., RMG network) and ongoing projects, as the combined results can provide an overview of the community structure and mechanisms along the GIT and allow for an integrated approach to the study of the GIT.

Launching the FI

Considering the growing number of shotgun metagenomic fecal data in cattle during the past 3 years (Noelle Noyes *et al.*, 2016a, b; Doster *et al.*, 2018; Zaheer *et al.*, 2018, 2019; Liu *et al.*, 2019; Rovira *et al.*, 2019; Salaheen *et al.*, 2019, 2020; Weinroth *et al.*, 2019; Gaeta *et al.*, 2020; Haley *et al.*, 2020; Lim *et al.*, 2020; Duarte *et al.*, 2021), launching the FI is timely, as its website (fecobiome.com) can serve as a central repository to organize metadata and facilitate meta-analyses to fully exploit the potential of such data.

For this purpose, we have developed a search tool at fecobiome.com (a beta version is already available online at: www.fecobiome.com/resources/) that will allow users to easily locate genomic and other data of interest. Researchers can submit the metadata of their studies through the website (<https://www.fecobiome.com/contact/submit-data/>). Fecal microbial isolates are already available at the National Research Institute for Agriculture, Food and the Environment (France), the Institute for Global Food Security at Queen's University Belfast (Ireland), and the Danish Technological Institute (Denmark) and the metainformation of these isolates will be shortly available at our website. Similarly, a list of protocols and reagents along with research labs that can provide them will also be made publicly available through www.fecobiome.com.

The FI website will serve as a platform for connection and communication among members (an internet forum is an upcoming feature), where members can exchange ideas related to our initiative. The website administrators will be responsible for regulating the threads and for monthly news updates. An Internet forum can also facilitate communication between members, which will initially take place through the website. However, in the coming years, the FI will actively pursue funding for organizing meetings and workshops where members can interact in person.

We invite all interested parties to join the FI. It will be a dynamic site and, through the continuous feedback of members, the FI will adapt to the needs of fecobiome related research in the coming years.

Disclosure Statement

No competing financial interests exist.

Funding Information

This publication was supported in part by a start package grant from the National Research Institute for Agriculture, Food and the Environment.

References

- Alipour MJ, Jalanka J, Pessa-Morikawa T, Kokkonen T, Sato-kari R, Hynönen U, *et al.* The composition of the perinatal intestinal microbiota in cattle. *Sci Rep* 2018;8:1–14.
- Bergmann GT. Microbial community composition along the digestive tract in forage- and grain-fed bison. *BMC Vet Res* 2017;13:253.
- Bernard G, Pathmanathan JS, Lannes R, Lopez P, Baptiste E. Microbial dark matter investigations: How microbial studies transform biological knowledge and empirically sketch a logic of scientific discovery. *Genome Biol Evol* 2018;10:707–715.
- Bickhart DM, Watson M, Koren S, Panke-Buisse K, Cersosimo LM, Press MO, *et al.* Assignment of virus and antimicrobial resistance genes to microbial hosts in a complex microbial community by combined long-read assembly and proximity ligation. *Genome Biol* 2019;20:153.
- Bourtzis K, Miller TA. *Insect Symbiosis*, 1st ed. Boca Raton: CRC Press, 2003.
- Butt RL, Volkoff H. Gut microbiota and energy homeostasis in fish. *Front Endocrinol (Lausanne)* 2019;10:9.
- Comtet-Marre S, Parisot N, Lepercq P, Chaucheyras-Durand F, Mosoni P, Peyretaille E, *et al.* Metatranscriptomics reveals the active bacterial and eukaryotic fibrolytic communities in the rumen of dairy cow fed a mixed diet. *Front Microbiol* 2017;8:67.
- Dodd D, Mackie RI, Cann IKO. Xylan degradation, a metabolic property shared by rumen and human colonic Bacteroidetes. *Mol Microbiol* 2011;79:292–304.
- Doster E, Rovira P, Noyes NR, Burgess BA, Yang X, Weinroth MD, *et al.* Investigating effects of tulathromycin metaphylaxis on the fecal resistome and microbiome of commercial feedlot cattle early in the feeding period. *Front Microbiol* 2018;9:1715.
- Duarte ASR, Röder T, Van Gompel L, Petersen TN, Hansen RB, Hansen IM, *et al.* Metagenomics-based approach to source-attribution of antimicrobial resistance determinants—Identification of reservoir resistome signatures. *Front Microbiol* 2021;11:601407.
- Fohler S, Klein G, Hoedemaker M, Scheu T, Seyboldt C, Campe A, *et al.* Diversity of *Clostridium perfringens* toxinigenotypes from dairy farms. *BMC Microbiol* 2016;16:199.
- Gaeta NC, Bean E, Miles AM, de Carvalho DUOG, Alemán MAR, Carvalho JS, *et al.* A cross-sectional study of dairy cattle metagenomes reveals increased antimicrobial resistance in animals farmed in a heavy metal contaminated environment. *Front Microbiol* 2020;11:590325.
- Gardiner GE, Metzler-Zebeli BU, Lawlor PG. Impact of intestinal microbiota on growth and feed efficiency in pigs: A review. *Microorganisms* 2020;8:1886.
- Haley BJ, Kim S-W, Salaheen S, Hoving E, Van Kessel JAS. Differences in the microbial community and resistome structures of feces from preweaned calves and lactating dairy cows in commercial dairy herds. *Foodborne Pathog Dis* 2020; 17:494–503.

- Hannon SJ, Allan B, Waldner C, Russell ML, Potter A, Babiu LA, et al. Prevalence and risk factor investigation of *Campylobacter* species in beef cattle feces from seven large commercial feedlots in Alberta, Canada. *Can J Vet Res* 2009; 73:275–282.
- Hara N, Alkanani AK, Ir D, Robertson CE, Wagner BD, Frank DN, et al. The role of the intestinal microbiota in type 1 diabetes. *Clin Immunol* 2013;146:112–119.
- Holman DB, Gzyl KE. A meta-analysis of the bovine gastrointestinal tract microbiota. *FEMS Microbiol Ecol* 2019;95:fiz072.
- Holschbach CL, Peek SF. *Salmonella* in dairy cattle. *Vet Clin North Am Food Anim Pract* 2018;34:133–154.
- Huws SA, Creevey CJ, Oyama LB, Mizrahi I, Denman SE, Popova M, et al. Addressing global ruminant agricultural challenges through understanding the rumen microbiome: Past, present, and future. *Front Microbiol* 2018;9:2161.
- Jahnes BC, Sabree ZL. Nutritional symbiosis and ecology of host-gut microbe systems in the Blattodea. *Curr Opin Insect Sci* 2020;39:35–41.
- Kamada N, Chen GY, Inohara N, Núñez G. Control of pathogens and pathobionts by the gut microbiota. *Nat Immunol* 2013;14:685–690.
- Kameshwar AKS, Ramos LP, Qin W. Metadata analysis approaches for understanding and improving the functional involvement of rumen microbial consortium in digestion and metabolism of plant biomass. *J Genomics* 2019;7:31–45.
- Kanji S, Fonseka TM, Marshe VS, Srivatsakumar V, Hahn MK, Müller DJ. The microbiome-gut-brain axis: Implications for schizophrenia and antipsychotic induced weight gain. *Eur Arch Psychiatry Clin Neurosci* 2018;268:3–15.
- Keijser BJF, Agamennone V, van den Broek TJ, Caspers M, van de Braak A, Bomers R, et al. Dose-dependent impact of oxytetracycline on the veal calf microbiome and resistome. *BMC Genomics* 2019;20:65.
- Koike S, Kobayashi Y. Fibrolytic rumen bacteria: Their ecology and functions. *Asian Australas J Anim Sci* 2009;2009:131–138.
- Liberti J, Engel P. The gut microbiota—Brain axis of insects. *Curr Opin Insect Sci* 2020;39:6–13.
- Lim S-K, Kim D, Moon D-C, Cho Y, Rho M. Antibiotic resistomes discovered in the gut microbiomes of Korean swine and cattle. *Gigascience* 2020;9:giaaa043.
- Liu J, Taft DH, Maldonado-Gomez MX, Johnson D, Treiber ML, Lemay DG, et al. The fecal resistome of dairy cattle is associated with diet during nursing. *Nat Commun* 2019;10:4406.
- Malmuthuge N, Guan LL. Understanding the gut microbiome of dairy calves: Opportunities to improve early-life gut health. *J Dairy Sci* 2017;100:5996–6005.
- Matthews C, Crispie F, Lewis E, Reid M, O'Toole PW, Cotter PD. The rumen microbiome: A crucial consideration when optimising milk and meat production and nitrogen utilisation efficiency. *Gut Microbes* 2019;10:115–132.
- Mizrahi I, Wallace RJ, Moraïs S. The rumen microbiome: Balancing food security and environmental impacts. *Nat Rev Microbiol* 2021;19:553–566.
- Moran J. How the rumen works. Tropical Dairy Farming Feeding Management for Small Holder Dairy Farmers in the Humid Tropics. In: Moran J (ed.) Melbourne: CSIRO Publishing, 2005, pp. 41–49.
- Morgavi DD, Popova M, Yanez-Ruiz D, Forano E. Colonization and establishment of the rumen microbiota—Opportunities to influence productivity and methane emissions. In: McSweeney C, Mackie RI (eds.) *Improving Rumen Functions*. S.l.: Burleigh Dodds Science Publishing Limited, 2020.
- Nagy B, Fekete PZ. Enterotoxigenic *Escherichia coli* (ETEC) in farm animals. *Vet Res* 1999;30:259–284.
- Naylor SW, Low JC, Besser TE, Mahajan A, Gunn GJ, Pearce MC, et al. Lymphoid follicle-dense mucosa at the terminal rectum is the principal site of colonization of enterohemorrhagic *Escherichia coli* O157:H7 in the bovine host. *Infect Immun* 2003;71:1505–1512.
- Noyes Noelle R, Yang X, Linke LM, Magnuson RJ, Cook SR, Zaheer R, et al. Characterization of the resistome in manure, soil and wastewater from dairy and beef production systems. *Sci Rep* 2016a;6:24645.
- Noyes Noelle R, Yang X, Linke LM, Magnuson RJ, Dettenwanger A, Cook S, et al. Resistome diversity in cattle and the environment decreases during beef production. *Elife* 2016b;5: e13195.
- Palmer JE, Whitlock RH, Benson CE, Becht JL, Morris DD, Acland HM. Comparison of rectal mucosal cultures and fecal cultures in detecting *Salmonella* infection in horses and cattle. *Am J Vet Res* 1985;46:697–698.
- Pitta DW, Indugu N, Kumar S, Vecchiarelli B, Sinha R, Baker LD, et al. Metagenomic assessment of the functional potential of the rumen microbiome in Holstein dairy cows. *Anaerobe* 2016;38:50–60.
- Rovira P, McAllister T, Lakin SM, Cook SR, Doster E, Noyes NR, et al. Characterization of the microbial resistome in conventional and “Raised Without Antibiotics” beef and dairy production systems. *Front Microbiol* 2019;10:1980.
- Salaheen S, Kim SW, Hovingh E, Van Kessel JAS, Haley BJ. Metagenomic analysis of the microbial communities and resistomes of veal calf feces. *Front Microbiol* 2020;11:609950.
- Salaheen S, Kim SW, Karns JS, Hovingh E, Haley BJ, Van Kessel JAS. Metagenomic analysis of the fecal microbiomes from *Escherichia coli* O157:H7-shedding and non-shedding cows on a single dairy farm. *Food Control* 2019; 102:76–80.
- Sapountzis P, Segura A, Desvaux M, Forano E. An overview of the elusive passenger in the gastrointestinal tract of cattle: The Shiga toxin-producing *Escherichia coli*. *Microorganisms* 2020;8:877.
- Seshadri R, Leahy SC, Attwood GT, Teh KH, Lambie SC, Cookson AL, et al. Cultivation and sequencing of rumen microbiome members from the Hungate1000 Collection. *Nat Biotechnol* 2018;36:359–367.
- Stenkamp-Strahm C, McConnel C, Magzamen S, Abdo Z, Reynolds S. Associations between *Escherichia coli* O157 shedding and the faecal microbiota of dairy cows. *J Appl Microbiol* 2018;124:881–898.
- Stewart RD, Auffret MD, Warr A, Wiser AH, Press MO, Langford KW, et al. Assembly of 913 microbial genomes from metagenomic sequencing of the cow rumen. *Nat Commun* 2018;9:870.
- Teseo S, van Zweden JS, Pontieri L, Kooij PW, Sørensen SJ, Wenseleers T. The Scent of Symbiosis: Gut bacteria may affect social interactions in leaf-cutting ants. *Anim Behav* 2019;150:239–254.
- Wang L, Qu K, Li X, Cao Z, Wang X, Li Z, et al. Use of bacteriophages to control *Escherichia coli* O157:H7 in domestic ruminants, meat products, and fruits and vegetables. *Foodborne Pathog Dis* 2017;14:483–493.
- Wang O, McAllister TA, Plastow G, Stanford K, Selinger B, Guan LL. Interactions of the hindgut mucosa-associated microbiome with its host regulate shedding of *Escherichia coli* O157:H7 by cattle. *Appl Environ Microbiol* 2018;84: e01738-17.

- Weinroth MD, Martin JN, Doster E, Geornaras I, Parker JK, Carlson CR, *et al.* Investigation of tylisin in feed of feedlot cattle and effects on liver abscess prevalence, and fecal and soil microbiomes and resistomes. *J Anim Sci* 2019;97:4567–4578.
- Xu Y, Dugat-Bony E, Zaheer R, Selinger L, Barbieri R, Munns K, *et al.* *Escherichia coli* O157:H7 super-shedder and non-shedder feedlot steers harbour distinct fecal bacterial communities. *PLoS One* 2014;9:e98115.
- Zaheer R, Dugat-Bony E, Holman D, Cousteix E, Xu Y, Munns K, *et al.* Changes in bacterial community composition of *Escherichia coli* O157:H7 super-shedder cattle occur in the lower intestine. *PLoS One* 2017;12:e0170050.
- Zaheer R, Lakin SM, Polo RO, Cook SR, Larney FJ, Morley PS, *et al.* Comparative diversity of microbiomes and Resistomes in beef feedlots, downstream environments and urban sewage influent. *BMC Microbiol* 2019;19:197.
- Zaheer R, Noyes N, Ortega Polo R, Cook SR, Marinier E, Van Domselaar G, *et al.* Impact of sequencing depth on the characterization of the microbiome and resistome. *Sci Rep* 2018;8:5890.

Address correspondence to:
Panagiotis Sapountzis, PhD
Université Clermont Auvergne
INRAE, UMR 0454 MEDIS
Clermont-Ferrand 63000
France

E-mail: panagiotis.sapountzis@inrae.fr