



Revisiting the use of reeds on organic farms in the Atlantic marshes

Daphné Durant, Anne Farruggia, Alexandre Tricheur, Michel Prieur, Pierre Roux

► To cite this version:

Daphné Durant, Anne Farruggia, Alexandre Tricheur, Michel Prieur, Pierre Roux. Revisiting the use of reeds on organic farms in the Atlantic marshes. Organic World Congress, Sep 2021, Rennes, France. hal-03523256

HAL Id: hal-03523256

<https://hal.inrae.fr/hal-03523256>

Submitted on 12 Jan 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



FROM ITS ROOTS
organic inspires life.

 **Organic World**
Congress **2021**

FRANCE

FORUM: Science

Innovative designs for transitions and research



**Organic World
Congress 2021**

FRANCE

**Organic World
Congress 2021**

FRANCE

WWW.OWC.IFOAM.BIO/2021

REVISITING THE USE OF REEDS ON ORGANIC FARMS IN THE ATLANTIC MARSHES

Daphné DURANT

INRAE, DSLP, F-17450 Saint Laurent de la Prée, FRANCE

D. DURANT, A. FARRUGGIA, A. TRICHEUR




**RÉPUBLIQUE
FRANÇAISE**
*Liberté
Égalité
Fraternité*

INRAE

The research farm

Experimental farm belonging to



the French National Research Institute for
Agriculture, Food and Environment

Mixed crop-livestock farming system (160 ha),
organic farming since 2017



Transi'marsh was launched in 2009

**→ a system-experiment dedicated to the agroecological transition for
farms in marshes**

<https://www6.nouvelle-aquitaine-poitiers.inrae.fr/dslp/>

Context

The herd spends 5 months of the year in a building (early November → early April) in free stalling

Needs for straw:



120-130 t / year



Tracking down innovative practices
funding from the Région Nouvelle-Aquitaine



→ exploring alternative sources of bedding

One of the principles of agroecology:

Preserving the natural resources present on a farm (by maintaining their renewal capacity) and the associated ecosystem services


**What natural resource present in marshlands could
be used for bedding?**



Why not the common reed?
Phragmites australis



The experimental design



- Measurements/observations :
- animals' cleanliness
 - temperature of bedding
 - level of soiling of bedding
 - temperature and humidity in the sheds



Results

● Animal cleanliness:

2018: the quantities of bedding (straw or reed) were not sufficient to keep the animals clean

2019: the animals were kept cleaner than in 2018. By increasing the amount of bedding (~ 5 kg/m² of bedding at each mulching, with 3 mulchings per week), no significant difference in the cleanliness of cows between straw and reed bedding



→ Reeds used as bedding are as effective as straw



- Cost of reed: €53 / t (in 2018) and €92 / t (in 2019)

It can compete with straw economically

It is at best half the price, and at worst about the same price as buying straw 'delivered to the yard' (~ €100 / t)



→ The closer the reed bed is to the farm, the more competitive it is



What is reed worth as pasture fodder?

A test of grazing on a small reed bed (~ 2 000 m²) in August 2018...



The composition of simulated bites:

	Fodder unit	Crude protein content	Clutter unit
Reed	0.61	108 g/kg DW	1.54
Permanent grassland	0.61	60 g/kg DW	1.29

Conclusion and implications of this study

Reed is a good alternative to cereal straw

Its cost can be competitive compared to straw

Reed can also constitute a good additional fodder in summer as this plant has interesting nutritive value

➔ Reeds could be given a place on organic livestock farms that wish to make better use of natural resources, while also preserving them and biodiversity!

Many advantages of this natural resource

Provided that the reeds are **'managed' sustainably**:

Respecting the biological cycle of the plant by harvesting late in the season enables the reed beds to retain their role as a habitat for many animal species



The trials, visited by farmers (December 2019)

Many thanks to:

- Michel Prieur (INRAE) and Pierre Roux (INRAE) as contributors
- Matthieu Marquet and 4 farmers from the *Parc Naturel Régional de Brière*
- The *Fédération des Chasseurs de Charente-Maritime* (cabane de Moins)
- The Charente-Maritime *Département*



Utilization of Common Reed (*Phragmites australis*) as Bedding for Housed Suckler Cows: Practical and Economic Aspects for Farmers

Daphné Durant *, Anne Farruggia and Alexandre Tricheur

INRAE, DSLP, F-17450 Saint Laurent de la Prée, France; anne.farruggia@inrae.fr (A.F.); alexandre.tricheur@inrae.fr (A.T.)

* Correspondence: daphne.durant@inrae.fr; Tel.: +33-05-46-82-10-50

Received: 29 September 2020; Accepted: 20 November 2020; Published: 26 November 2020



Abstract: The common reed (*Phragmites australis*) has long been used in wetlands of the French Atlantic coast as fodder and bedding or roof thatching, among other uses. This article explores the practical and economic aspects of utilizing common reed for housing suckler cows compared to straw. Based on a study conducted over two years on a research farm of the French National Research Institute for Agriculture, Food and the Environment (INRAE), located in the marshes of Rochefort-sur-Mer, we show that reed is a good alternative to cereal straw and its cost is quite competitive compared to straw; the closer the reed bed is to the farm, the more competitive it is. By mobilizing the concept of restoration of natural capital, we lay the foundations for a debate on a possible revival of this ancient practice, with the idea that ecological restoration of reed beds can benefit biodiversity and the economy of wetlands farms.

Keywords: mixed crop-livestock farms; bedding; restoration of natural capital; reed beds; wetlands; bioeconomy

1. Introduction

In 2019, approximately half of the 17.3 million tons of cereal straw produced in France was used as bedding for livestock [1]. Many livestock or mixed crop-livestock farms are not independent in terms of straw for housing their animals. Yet straw is an agricultural resource that in some cases

Durant et al. (2020) Resources 9, 140

DOI: 10.3390/resources9120140



**Organic World
Congress 2021**

FRANCE

SEPTEMBER 6TH TO 10TH 2021 IN RENNES

AT THE COUVENT DES JACOBINS • RENNES MÉTROPOLE CONFERENCE CENTRE

THE WORLD ORGANIC CONGRESS IN 2021, IT'S HERE!

WWW.OWC.IFOAM.BIO/2021

Thank you!

FOR MORE INFORMATION PLEASE CONTACT:

papers@owc2020.bio