



Specificity of damage to agriculture due to storms: first recommendations to produce flood damage functions

Pauline Brémond, Frédéric Grelot, Daphné Durant

► To cite this version:

Pauline Brémond, Frédéric Grelot, Daphné Durant. Specificity of damage to agriculture due to storms: first recommendations to produce flood damage functions. European Geosciences Union, Apr 2019, Vienna, Austria. 2022. hal-03523322

HAL Id: hal-03523322

<https://hal.inrae.fr/hal-03523322>

Submitted on 13 Jul 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Specificity of damage to agriculture due to storms: first recommendations to produce flood damage functions

Pauline Brémond¹, Frédéric Grelot¹ and Daphné Durant²

1. Background

1.1 Storm Xynthia, 2010

French littoral and particularly Vendée and Charente-Maritime departments have been seriously impacted by storm Xynthia which occurred on the 28th of February 2010.

32 000 ha of agricultural lands representing 956 farms were flooded with saline water. 55% of farms impacted had more than 50% of their land plots impacted by saline submersion. The main crops impacted by Xynthia were cereal crops or mixed crop-livestock farming systems and viticulture. French littoral is protected by dykes: first rank dykes protect mainly agricultural areas whereas urban areas are rather enclosed by second rank dykes. A lot of infrastructures, in particular first rank dykes have been destroyed by Xynthia.

1.2 Reconstruction of new protections / Cost Benefit Analysis

After Xynthia, local communities asked for fundings to reconstruct and implement new protections.

In France, funding from the State needs to be asked through the procedure PAPI (Programs of Actions to Prevent Floodings). In this PAPI, Cost Benefit Analysis (CBA) is mandatory.

As a consequence, to conduct CBA, Avoided Damage needs to be evaluated. In France, National Flood Damage Functions have been elaborated for non saline floodings and for every category of assets (dwellings, economic activities, agriculture and public infrastructures).

The adaptation of Flood Damage Functions have been proposed for saline floodings for dwelling and economic activities but not yet for agriculture.

1.3 Evaluation of agricultural damage

French National Damage Functions for non saline context have been developed by IRSTEA (Agenais et al, 2013). The approach relies on a conceptual model developed by Brémond et al. (2013) which suggests to distinguish damage to plots (crops, vegetal material, soil and equipment) and damage to farm buildings (building and equipment).

French National Damage functions include damage to plots by considering loss of added value induced by the flood on crops (loss of yield and technical operations required after flooding), vegetal material and soil. All the information collected on loss of yield and recovery actions after flooding have been collected from surveys with agricultural experts (30). Focus groups by category of crop have been organised to validate the Damage Functions. floodam-agri® is the model developed by IRSTEA to simulate agricultural damage due to non saline flooding (fig. 3).




Figure 1: French littoral impacted by Xynthia storm in 2010



Figure 2: Ile de Ré during Xynthia, 2010




Figure 3: Example of flood damage function

2. Research and operational issues

1. To what extent do the mechanisms that induce damage due to saline flooding are specific?
2. To what extent do mid and long term effects should be considered?
3. How long is the recovery of farm after submersion?
4. Which adaptations of the damage functions to agriculture should be done?

3. Method to analyze specificity of saline floodings

We conducted interviews with 8 farmers, 8 agricultural experts and 2 cooperatives in Vendée and Charente-Maritime that have been impacted by Xynthia or have supported farmers' recovery after Xynthia. Those interviews conducted 8 years after Xynthia allowed us to consider **short, mid and long term effects** as well as the **duration of recovery**. Two interview guides have been used: one for the farmers and one for the agricultural experts. All the interviews were transcribed and analysed.

4. Results

4.1 General result

No consolidated expertise on mid and long term effects of storm Xynthia on agriculture have been carried out. Only one precise feedback experience has been found on the INRA experimental farm of Saint-Laurent-de-la-Prée (Durant et al, 2018). The only other submersion that occurred in this area is Martin (December, 1999). With this only two events, it is difficult to analyze potential effects on crops for every season.

4.2 Specificity of damage, specific actions and duration of the recovery



Figure 5: Impact of salinity on soil structure



Figure 6: Mud deposit on soil

Damage to soil are specific due to salinization (fig. 5) and mud deposit composed of sand and algae (fig. 6). To evacuate salt, gypsum has been applied on fields (fig. 7). From our interviews, 20t / ha of gypsum during 3 years have been applied which represents a total cost of 990 €/ha. Efficacy of gypsum also depended on whether rain occurred after treatment.

On vineyard, gypsum could not be applied due to technical constraints and organic matter has been applied.

Damage to drainage equipment (fig. 8) is also specific. Crops along the littoral are equipped with a system of drainage which relies on pump engines and canals. Reparation costs should be considered in damage functions.

Damage to cereals / leguminous plants

Crop losses on winter durum wheat after Xynthia were 100%. Depending on crop type, crop losses varied between 50% and 100% the year Xynthia occurred.

Contrarily to what happens after a non saline flooding, crop losses were endured the years after Xynthia. Effects on yield have been observed between five to seven years after Xynthia due to soil salinity.

Damage to grassland / livestock feeding

Fatalities of cattle have been observed. However, most of barns were out of the area flooded by Xynthia. The main effect on livestock farms was due to flood impacts on grassland. The shortfall induced for feeding cattle has been compensated by buying fodder. The duration of recovery of grassland fields was about three years. Specific damage functions which take into account this linkage should be developed.

Damage to vineyard (fig. 11)

Vineyard mortality due to saline water differs from mortality observed with non saline water. In non saline context, except in spring, mortality is very low. After Xynthia, vineyard mortality occurred and was high and delayed in time (sometimes two years after). The factors that influenced vineyard mortality were: the age of the vineyard, the type of soil and duration of submersion. The duration of the recovery varies between two and seven years.

Damage to potatoes. Potato crop is often coupled with viticulture in this area. After Xynthia, crop loss was 100%. However, no effect was observed the year after on crops.



Figure 4: Fields in Saint Laurent de la Prée



Figure 7: Gypsum application



Figure 8: Drainage equipment in Saint Laurent de la Prée



Figure 9: Crops after Xynthia



Figure 10: Grassland flooded after Xynthia



Figure 11: Vinegrowers in their field after Xynthia

6. Conclusions / Key messages

1. Lack of data on other submersion events to produce damage functions for every season
2. Need to introduce long term effects for non perennial crops whereas long term effects existed only for perennial crops
3. Need to introduce higher rate of vineyard mortality
4. Need to develop damage functions for mix crop livestock farming systems (do not exist yet for non saline flooding)

Bibliography

Agenais, A.-L., et al. (2013). Dommages au secteur agricole : Guide méthodologique et fonctions nationales, GT AMC inondation: 321.
Brémond, P. et al. (2013). "Flood damage assessment on agricultural areas: review and analysis of existing methods." NHESS 13, 2493-2512. doi:10.5194/nheiss-13-2493-2013
Durant, D., et al. (2018). "Impact of storm Xynthia in 2010 on coastal agricultural areas: the Saint Laurent de la Prée research farm's experience." Journal of Coastal Conservation.