

The challenges of modelling mixed management grasslands in North Spain under climate change

Alfredo Rodríguez, Iñigo Gómara, Gianni Bellocchi, Raphaël Martin, Adela Martínez-Fernández, Alfonso Caballal, Jordi Doltra, Agustín del Prado, Margarita Ruiz-Ramos

▶ To cite this version:

Alfredo Rodríguez, Iñigo Gómara, Gianni Bellocchi, Raphaël Martin, Adela Martínez-Fernández, et al.. The challenges of modelling mixed management grasslands in North Spain under climate change. EGU General Assembly 2022, Apr 2022, Vienne, Austria. hal-03530298

HAL Id: hal-03530298 https://hal.inrae.fr/hal-03530298v1

Submitted on 17 Jan2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License



EGU22-12103 EGU General Assembly 2022 © Author(s) 2022. This work is distributed under the Creative Commons Attribution 4.0 License.



The challenges of modelling mixed management grasslands in North Spain under climate change

Alfredo Rodríguez¹, Iñigo Gómara², Giani Bellocchi³, Raphael Martin³, Adela Martínez-Fernández⁴, Alfonso Caballal⁴, Jordi Doltra⁵, Agustín del Prado⁶, and Margarita Ruiz-Ramos⁷

¹Universidad de Castilla-La Mancha, Department of Economic Analysis and Finances, Toledo, Spain

(alfredo.rodriguez@uclm.es)

²Departamento de Física de la Tierra y Astrofísica, Universidad Complutense de Madrid

³Grassland Ecosystem Research, French National Institute for Agricultural Research, INRAe

⁴Servicio Regional de Investigación y Desarrollo Agroalimentario (SERIDA), Asturias

⁵Centro de Investigación y Formación Agrarias (CIFA), Santander, Spain

⁶Basque Centre for Climate Change

⁷CEIGRAM, Universidad Politécnica de Madrid

Permanent grasslands are a very relevant cropping system in the North of Spain and support the main dairy farms in the country. Adaptation to climate change will be required given the projected changes of regional precipitation. To support such adaptation, modelling of these systems to generate high quality projections of the system performance is required. In the region to be simulated, grasslands are managed with a mixture of cuts and grazing. Several issues hinder the modelling of this type of systems: 1) the available data of grazing intensity presents large uncertainties; 2) there are few grassland models that allows flexibility to define a variable combination of cuts and grazing; 3) soil heterogeneity. This exercise has the goal of exploring the potential of PASIM to simulate mixed management grasslands. The model has been successfully applied in a permanent grassland in the French Massif Central, for a management that only contemplated cuts without grazing (Gómara et al., 2020).

The model was calibrated using data from Villaviciosa (Asturias, Spain, 5° 26' 27" W, 43° 28' 50" N, 10 m a.s.l.), located at northern Spain with a temperate climate. This calibration was used to simulate several grasslands locations distributed along the Cantabrian Sea. The soil information was obtained from Trueba et al. (2000). The model was configured for the optimum management for mowing and nitrogen fertilization. The 1976-2005 period and the 2030-2059 period were selected. For the future period two representative concentration pathways emission scenarios (RCP, van Vuuren et al., 2011) were selected (i.e. RCP4.5 and RCP8.5). An ensemble of climate models will be used from the Coordinated Regional Climate Downscaling Experiment (CORDEX, Giorgi and Gutowski, 2015) previously bias-adjusting them by using the European observational database EOBS (Haylock et al., 2008) with the empirical quantile mapping method included at the climate4R R package (Iturbide et al., 2019).

Modelling was challenging due to a combination of complexity (many processes involved) and uncertainty (observed data are difficult to generate). The results of the simulation exercise allow for assessing PASIM skill to reproduce the performance of these complex systems, as well as to determine the main weakness of the model and the observational data and field experiments required to develop the corresponding model improvements.

References

Giorgi, F. and Gutowski, W.J., 2015. Annual Review of Environment and Resources, 40(1): 467-490. Gómara I, Bellocchi G, Martin R, Rodríguez-Fonseca B, Ruiz-Ramos M, 2020. Agricultural and Forest Meteorology, 280, 107768.

Haylock, M.R., Hofstra, N., Klein Tank, A.M.G., Klok, E.J., Jones, P. and New, M., 2008. J. Geophys. Res., 113: D20119.

Iturbide, M., Bedia, J., Herrera, S., Baño-Medina, J., Fernández, J., Frías, M.D., Manzanas, R., San-Martín, D., Cimadevilla, E., Cofiño, A.S. and Gutiérrez, J.M., 2019. Environ. Modell. Softw., 111: 42-54.

Trueba, C., Millán, R., Schimd, T, Lago, (2000). CIEMAT. ISBN: 84-7834-370-9. Madrid.

van Vuuren, D.P., Edmonds, J., Kainuma, M., Riahi, K., Thomson, A., Hibbard, K., Hurtt, G.C., Kram, T., Krey, V., Lamarque, J.-F., Masui, T., Meinshausen, M., Nakicenovic, N., Smith, S.J. and Rose, S.K., 2011. Clim. Change, 109: 5–31.