



HAL
open science

Potential of brassica cover crop and biofumigation to reduce *Verticillium dahliae* germination and Sunflower *Verticillium Wilt*

Neïla Ait Kaci Ahmed, Dessere Diane, Desplanques Jérémy, Galaup Benoit, Dechamp-Guillaume Grégory, Celia Seassau

► To cite this version:

Neïla Ait Kaci Ahmed, Dessere Diane, Desplanques Jérémy, Galaup Benoit, Dechamp-Guillaume Grégory, et al.. Potential of brassica cover crop and biofumigation to reduce *Verticillium dahliae* germination and Sunflower *Verticillium Wilt*. Biofumigation 7 Symposium, Mar 2021, Crans-Montana, Switzerland. hal-03537910

HAL Id: hal-03537910

<https://hal.inrae.fr/hal-03537910>

Submitted on 20 Jan 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Potential of brassica cover crop and biofumigation to reduce *Verticillium dahliae* germination and Sunflower Verticillium Wilt

Ait Kaci Ahmed Neila¹, Dessere Diane¹, Desplanques Jérémy¹, Galaup Benoit¹, Dechamp-Guillaume Grégory², Seassau Célia³

¹AGIR, Université de Toulouse, INRAE, INPT, Castanet-Tolosan, France, ²AGIR, Université de Toulouse, INRAE, INPT, INP-ENSAT, Castanet-Tolosan, France, ³AGIR, Université de Toulouse, INRAE, INPT, INP-PURPAN, Castanet-Tolosan, France

Corresponding author : celia.seassau@purpan.fr

AIMS

1. Evaluate *in vitro* the toxicity of 5 selected brassica on *V. dahliae* germination and development
2. Assess *in situ* the biofumigant potential of 3 brassica on Sunflower Verticillium Wilt

CONTEXT



Sunflower Verticillium Wilt (SVW) is caused by the soil-borne fungal pathogen *Verticillium dahliae*. The widespread of the disease is due to short rotations with sunflower. Currently, there is no agronomic solution except for varietal resistance.

This research may provide an interest of **brassica cover crops** and **biofumigation** to **reduce SVW**, instead of leaving the soil bare in winter.



METHODS




Field study, INRAE (Auzeville, France)

- Microsclerotia (MS) or developed *V. dahliae* on growing media (PDA) were exposed to 5 grinded brassica or no brassica for 21 days (Fig. 1). Brassica were chosen for their contrasted profile in GSL.



- The radial growth of the fungus was measured to calculate the AUDPC.

- Sole crop of brassica were cultivated between wheat and sunflower or the soil was left bare (control) for two years.
- Brassica were brown mustard, fodder radish, turnip rape. Biofumigation () was carried out at bud stage.
- Brassica were sown in september et biofumigation in december



- The incidence and severity of SVW were assessed weekly on sunflower for each treatment (2016, n=30 ; 2017, n=75).

RESULTS

- Brassica reduced ($P < 0,01$) the development of *V. dahliae* from MS up to 90% with turnip rape (**Fig. 1**) and the mycelium development up to 90% (brown mustard, data not shown) compared to the control without brassica.
- Effective suppression of *V. dahliae* using mustards is in line with Olivier et al. (1999) and Neubauer et al. (2014).




Fig.1 – AUDPC of the radial growth of *V. dahliae* MS on PDA exposed to grinded brassica or no brassica (control) for 21 days (n = 30 per treatment)

- Brassica cover crop and biofumigation reduced ($P < 0,01$) the incidence and severity of SVW compared to bare soil both years (**Fig. 2**). The highest reduction of SVW was observed after a fodder radish (up to -43% of SVW).
- Similar reduction of plant disease were observed on vegetables crops (Subbarao et al., 2007 ; Larkin et al., 2010).




Fig.2 – Disease severity index of SVW on sunflower after a bare soil or a Brassica cover crop + biofumigation for 2 years of field trials

CONCLUSION

- Grinded Brassica reduced the germination and the development of *V. dahliae* on growing media
- Biofumigation with brassica reduced SVW compared to a bare soil in the field trials both years
- Fodder radish showed the most promising effects on *V. dahliae* regulation.

References

- Larkin RP, Griffin TS, Honeycutt CW. 2010. Rotation and cover crop effects on soilborne potato diseases, tuber yield, and soil microbial communities. *Plant Dis* 94(12): 1491–1502.
 Neubauer C, Heitmann B, Müller C. 2014. Biofumigation potential of Brassicaceae cultivars to *Verticillium dahliae*. *Eur J Plant Pathol* 140(2): 341–352.
 Olivier C, Vaughn SF, Mizubuti ES, Loria R. 1999. Variation in allyliso thiocyanate production within Brassica species and correlation with fungicidal activity. *J Chem Ecol* 25(12): 2687–2701.
 Subbarao KV, Kabir Z, Martin FN, Koike ST. 2007. Management of soilborne diseases in strawberry using vegetable rotations. *Plant Dis* 91(8): 964–972.

Acknowledgments :

INRAE technical staff, Didier Raffailac, Eric Lecloux and the UE Auzeville who carried out field trials