



HAL
open science

Are forest reserves or deadwood retention key elements for bryophyte diversity?

Marion Gosselin, Yann Dumas, Camille Fabbri, Serge Cadet, Nicolas Debaive, Yoan Paillet, Frédéric Gosselin

► To cite this version:

Marion Gosselin, Yann Dumas, Camille Fabbri, Serge Cadet, Nicolas Debaive, et al.. Are forest reserves or deadwood retention key elements for bryophyte diversity?. International Conference of Ecological Sciences, SFE2, Oct 2018, Rennes, France. hal-03540621

HAL Id: hal-03540621

<https://hal.inrae.fr/hal-03540621>

Submitted on 24 Jan 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Are forest reserves or deadwood retention key elements for bryophyte diversity?

Gosselin, M., Dumas, Y., Fabbri, C., Cadet, S.,
Debaive, N., Paillet, Y., Gosselin, F.



Irstea – Nogent-s/-Vernisson, France



ONF, France

Office National des Forêts



RNF – Dijon, France

Réserves
Naturelles
DE FRANCE



Strategy for forest biodiversity conservation

Forest reserves **left unharvested** are a central part of the strategy for biodiversity conservation...

→ Land sparing (segregation)



... even though other approaches can improve biodiversity by **integrating biodiversity-friendly practices within management** (extending rotations, deadwood, pioneer stages ...)

→ Land sharing (integration)






State of knowledge

European (Paillet et al. 2010 *Conserv. Biol.*) and Global meta-analyses revealed a diversity of taxonomic answers to forest harvesting cessation

- ↪ **Trends towards mostly negative effect of forest harvesting on local species richness**
- ↪ **... with strong variations between taxa...**


Global meta-analysis (2014)



State of knowledge

The European meta-analysis (Paillet et al. 2010 *Conserv. Biol.*) also revealed important gaps


- *less temperate studies*
- *sampling often problematic (e.g. pseudoreplication)*
- *explanatory factors often not incorporated*



Questions

↪ Do the meta-analyses results account for French temperate forests?


Species richness



↪ Are there some key structural variables behind the reserve effect ?

↪ For bryophytes, do climatic variables play a role in addition to or interaction with management or stand structural variables (Raabe et al. 2010)?

A multi-site sampling design



- ↳ 213 stands in 15 French forests
- ↳ Balanced between managed and unmanaged stands (>20 years), on similar site types (topography, soil)

Time since the last harvest

(min Max)


MAN: 9 ± 12 years (0 49)

UNM: 46 ± 38 years (8 148)

Models and magnitude analysis


- Bayesian Generalized linear mixed effects models to explain the **richness** of each taxonomic group
- Simulation of a change ΔX in the explanatory variable to assess the **magnitude of the effect** (Barbier *et al.*, 2009) on the mean of the response variable

Strong effect



$$\frac{R_{\text{new}}}{R_{\text{init}}} \gg 1$$

Low effect



At a 10% threshold:

$$0.9 < \frac{R_{\text{new}}}{R_{\text{init}}} < 1.1$$


Notations: 10% threshold: + strong positive, – strong negative, 00 low
 20% threshold: ++, --, 0
 ? : no possible conclusion

Reserves highly enhance richness of forest bryophytes and rare fungi

Gosselin et al. (2014)
Research Report

n = 213 plots







(9 lowland forests,
6 mountain forests)



Multiplicative factor of the change MAN ⇒ UNM

Three taxonomic groups with strong + to ++ effects in their best model

Criteria:
DICmarg

 <p>Bryophytes Forest</p>	Volume of large deadwood	
 <p>Fungi All</p>	Deadwood volume	
 <p>Rare</p>	Proximity to protected area	

Gosselin et al.,
Rdvt ONF, (2017)

The MAN vs UNM model was not the best model for these groups.

Bryophytes: do climatic data enhance the models?

1) Forest bryophyte group

The best model remains the simple « Volume of large deadwood » threshold model

Best Models	Type	DICm	Variable	Sign.	Magn.
Large deadwood volume (LDV)	┌	529.2	LDV	***	++
Deadwood volume + Humidity	/	529.9	DV	***	+
			Hum	*	0
Deadwood volume + Light	∩	532.5	DV	***	+
			Light	ns	0
Deadwood volume* Humidity	/	533	DV	***	+
			Hum	*	0
			DV*hum	ns	00
Deadwood volume	/	535	DV	***	+
Deadwood volume * Light	/	535.5	DV	***	+
			Light	ns	0
			DV*Light	ns	00

Bryophytes: do climatic data enhance the models?

The best model remains the simple « Volume of large deadwood » threshold model

1) Forest bryophyte group






Best Models	Type	DICm	Variable	Sign.	Magn.
Large deadwood volume (LDV)	┌	529.2	LDV	***	++
Deadwood volume + Humidity	/	529.9	DV	***	+
			Hum	*	0
Deadwood volume + Light	/		DV	***	+
			Light	ns	0
Deadwood volume* Humidity	/		DV	***	+
			Hum	*	0
			DV*hum	ns	00
Deadwood volume	/	535	DV	***	+
Deadwood volume * Light	/	535.5	DV	***	+
			Light	ns	0
			DV*Light	ns	00

Climatic variables enhance other models in addition to (or interaction with) deadwood volume, with significant though negligible effects

Bryophytes: do climatic data enhance the models?

2) All bryophytes

The best model remains the simple
« **Volume of deadwood** »
sigmoid model - Magn = nc





Best Models	Type	DICm	Variable	Sign.	Magn.
Deadwood volume (DV)		734.9	DV	***	nc
Deadwood volume * Altitude		735.5	DV	***	nc
			Alt	ns	0
			DV*Alt	ns	00
Deadwood volume + Humidity		736.1	DV	***	0
			Hum	ns	00
Large living tree volume (LV)		736.2	LV	***	0
Deadwood volume + Light		736.5	DV	***	+/nc/0
			Light	ns	00/00/00
Mean living tree diameter (Dq)		535.5	Dq	***	0/00/00

Bryophytes: do climatic data enhance the models?

2) All bryophytes

The best model remains the simple « Volume of deadwood » sigmoid model - Magn = nc



Best Models	Type	DICm	Variable	Sign.	Magn.
Deadwood volume (DV)		734.9	DV	***	nc
Deadwood volume * Altitude		735.5	DV	***	nc
			Alt	ns	0
			DV*Alt	ns	00
Deadwood volume + Humidity			DV	***	0
			Hum	ns	00
Large living tree volume (LV)			LV	***	0
Deadwood volume + Light		736.5	DV	***	+/nc/0
			Light	ns	00/00/00
Mean living tree diameter (Dq)		535.5	Dq	***	0/00/00

Topographic and Climatic variables appeared among the best models in addition to (or interaction with) deadwood volume, with low effects


Large living trees and Mean diameter also were among the best models

Forest bryophytes: possible applications from the best model



Forest Bryophytes

Sp. richness




Mean value in managed plots : 5 m³/ha



Conclusion

⇒ **Strong positive effect of reserves on forest bryophytes and red-listed lignicolous fungi**




⇒ **Best models for bryophytes = deadwood metrics**

→ Climatic data enhance the models explaining forest bryophytes richness in addition to deadwood or living wood metrics, but the strongest effects are those of deadwood volume.

Discussion

⇒ The utility of segregative and of integrative practices is confirmed for the richness of these taxa



- Limits : very few old or big reserves (recent policy)
- Possible applications need to be discussed with forest managers

+ 5 forest bryophyte species

GAIN

+ 2 to 7 m³/ha large deadwood

COST

Interest ? Feasibility ?



😊😊 Many thanks to 😊😊

- 😊 Your attention!
- 😊 French Ministry of Ecology & ONF for funding
- 😊 All the persons (~100) that were involved at some point in the project