



**HAL**  
open science

## Unlocking genomic diversity of pepper (*Capsicum* spp.) collections held in genebanks: perspectives for breeding and germplasm management

Tripodi P, L Barchi, Acquadro A, Schafleitner R, J. Prohens, M.J. Diez, Börner A, Bovy As, Jérémy Salinier, Boyaci Hf, et al.

### ► To cite this version:

Tripodi P, L Barchi, Acquadro A, Schafleitner R, J. Prohens, et al.. Unlocking genomic diversity of pepper (*Capsicum* spp.) collections held in genebanks: perspectives for breeding and germplasm management. . 17th Eucarpia Meeting on Genetics and Breeding of Capsicum and Eggplant, Sep 2019, Avignon, France. pp.1-3. hal-03551764

**HAL Id: hal-03551764**

**<https://hal.inrae.fr/hal-03551764>**

Submitted on 1 Feb 2022

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Tripodi P, Barchi L, Acquadro A, Schafleitner R, Prohens J, Diez MJ, Börner A, Bovy A, Salinier J, Boyaci HF, Pasev G, Brouwer M, Wallace T, Kale S, Esposito S, Brandt R, Portis E, Finkers R, Lanteri S, Paran I, Lefebvre V, Giuliano G, Stein N. **2019**. Unlocking genomic diversity of pepper (*Capsicum* spp.) collections held in genebanks: perspectives for breeding and germplasm management | BG-O/03. **17th Eucarpia Meeting on Genetics and Breeding of Capsicum and Eggplant**, 11-13 septembre 2019, Avignon (France). p.178-179.

Tripodi P (Orateur),

[Communication orale + Résumé] [Communication orale invitée + Résumé]

URL du congrès ou éditeur	<a href="https://symposium.inra.fr/eucarpiacapsicumeggplant2019/">https://symposium.inra.fr/eucarpiacapsicumeggplant2019/</a>
---------------------------	---

## Title

Unlocking genomic diversity of pepper (*Capsicum* spp.) collections held in genebanks: perspectives for breeding and germplasm management

## Authors

Tripodi P1 , Barchi L2 , Acquadro A2 , Schafleitner R3 , Prohens J4 , Diez MJ4 , Börner A5 , Bovy A6 , Salinier J7 , Boyaci HF8 , Pasev G9 , Brouwer M6 , Wallace T5 , Kale S5 , Esposito S1 , Brandt R5 , Portis E2 , Finkers R6 , Lanteri S2 , Paran I10 , Lefebvre V7 , Giuliano G1 , Stein N5

1 CREA Research Centre for Vegetable and Ornamental Crops, Pontecagnano Faiano (SA), Italy. 2 DISAFA, University of Torino, Torino, Italy. 3 AVRDC The World Vegetable Centre, Taiwan. 4 Instituto de Conservación y Mejora de la Agrodiversidad Valenciana (COMAV), Universitat Politècnica de València, Valencia, Spain. 5 Leibniz Institute of Plant Genetics and Crop Plant Research (IPK) Gatersleben, Germany. 6 Wageningen University & Research WUR, Wageningen, The Netherlands. 7 INRA, UR1052 GAFL, Avignon, France. 8 Bati Akdeniz Agricultural Research Institute BATEM, Antalya, Turkey. 9 Maritsa Vegetable Crops Research Institute, Plovdiv, Bulgaria 10The Volcani Centre, Bet Dagan, Israel. 11Agenzia Nazionale per le Nuove Tecnologie, l'Energia e lo Sviluppo Economico Sostenibile (ENEA), Roma, Italy

## Abstract

**BACKGROUND** The genetic diversity present in cultivars, landraces and wild relatives of modern crops, represents the source of agriculturally important genes to be discovered in our attempts of making crops more resilient and better adapted to the challenges of modern agriculture. In this context, genebanks are pivotal for the collection, preservation and distribution of germplasm; however, better knowledge of the available genetic variability would help improve the management of plant genetic resources. Here we report the analysis of genome-wide genotyping-by-sequencing (GBS) data on 9,659 pepper accessions retrieved from the major European (CGN, INRA, IPK, UPV) and Asian (AVRDC) genebanks, universities and research centres. The collection analyzed is so far the largest and the most diverse ever studied in pepper. The main objectives of this study are to: a) evaluate the genetic diversity using a large set of single-nucleotide polymorphism (SNP) data; b) detect the overlaps within collections and between genebanks; c) detect misclassification of accessions; d) develop core collections for association mapping studies. **MATERIALS & METHODS** Germplasm studied comprised 7,253 *C. annum* accessions, 1,795 accessions belonging to the other four domesticated species (*C. frutescens*, *C. chinense*, *C. baccatum*, *C. pubescens*) and 43 accessions belonging to the wild species (Table 1). About 3% of the collection was

represented by unknown species. Passport information and 42 phenotypic descriptors for plant, flower and fruit traits have been provided by genebank curators in order to enhance the description of plant material. A two restriction enzyme GBS protocol was used [1]; PstI-MspI have been selected based on in silico digestion of the reference genome accession (CM334) [2]. RESULTS GeGBS data provided more than 10,000 SNPs evenly distributed on the genome and used to characterize genetic diversity, population structure and phylogenetic relationships. The analysis allowed identifying five main clusters (K) corresponding to the most represented Capsicum species in the whole collection (Figure 1). *Capsicum annum* was subdivided into two sub clusters and include admixed samples. Three other clusters were observed for *C. baccatum*, *C. frutescens* and *C. chinense*, respectively. The remaining species, including those corresponding to the group of purple flowers clustered in an admixed group. The principle component analysis (PCA) plot for the first two principal components suggests a geographical differentiation of the Caribbean, Central Asia and European gene-pools, and a distinction of the three main clades according to the most updated classification [3]. Moreover, the observed ~1,000 admixed accessions, and the incomplete separations between *C. frutescens* and *C. chinense*, revealed possible taxonomic misclassifications in the genebanks or gene flow between species resulting in introgressions. The genetic and phenotypic variation existing in the collection studied will be exploited for genome-wide association studies (GWAS). 178 Capsicum and Eggplant EUCARPIA Meeting 2019 | BG-O/03 DISCUSSION & CONCLUSION The genetic information provided in this study is appropriate for a deep investigation of the diversity present across the different Capsicum collections maintained by genebanks. Appropriate strategies including “germplasm genomics” can optimize the management of resources, the exchange of material between genebanks and their use for breeding purposes. Beyond genetic diversity, data can be used for the development of core collections to be exploited in GWAS, facilitating the identification of genomic regions associated with valuable traits and enhancing the efforts made by genebanks in the extensive collection and characterization of Capsicum germplasm resources.

REFERENCES [1] Elshire et al., 2011, Plos One 6: e19379. doi.org/10.1371/journal.pone.0019379 [2] Kim et al., 2014, Nature Genetics 46:270–278 [3] Carrizo-Garcia et al., 2016, Ann. Bot. 118:35-51. doi: 10.1093/aob/mcw079 Figure 1. Population structure of the Capsicum germplasm collection. Five subpopulation clusters inferred by STRUCTURE are represented by different colors : 1) *C. baccatum* ; 2) *C. annum* ; 3) *C. chinense* and *C. frutescens* ; 4) *C. annum* ; 5) *C. chinense*, *C. frutescens* and *C. annum* ; adm = admixed 179 BG-O/03 | Capsicum and Eggplant EUCARPIA Meeting 2019 Table 1. Summary of plant material provided by genebanks and used in this study

ACKNOWLEDGEMENTS Lorenzo Barchi and Arnaud Bovy thank the Agropolis Fondation and the SFR Tersys for covering the registration fees for the 17th EUCARPIA Meeting on Genetics and Breeding of Capsicum and Eggplant. Gancho Pasev thanks the French “Ministère de l’Enseignement Supérieur, de la Recherche et de l’Innovation (program ACCES de CAMPUS France)” for covering the travel and accommodation expenses for the 17th EUCARPIA Meeting on Genetics and Breeding of Capsicum and Eggplant.




Figure 1. Population structure of the Capsicum germplasm collection. Five subpopulation clusters inferred by STRUCTURE are represented by different colors : 1) *C. baccatum* ; 2) *C. annuum* ; 3) *C. chinense* and *C. frutescens* ; 4) *C. annuum* ; 5) *C. chinense*, *C. frutescens* and *C. annuum* ; adm = admixed

Species	N° of accessions sequenced	AVRDC	IPK	UPV	INRA	WUR	ARO	CREA	Uniba	BATEH	NYCRI
<i>C. annuum</i>	7523	4166	908	923	639	264	215	209	94	40	38
<i>C. annuum</i> var. <i>pubescens</i>			4					3			
<i>C. pubescens</i>	728	474	182	10	39	15	21	8	1		
<i>C. chinense</i>	627	325	50	7	104	71	31	37	2		
<i>C. baccatum</i>	395	101	3	24	86		8		1		
<i>C. baccatum</i> var. <i>baccatum</i>		3	3			7		3			
<i>C. baccatum</i> var. <i>pendulum</i>		115	23			17		2			
<i>C. pubescens</i>	45	8	21		7	2	2	2	3		
<i>C. chinense</i>	26	10	5			6	4	1			
<i>C. eximium</i>	8	2	4		2						
<i>C. prostratum</i>	7		5*					1	1		
<i>C. oense</i>	1		1								
<i>C. tomat</i>	1					1					
Undefined	298	286	7	3	1				1		
<b>TOTAL</b>	<b>9659</b>	<b>5490</b>	<b>1194</b>	<b>879</b>	<b>878</b>	<b>383</b>	<b>281</b>	<b>265</b>	<b>103</b>	<b>40</b>	<b>38</b>

\* Listed as *C. baccatum* subsp. *prostratum*

Table 1. Summary of plant material provided by genebanks and used in this study