



Many different recipes but one flavor: A universal nested structure of quantitative interactions between plant genotypes and their parasites, including viruses

Benoît Moury, Jean-Marc Audergon, Sylvie Baudracco-Arnas, Safa Ben Krime, François Bertrand, Nathalie Boissot, Mireille Buisson, Valérie Caffier, Melissa Cantet, Sylvia Chanéac, et al.

► To cite this version:

Benoît Moury, Jean-Marc Audergon, Sylvie Baudracco-Arnas, Safa Ben Krime, François Bertrand, et al.. Many different recipes but one flavor: A universal nested structure of quantitative interactions between plant genotypes and their parasites, including viruses. 18. Rencontres de virologie végétale (RVV 2021), INRAE; CIRAD, Sep 2021, Aussois, France. hal-03553180

HAL Id: hal-03553180

<https://hal.inrae.fr/hal-03553180>

Submitted on 3 Feb 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Many different recipes but one flavor: A universal nested structure of quantitative interactions between plant genotypes and their parasites, including viruses

Benoît Moury¹, Jean-Marc Audergon², Sylvie Baudracco-Arnas³, Safa Ben Krima⁴, François Bertrand⁵, Nathalie Boissot², Mireille Buisson⁶, Valérie Caffier⁷, Mélissa Cantet², Sylvia Chanéac⁸, Carole Constant⁹, François Delmotte¹⁰, Catherine Dogimont², Juliette Doumayrou¹, Frédéric Fabre¹⁰, Sylvain Fournet¹¹, Valérie Grimault¹², Thierry Jaunet¹³, Isabelle Justafré¹⁴, Véronique Lefebvre², Denis Losdat¹⁵, Thierry Marcel⁴, Josselin Montarry¹¹, Cindy E. Morris¹, Mariem Omrani^{1,2}, Manon Paineau¹⁰, Sophie Perrot¹², Marie-Laure Pilet-Nayel¹¹, Youna Ruellan²

¹*Pathologie Végétale, INRAE, 84140 Montfavet, France*

²*GAFL, INRAE, 84140, Montfavet, France*

³*Laboratoires ASL, 755 chemin de Meinajaries, 84140 Montfavet, France*

⁴*University of Paris-Saclay, INRAE, AgroParisTech, UMR BIOGER, 78850 Thiverval-Grignon, France*

⁵*Bayer Seeds SAS, Chemin de Roquemartine Mas Lamy, 13670, Saint-Andiol, France*

⁶*GAUTIER SEMENCES, Route d'Avignon, 13630 Eyragues, France*

⁷*Univ Angers, Institut Agro, INRAE, IRHS, SFR QUASAV, 49000 Angers, France*

⁸*TAKII FRANCE SAS, 660 Chemin de la Crau, 13630 EYRAGUES, France.*

⁹*Sakata Vegetables Europe, Domaine de Sablas, rue du moulin, 30620 Uchaud, France*

¹⁰*SAVE, INRAE, Bordeaux Sciences Agro, ISVV, 33140 Villenave d'Ornon, France*

¹¹*IGEPP, INRAE, Institut Agro, Univ. Rennes, 35653 Le Rheu, France*

¹²*GEVES, 25 rue Georges Morel, CS 900024, 49071 Beaucouzé, France*

¹³*HM.Clause, 1 chemin du Moulin des Ronzières, 49800 La Bohalle, France*

¹⁴*Vilmorin & Cie, Mas Pazac, 30210 Ledenon, France*

¹⁵*RIJK ZWAAN France, La Vernède, 30390 Aramon, France*

Understanding the relationships between host range and pathogenicity for parasites, and between the efficiency and scope of immunity for hosts are essential to implement efficient disease control strategies. In the case of plant parasites, most studies have focused on describing qualitative interactions and a variety of genetic and evolutionary models has been proposed in this context. Although plant quantitative resistance benefits from advantages in terms of durability, we presently lack models that account for quantitative interactions between plants and their parasites and the evolution of these interactions.

Nestedness and modularity are important features to unravel the overall structure of host-parasite interaction matrices. We analysed these two features on 32 matrices of quantitative pathogenicity trait data gathered from 15 plant-parasite pathosystems including viruses, fungi, oomycetes, bacteria, nematodes, and insects. We observed significant modularity in only six of the 32 matrices, with two or three modules detected. In contrast, we found high and significant nestedness in 30 of the 32 matrices. Consistent with the widespread nestedness, we showed additionally that: (i) pathogenicity trait values can be mostly explained by a parasite strain effect and a plant accession effect, with minor parasite-plant interaction effect; (ii) there are positive correlations between the efficiency and scope of the resistance of plant genotypes and (iii) between the host range breadth and pathogenicity level of parasite strains. This latter result questions the efficiency of strategies based on the deployment of several genetically-differentiated cultivars of a given crop species in the case of quantitative plant immunity.