



**HAL**  
open science

## Lettuce lines with potyvirus resistance: differential set for study of strains and new genitors of resistance

Brigitte Maisonneuve, Yannick Bellec, Eric Martin, Hervé Lot, Patrick Gognalons, Benoît Moury

### ► To cite this version:

Brigitte Maisonneuve, Yannick Bellec, Eric Martin, Hervé Lot, Patrick Gognalons, et al.. Lettuce lines with potyvirus resistance: differential set for study of strains and new genitors of resistance. 9. International conference on genetics and breeding of leafy vegetables. Eucarpia Leafy Vegetables 2019, Jun 2019, Olomouc, Czech Republic. JOLA, v.o.s., pp.44-45, 2019, 9th International Conference on Genetics and Breeding of Leafy Vegetables. Eucarpia Leafy Vegetables 2019. Programme and Proceedings of Abstracts. hal-03553254

**HAL Id: hal-03553254**

**<https://hal.inrae.fr/hal-03553254>**

Submitted on 3 Feb 2022

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial - NoDerivatives 4.0 International License

## Lettuce lines with potyvirus resistance: differential set for strains studies and genitors of new resistance

Maisonneuve B.<sup>a</sup>, Bellec Y.<sup>b</sup>, Martin E.<sup>a</sup>, Lot H.<sup>c</sup>, Gognalons P.<sup>c</sup> and Moury B.<sup>c</sup>

<sup>(a)</sup>Inra, UR1052, GAFL (Génétique et Amélioration des fruits et Légumes), CS 60094, 84143 Montfavet, France (e-mail: [brigitte.maisonneuve@inra.fr](mailto:brigitte.maisonneuve@inra.fr))

<sup>(b)</sup>Inra, Génétique et Amélioration des Plantes, 78026 Versailles Cedex, <sup>(c)</sup>Inra, Pathologie Végétale, 84140 Montfavet, France

*Lettuce mosaic virus* (LMV; genus *Potyvirus*) is distributed worldwide. This potyvirus could be destructive for lettuce crops and some strains are seed-transmitted. Two resistance alleles of the *mo1* gene were identified: *mo1*<sup>1</sup> from Gallega de Invierno was used in European breeding and *mo1*<sup>2</sup> from PI 251245, a wild Egyptian lettuce, was used in breeding in the USA. Among different LMV isolates collected in Europe, both alleles conferred a partial resistance and untransmissibility by seeds to the common isolate LMV-0, and were overcome by isolates LMV-E and LMV-13. However, these genes showed a differential reaction to isolates LMV-1 and LMV-9, since only *mo1*<sup>2</sup> was resistant to isolates LMV-1 and LMV-9. To study new LMV isolates or new potyviruses, the set of differential hosts includes butterhead genotypes for susceptibility and for *mo1*<sup>1</sup> and iceberg genotypes for *mo1*<sup>2</sup>. The growth of these lettuce types could be different; moreover iceberg lettuce is more difficult for inoculation and symptom observation due to the leaf structure and head shape. Besides that material, a new resistance gene (*Mo3*), efficient against all LMV isolates, has been identified in *L. virosa* PIVT1398 and introgressed into butterhead-type lettuce. Projects were developed to create a set of near-isogenic butterhead lettuce lines useful to characterize LMV and potyvirus pathotypes.

To create near-isogenic lines with *mo1*, a backcrossing programme was used for introducing the *mo1*<sup>1</sup> gene from the butterhead variety Mantilia or the *mo1*<sup>2</sup> gene from the iceberg variety Salinas88 into LMV-susceptible Girelle and Mariska butterhead cultivars. Between backcrosses (BC), resistance tests were made on F<sub>2</sub> plants with LMV-0 for *mo1*<sup>1</sup> screening and LMV-9 for *mo1*<sup>2</sup>. With this method, four lines with butterhead phenotypes were obtained after four or five BC for *mo1*<sup>1</sup> and eight BC for *mo1*<sup>2</sup>; these lines are named Girelle-*mo1*<sup>1</sup>, Girelle-*mo1*<sup>2</sup>, Mariska-*mo1*<sup>1</sup> and Mariska-*mo1*<sup>2</sup>. That material could be useful for research on new LMV isolates or new potyviruses.

The *Mo3* was introgressed from *L. virosa* into two butterhead lettuce by backcrosses after interspecific F<sub>1</sub> hybrids (Maisonneuve et al, 2018. Plant Pathol). The screenings for *Mo3* were made by tests of resistance to LMV-13, LMV-0 or LMV-9. The two near-isogenic lines created, named Girelle-*Mo3* and Mantilia-*Mo3*, were very useful to demonstrate the efficiency of *Mo3* against a new potyvirus present in South-East France: lettuce Italian necrotic virus (LINV). Some physiological disorders (necrotic streaks on stem or corky stem basis), resulting probably from interspecific crosses, were still observed in bolting and flowering despite seven or nine BC by lettuce cultivars. Nevertheless these lines could be interesting for research and as genitors for breeding.

All *mo1*<sup>+</sup>, *mo1*<sup>1</sup>, *mo1*<sup>2</sup> and *Mo3* near-isogenic lines are released for research laboratories.

### Acknowledgements

This research was supported by grants of the French Ministry of Agriculture (project LMV-Most). The authors thank Michel Pitrat for scientific advises and Loïc Jean for seed production.