



Exosomes, adherent-invasive Escherichia coli (AIEC) and Crohn's disease

Hang Nguyen

► To cite this version:

Hang Nguyen. Exosomes, adherent-invasive Escherichia coli (AIEC) and Crohn's disease. 1ères rencontres clermontoises sur les Vésicules Extracellulaires, Laurent-Emmanuel MONFOULET-INRAE-UNH (Organisateur), Feb 2022, Clermont-Ferrand, France. 10.17180/ev22-co04 . hal-03561327

HAL Id: hal-03561327

<https://hal.inrae.fr/hal-03561327>

Submitted on 8 Feb 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial - NoDerivatives 4.0
International License

Exosomes, adherent-invasive *Escherichia coli* (AIEC) and Crohn's disease

Hang Nguyen

M2iSH (Microbes, intestin, inflammation et Susceptibilité de l'Hôte), U1071 Inserm/Université Clermont Auvergne, INRAe USC-2018

The general objective of our research is to better understand the role of microbiota, in particular some pathobiont bacteria (e.g., *E. coli*) in the etiology of chronic intestinal diseases, such as Crohn's disease (CD). Our research group is the first to report that the ileal mucosa of CD patients is abnormally colonized by adherent-invasive *E. coli* (AIEC), which can invade and replicate inside host cells, inducing inflammation. We have shown that upon AIEC infection, host cells induce autophagy as a key defense mechanism to restrain AIEC intracellular replication. However, AIEC can subvert autophagy by up-regulating the levels of miRNAs 30c and 130a, which target and inhibit expression of ATG5 and ATG16L1, respectively, key proteins required for autophagy activation. This in turn leads to increases in AIEC intracellular replication and AIEC-induced inflammation (Nguyen *et al.*, Gastroenterology 2014. PMID: 24148619).

Recently, we showed that exosomes, extracellular vesicles of 30-100 nm implicating in cell-to-cell communication, are new mediators of host-AIEC interaction with their capacity to activate innate immune responses and to subvert the control of AIEC replication by host cells (Carriere *et al.*, Inflamm Bowel Dis. 2016. PMID: 26595556). Mechanistically, upon AIEC infection, host cells secrete exosomes that can transfer miR-30c and miR-130a to recipient cells, thereby inhibiting autophagy in the later. This consequently leads to AIEC abnormal replication in host cells (Larabi *et al.*, Gut microbes 2020. PMID: 32583714).

The on-going research is focusing on (i) the analysis of mRNA, miRNA and protein composition of the exosomes secreted by AIEC-infected cells and their functional effects, together with the underlying mechanism, in recipient cells, and (ii) establishing a serum-derived exosomal miRNA signature to predict AIEC colonization in CD. In summary, exosomes and their cargos are important mediators in host-AIEC interaction, and the strategy targeting them could have important clinical applications.