



HAL
open science

Fluorescent Labelling of membranes and analysis by Malvern-Panalytical Fluo-NTA

Aymeric Audfray

► **To cite this version:**

Aymeric Audfray. Fluorescent Labelling of membranes and analysis by Malvern-Panalytical Fluo-NTA. 1ères rencontres clermontoises sur les Vésicules Extracellulaires, Laurent-Emmanuel MONFOULET-INRAE-UNH (Organisateur), Feb 2022, Clermont-Ferrand, France. 10.17180/ev22-co06 . hal-03561355

HAL Id: hal-03561355

<https://hal.inrae.fr/hal-03561355>

Submitted on 8 Feb 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial - NoDerivatives 4.0
International License

Fluorescent Labelling of membranes and analysis by Malvern-Panalytical Fluo-NTA

Aymeric Audfray PhD, Field Application Scientist, France, Malvern-Panalytical

As researchers push to gain further insight into extracellular vesicles (EVs), the fundamental need to understand the purity of the sample remains. Depending on the EV isolation technique, the presence of co-purifying proteins in EV samples can be a concern when measuring EV concentration. Protein aggregates that scatter light can be misidentified as vesicles when using Nanoparticle Tracking Analysis (NTA). Addressing this issue is particularly important when studying EV heterogeneity and distinct EV subpopulations.

To overcome this problem, we labelled EVs in mixed samples with CellMaskTMOrange® Plasma membrane dye and characterized EVs using NanoSight NTA (“Nanoparticle Tracking Analysis”) in fluorescence mode (F-NTA). EV labelling with this dye is quick, easy and efficient, and not affected by presence of co-isolated proteins. F-NTA can be a powerful tool in confirmation of the presence of pure EVs, and subpopulations in fractionated samples.