



HAL
open science

A soil biodiversity survey coupled with the national soil quality monitoring network?

Camille Imbert, Lucia Santorufo, Carole Ortega, Claudy Jolivet, Apolline Auclerc, Nolwenn Bougon, Yvan Capowiez, Nathalie Cheviron, Daniel Cluzeau, Jérôme Cortet, et al.

► **To cite this version:**

Camille Imbert, Lucia Santorufo, Carole Ortega, Claudy Jolivet, Apolline Auclerc, et al.. A soil biodiversity survey coupled with the national soil quality monitoring network?. Global Symposium on Soil Biodiversity, FAO, Apr 2021, Rome, Italy. pp.46-52. hal-03576462

HAL Id: hal-03576462

<https://hal.inrae.fr/hal-03576462v1>

Submitted on 16 Feb 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

**A soil biodiversity survey coupled with the national soil
quality monitoring network?**

Camille Imbert^{1}, Lucia Santorufo², Carole Ortega³, Claudy Jolivet⁴, Apolline Auclerc⁵, Nolwenn Bougon⁶, Yvan Capowiez⁷, Nathalie Cheviron⁸, Daniel Cluzeau⁹, Jérôme Cortet¹⁰, Gaëlle Deronzier¹¹, Mickaël Hedde¹², Antoine Lévêque¹³, Florence Maunoury-Danger¹⁴, Laurent Palka¹⁵, Guénola Pérès¹⁶, Lionel Ranjard¹⁷, Benoît Vanhée¹⁸, Cécile Villenave¹⁹, Stanislas Wroza²⁰, Antonio Bispo²¹*

¹INRAE Infosol, F-45000 Orléans, France. Camille.imbert@inrae.fr

²Department of Biology, University of Naples Federico II, 80126 Naples, Italy. lucia.santorufu@unina.it

³INRAE Infosol, F-45000 Orléans, France. Carole.ortega@orleans-metropole.fr

⁴INRAE Infosol, F-45000 Orléans, France. claudy.jolivet@inrae.fr

⁵INRAE Université de Lorraine ENSAIA, LES, F-54500 Vandœuvre-les-Nancy, France. apolline.auclerc@univ-lorraine.fr

⁶OFB, F-94300 Vincennes, France. Nolwenn.bougon@ofb.gouv.fr

⁷INRAE, EMMAH, F-84000 Avignon, France. Yvan.capowiez@inrae.fr

⁸INRAE, EcoSys, Plateforme Biochem-Env, F-78000 Versailles, France.

46enoit46e.cheviron@inrae.fr

⁹Université de Rennes, ECOBIO, F-35380 Plélan le grand, France. 46enoit.cluzeau@univ-rennes1.fr

¹⁰Université Paul Valéry Montpellier 3, CEFÉ, F-34000 Montpellier, France. jerome.cortet@univ-montp3.fr

¹¹OFB, F-94300 Vincennes, France. gaelle.deronzier@afbiodiversite.fr

¹²INRAE Montpellier SupAgro Cirad, Eco&Sols, F-34000 Montpellier, France. mickael.hedde@inrae.fr

¹³OFB CNRS MNHN Patrinat, F-75005 Paris, France. Antoine.leveque@mnhn.fr

¹⁴INRAE Université de Lorraine-ENSAIA, LIEC, F-57000 Metz, France. Florence.maunoury-danger@univ-lorraine.fr

¹⁵MNHN, CESCO, F-75005 Paris, France. palka@mnhn.fr

¹⁶INRAE, Agrocampus Ouest, SAS, F-35000 Rennes, France. guenola.peres@agrocampus-ouest.fr

¹⁷INRAE, Agroécologie, F-21000 Dijon, France. lionel.ranjard@inrae.fr

¹⁸Université Catholique de Lille, Ecologie et biodiversité F-59000 Lille. 46enoit.vanhee@univ-catholille.fr

¹⁹Elisol Environnement, F-30111 Congénies, France. cecile.villenave@elisol-environnement.fr

²⁰OFB, F-94300 Vincennes, France. Stanislas.wroza@afbiodiversite.fr

²¹INRAE Infosol, F-45000 Orléans. antonio.bispo@inrae.fr

Abstract summary

Despite its importance for human activities, soil biodiversity remains largely unknown and threatened. Aware of the need for large-scale monitoring, we studied the feasibility to add a soil biodiversity survey (RMBS-Biodiversity) to the already existing French soil quality monitoring network (RMQS). We worked with soil biodiversity experts to couple RMBS-Biodiversity with RMQS i.e. to choose soil taxons and functions, the methodologies and the sampling design. According to the experts, the sampling design of RMQS fits with a soil biodiversity survey. We propose sampling protocols for *i*) bacteria and fungi, *ii*) protists, *iii*) nematodes, *iv*) mesofauna (particularly springtails), *v*) below-ground macrofauna (earthworms), *vi*) surface macrofauna (beetles and spiders), and considering functions, *vii*) soil porosity, *viii*) enzymatic activity and *ix*) organic matter degradation. The RMBS-Biodiversity data would be used to assess the distribution and monitor the changes in soil biodiversity at the scale of the French territory. In addition, coupling data providing from RMBS with data from RMQS-Biodiversity, allows deciphering links between biodiversity,

physical-chemical characteristics of the soil and human activities (practices, heavy metal loading, and pesticide residues for instance).

Keywords: Large-scale survey, soil taxons, soil functions, soil physical-chemical characteristics, practices, biodiversity distribution

Introduction, scope and main objectives

Soil biodiversity is essential for human activities and particularly for agriculture, by governing soil structure, organic matter degradation, nutrient and water cycles (Wagg *et al.*, 2014). Despite of its importance, soil biodiversity remains largely unknown and threatened by human activities (Decaëns, 2010; Orgiazzi *et al.*, 2016).

From the early 1990's, research programs are launched in order to know better soil biodiversity. In France, several surveys exist or have existed, varying according to sampling design, spatio-temporal breadth and studied taxon and functions: ENVASSO (Bispo *et al.*, 2009), EcoFINDERS (CORDIS, 2015), LUCAS (Toth, Jones & Montanarella, 2013), Bioindicator programme (Bispo, Grand et Galsomies, 2009) and Landmark (LANDMARK, 2015). It remains the need to a global surveillance network of soil biodiversity based on standardized methodologies, covering the whole French territory (metropolitan and ultra-marine areas), including almost the main soil biodiversity components as it is done elsewhere (Dragicevic, 2008; McKenzie, Van Leeuwen et Pinder, 2009; Rutgers, 2011; Schmidt *et al.*, 2011; CENBAM, 2012).

Building a large-scale survey with several methodologies needs a strong organization and substantial resources to cover large spatiotemporal scales. It could be advantageous to lean on an existing survey, already operational. Beyond the organization benefit, this coupling allows to study the links between different data with a high statistical power. Besides, linking different biodiversity surveys is one of the main goals of the French Biodiversity Office (OFB) (Touroult *et al.*, 2017). However, it is necessary to assess if the sampling design of the existing survey could fit with the new one. For example, the grid pixel size and the number of studied size can impact the results (Nielsen *et al.*, 2009; Soberón *et al.*, 2007).

Aware of the need for large-scale monitoring and of the opportunity of this coupling for research issues, we studied the feasibility to add a soil biodiversity survey to the already existing French soil quality monitoring network (RMQS). The objectives of the present work were: 1) Constitute an expert group involving in the feasibility evaluation of a long-term soil biodiversity survey, 2) Choose protocols and assess human, financial and technical needs and 3) Propose a sampling design accommodating all the biodiversity methodologies and the pedologic measurements.

Methodology

In order to study the feasibility to add a soil biodiversity survey (RMBS) to the already existing RMQS, we first installed a group of

experts and discussed the way we could adapt the existing sampling design to biodiversity measurements.

RMQS

The first campaign of the RMQS took place from 2000 to 2015 in continental France and overseas territories (French Guiana, West Indies, Reunion and Mayotte islands) and the second campaign began in 2016. The RMQS is based on the monitoring of 2,240 sites spread over a 16*16 km grid on French territory. These sites can be agricultural plots, meadows, urban gardens, wastelands, forests, vineyards and orchards and other types of natural environments.

For each site, a sampling area of 20m*20m, divided into 2m*2m plots, was selected. Monitoring is carried out on 25 of these plots (those with plot n°2 for the second RMQS campaign, Figure 1). A pedological profile is carried out near the sampling area.




Figure 1: Sampling design of the RMQS

The data collected correspond to the physico-chemical characteristics of the soil, including contaminants, organic carbon stocks and human activities (agricultural or soil management practices). Some biodiversity monitoring has already been done on RMQS plots in three projects. Flora and forestry characteristics are measured since 2000 in plots of the ICP forest network. In the ECOMIC-RMQS project, microorganism community was determined for all RMQS plots. Similarly, fungi analysis is in progress. In the RMQS-Biodiv, several taxa (microorganisms, mites, springtails, nematodes, earthworms) and functions (organic matter degradation and enzymatic activity) were sampled in 107 Brittany sites. More recently, the enzymatic activity is measured on RMQS soil samples collected since 2016 (5 enzymes in 2016 and 2017, 10 enzymes from 2018).

Setting up the soil biodiversity survey

The OFB expressed the need to build a survey of soil biodiversity, which will be part of the larger survey of terrestrial biodiversity. The first step is to demonstrate the feasibility of such surveillance, taking into account scientific and practical aspects. Thus, we contacted around 30 soil biodiversity experts to install a working group where we built a survey. The questionnaire included 59 questions divided into 4 main categories: sampling (16 questions), laboratory and data analysis (23 questions), costs (5 questions) and interpretation of the results (15 questions). The survey was used to gather the opinions and knowledge of the experts (e.g. usefulness of the taxon, budget, way of implementation, interpretation limits) and collect the details of both field and lab methods for each taxon and function. We circulated the survey within the group. We performed face-to-face interviews to discuss more practical aspects (such as field and analysis time, workload, coordination with other teams, analysis difficulties) and collect suggestions for the most efficient way of working. Finally, during the 4 plenary meetings we all together i) discuss the results of the questionnaire, ii) study the feasibility of coupling RMBS with RMQS, iii) choose soil taxa, functions and associated methods, iv) figure out the sampling design and v) list practical needs (labour supply, costs, material, time). Such meetings were also used to explain the aims of the project, present the monitoring network, the advancement of the work, make a synthesis of the different information and reach agreements.

Results

During the last two years, 21 completed answers and 2 uncompleted answers to the questionnaire were received, together with 14 interviews. We discussed the results and shared our views during 4 plenary meetings with all the experts.

The working groups composed by the contacted experts, is representative of almost all the soil biodiversity taxonomic groups, and some functions. According to experts, the sampling design of RMQS seems be appropriate to perform the soil biodiversity survey. However, some aspects of the RMQS protocols were discussed to be adapted to the RMBS-Biodiversity. We decided that the frequency (10-15 years) and season of sampling, and the sampling grid (16*16 Km) of RMQS fits to the objectives of RMBS-Biodiversity.

We collectively propose a survey of i) bacteria and fungi, ii) protists, iii) nematodes, iv) below-ground mesofauna (springtails mainly), v) below-ground macrofauna, vi) surface macrofauna (beetles and spiders) and mesofauna, and considering functions, vii) soil porosity, viii) enzymatic activity and ix) organic matter degradation (Tab. 1). Flora and sporocarp fungi should also be part of the RMBS

but discussions about sampling protocols are still in progress. The expert group still need to be set up for French ultramarine areas.

The development of RMBS-Biodiversity has also allowed federating a multidisciplinary network around soil biodiversity, composed of researchers (pedologists, agronomists and ecologists), botanical conservatories, nature managers, naturalist associations, the administration and private research offices.

Table 1: Taxons, functions and methodologies selected

Taxons- Functions	Data collection	Lab analysis
Bacteria and fungi	Composite sample	Mass sequencing
Protists	Composite sample	Sequencing
Nematodes	Composite sample	Taxonomic identification
Mesofauna	Soil corers	Taxonomic identification
Belowground macrofauna	Spade test and mustard application	Taxonomic identification
Surface macrofauna and mesofauna	Pitfall traps	Taxonomic identification
Soil porosity	Soil column	Soil column scan
Enzymatic Activity	Composite sample	Absorbance measure
Organic matter degradation	Cotton strips	Tension test

Discussion

After 2 years of discussion, meetings and literature review, we considered RMQS sampling design adapted to monitor soil biodiversity at large scale. As there are no studies at different scales of soil biodiversity, so it is difficult to define the most suitable sampling design density and therefore the most appropriate grid cell. Coupling a soil biodiversity survey with a project such as the RMQS, will certainly provide some answers to this question. Indeed, a soil biodiversity survey (RMBS-Biodiversity) using the RMQS tool represents a real opportunity to conduct a study on a national scale. Even if the study grid is quite large in relation to existing work (e.g., plot scale), it will provide a global vision at a national scale of soil biodiversity and thus provide a better understanding of the biogeography of soil organisms. Due to the lack of information about the wide scale distribution of soil organisms, the adaptation of RMQS protocols to soil organism's study should constitute a starting point to understand the main drivers influencing soil biodiversity at large scale.

The RMBS-Biodiversity will start with a testing phase in 2020. A soil expert group is engaged in this project and methodologies were chosen.

The next steps are to identify methodologies to survey Flora and fungi sporocarps and to define sampling designs in ultramarine territories.

Conclusions

Adding a soil biodiversity survey (RMBS-Biodiversity) to the RMQS seems feasible and will be tested. It is a promising project that will contribute to a better knowledge of soil biodiversity on the French territory and to understand links between different kind of soil biodiversity, physical-chemical characteristics of the soil and human activities.

Acknowledgements

The authors thank the French Biodiversity Office (OFB) for funding this project.

References

- Bispo, A., Cluzeau, D., Creamer, R., Dombos, M., Graefe, U., Krogh, P.H., Sousa, J.P., Peres, G., Rutgers, M. & Winding, A.** 2009. Indicators for monitoring soil biodiversity. *Integrated environmental assessment and management*, 5(4): 717-720.
- Bispo, A., Grand, C. & Galsomies, L.** 2009. Le programme ADEME "Bioindicateurs de qualité des sols". *Étude et gestion des sols*, 16(3-4): 145-158.
- CENBAM.** 2012. Programa de Pesquisa em Biodiversidade. In: *CENBAM Portal & PPbio* [online]. [Cited 02 July 2021]. <https://ppbio.inpa.gov.br/en/home>
- CORDIS.** 2015. Ecological Function and Biodiversity Indicators in European Soils (EcoFINDERS) In: *European Commission*. [online]. [Cited 23 July 2021]. <https://cordis.europa.eu/project/id/264465>
- Dragicevic, S.K.** 2008. Croatian soil monitoring programme. , p. 150. Zagreb, Croatian Environment Agency.
- Decaëns, T.** 2010. Macroecological patterns in soil communities. *Global Ecology and Biogeography*, 19(3): 287-302. <https://doi.org/10.1111/j.1466-8238.2009.00517.x>
- LANDMARK.** 2015. LAND Management: Assessment, Research, Knowledge base. In: *European Commission*. [online]. [Cited 23 July 2021]. <https://esdac.jrc.ec.europa.eu/projects/landmark>
- McKenzie, N.L., Van Leeuwen, S. & Pinder, A.M.** 2009. Introduction to the Pilbara biodiversity survey, 2002-2007. *Records of the Western Australian Museum, Supplement*, 78(1): 3-89.
- Nielsen, S.E., Haughland, D.L., Bayne, E. & Schieck, J.** 2009. Capacity of large-scale, long-term biodiversity monitoring programmes to detect

trends in species prevalence. *Biodiversity and Conservation*, 18(11): 2961-2978. <https://doi.org/10.1007/s10531-009-9619-1>

Orgiazzi, A., Panagos, P., Yigini, Y., Dunbar, M.B., Gardi, C., Montanarella, L. & Ballabio, C. 2016. A knowledge-based approach to estimating the magnitude and spatial patterns of potential threats to soil biodiversity. *Science of The Total Environment*, 545-546: 11-20. <https://doi.org/10.1016/j.scitotenv.2015.12.092>

Rutgers, M. 2011. Biological Indicators of Soil Quality (BISQ) in the Dutch Soil Monitoring Network. présenté à, 2011.

Schmidt, O., Keith, A.M. (Aidan), Arroyo, J., Bolger, T., Boots, B., Breen, J., Clipson, N., et al. 2011. *CréBeo Soil Biodiversity Project - baseline data, response to pressures, functions and conservation of keystone micro- and macro-organisms in Irish soils*. Wexford, Environmental Protection Agency.

Soberón, J., Jiménez, R., Golubov, J. & Koleff, P. 2007. Assessing completeness of biodiversity databases at different spatial scales. *Ecography*, 30(1): 152-160. <https://doi.org/10.1111/j.0906-7590.2007.04627.x>

Toth, G., Jones, A. & Montanarella, L. 2013. LUCAS topsoil survey methodology, data and results

Touroult, J., Chaumet, S., Poncet, L. & Siblet, J.P. 2017. *Diagnosis and recommendations for a strategy for acquiring continental naturalist knowledge*. Saint-Aignan de Grand Lieu, France, Val pg Pôle Graphique (also available at https://www.patrinat.fr/sites/patrinat/files/atoms/files/2018/10/spn_2017_-_10_-_strategieconnaissancetomeiweb.pdf)

Wagg, C., Bender, S.F., Widmer, F. & van der Heijden, M.G.A. 2014. Soil biodiversity and soil community composition determine ecosystem multifunctionality. *Proceedings of the National Academy of Sciences*, 111(14): 5266-5270. <https://doi.org/10.1073/pnas.1320054111>