



**HAL**  
open science

## **Biodiversity and vector-borne diseases: Host dilution and vector amplification occur simultaneously for Amazonian leishmaniases**

Arthur Kocher, Josselin Cornuault, Jean-charles Gantier, Sophie Manzi, Agathe Chavy, Romain Girod, Isabelle Dusfour, Pierre-Michel Forget, Marine Ginouves, Ghislaine Prevot, et al.

### ► To cite this version:

Arthur Kocher, Josselin Cornuault, Jean-charles Gantier, Sophie Manzi, Agathe Chavy, et al.. Biodiversity and vector-borne diseases: Host dilution and vector amplification occur simultaneously for Amazonian leishmaniases. *Molecular Ecology*, 2023, 10.1111/mec.16341 . hal-03576716

**HAL Id: hal-03576716**

**<https://hal.inrae.fr/hal-03576716v1>**

Submitted on 27 Jan 2025

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1 **Biodiversity and vector-borne diseases: host dilution and vector amplification**  
2 **occur simultaneously for Amazonian leishmaniases**

3 **Running title:** Biodiversity and Amazonian leishmaniases.

4 Arthur Kocher<sup>1,2,3,4\*</sup>, Josselin Cornuault<sup>5,6</sup>, Jean-Charles Gantier<sup>7</sup>, Sophie Manzi<sup>1</sup>, Agathe Chavy<sup>3,8</sup>, Romain  
5 Girod<sup>3</sup>, Isabelle Dusfour<sup>3</sup>, Pierre-Michel Forget<sup>10</sup>, Marine Ginouves<sup>8,9</sup>, Ghislaine Prévot<sup>8,9</sup>, Jean-François  
6 Guégan<sup>2,11</sup>, Anne-Laure Bañuls<sup>2</sup>, Benoit de Thoisy<sup>3,12</sup>, and Jérôme Murienne<sup>1</sup>

7 <sup>1</sup>Laboratoire Évolution et Diversité Biologique (UMR5174 EDB) - CNRS, IRD, Université Toulouse III  
8 Paul Sabatier - Toulouse, France.

9 <sup>2</sup>MIVEGEC, Université de Montpellier, IRD, CNRS, Montpellier, France.

10 <sup>3</sup>Institut Pasteur de la Guyane, Cayenne, France.

11 <sup>4</sup>Transmission, Infection, Diversification & Evolution Group, Max-Planck Institute for the Science of  
12 Human History, Kahlaische Str. 10, 07745 Jena, Germany.

13 <sup>5</sup>Real Jardín Botánico CSIC, Plaza Murillo 2, 28014, Madrid, Spain.

14 <sup>6</sup>ISEM, Université de Montpellier, CNRS, IRD, EPHE, Montpellier, France.

15 <sup>7</sup>Laboratoire des Identifications Fongiques et Entomo-parasitologiques, Mennecy, France.

16 <sup>8</sup>TBIP, Université de Guyane, 97300 Cayenne, France.

17 <sup>9</sup>Université de Lille, CNRS, Inserm, Institut Pasteur de Lille, U1019-UMR9017-CIIL Centre d'Infection et  
18 d'Immunité de Lille, 59000 Lille, France.

19 <sup>10</sup>Muséum National d'Histoire Naturelle, UMR-7179 MECADEV (Mécanismes Adaptatifs et Evolution),  
20 MNHN-CNRS, Brunoy, France.

21 <sup>11</sup>INRAE, Cirad, Université de Montpellier, UMR ASTRE, Montpellier, France.

22 <sup>12</sup>Association Kwata, Cayenne, French Guiana.

23 \*Correspondence: arthur.kocher@gmail.com

24 **Abstract**

25 Changes in biodiversity may impact infectious disease transmission through multiple  
26 mechanisms. We explored the impact of biodiversity changes on the transmission of  
27 Amazonian leishmaniases, a group of wild zoonoses transmitted by phlebotomine sand flies  
28 (Psychodidae), which represent an important health burden in a region where biodiversity is  
29 both rich and threatened. Using molecular analyses of sand fly pools and blood-fed dipterans,  
30 we characterized the disease system in forest sites in French Guiana undergoing different levels  
31 of human-induced disturbance. We show that the prevalence of *Leishmania* parasites in sand  
32 flies correlates positively with the relative abundance of mammal species known as  
33 *Leishmania* reservoirs. In addition, *Leishmania* reservoirs tend to dominate in less diverse  
34 mammal communities, in accordance with the dilution effect hypothesis. This results in a  
35 negative relationship between *Leishmania* prevalence and mammal diversity. On the other  
36 hand, higher mammal diversity is associated with higher sand fly density, possibly because  
37 more diverse mammal communities harbor higher biomass and more abundant feeding  
38 resources for sand flies, although more research is needed to identify the factors that shape  
39 sand fly communities. As a consequence of these antagonistic effects, decreased mammal  
40 diversity comes with an increase of parasite prevalence in sand flies, but has no detectable  
41 impact on the density of infected sand flies. These results represent additional evidence that  
42 biodiversity changes may simultaneously dilute and amplify vector-borne disease transmission  
43 through different mechanisms that need to be better understood before drawing generalities on  
44 the biodiversity-disease relationship.

45 **Keywords:** zoonotic disease, dilution effect, metabarcoding, iDNA, phlebotomine sand fly,  
46 Culicidae, amplification effect

47 **Introduction**

48 The current biodiversity crisis alters ecosystem functioning and human well-being through a  
49 variety of processes that are still debated (Cardinale et al., 2012; Oliver et al., 2015). In this  
50 context, the impact of biodiversity loss on infectious disease transmission and spread has  
51 become an important research topic during the last two decades. Particular attention has been  
52 given to the potential regulation of pathogens in ecosystems by the presence of species that are  
53 inefficient for their transmission, a mechanism referred to as the “dilution effect” (Keesing et  
54 al., 2010; Ostfeld & Keesing, 2012). Theoretical investigations as well as laboratory and field  
55 experiments have allowed defining the conditions under which a dilution effect is expected to  
56 occur (Dobson, 2004; Johnson, Hartson, Larson, & Sutherland, 2008; Johnson, Lund, Hartson,  
57 & Yoshino, 2009; Mihaljevic, Joseph, Orlofske, & Paull, 2014; Norman, Bowers, Begon, &  
58 Hudson, 1999; Ostfeld & LoGiudice, 2003; Roche, Dobson, Guégan, & Rohani, 2012; Rudolf  
59 & Antonovics, 2005; Suzán et al., 2009; Van Buskirk & Ostfeld, 1995), and empirical studies  
60 have suggested its existence in numerous disease systems (Clay, Lehmer, Jeor, & Dearing,  
61 2009; Derne, Fearnley, Lau, Paynter, & Weinstein, 2011; Ezenwa, Godsey, King, & Guptill,  
62 2006; Gilbert, Norman, Laurenson, Reid, & Hudson, 2001; Gottdenker, Chaves, Calzada,  
63 Saldaña, & Carroll, 2012; LoGiudice, Ostfeld, Schmidt, & Keesing, 2003; Ostfeld & Keesing,  
64 2000; Telfer et al., 2005; Weinstein, Titcomb, Agwanda, Riginos, & Young, 2017).  
65 Considering a local community of hosts composed of species that differ in their competence  
66 for a given pathogen (i.e., their ability to get infected and transmit), a dilution effect may occur  
67 if the presence of the least competent hosts reduces contact rates between the most competent  
68 hosts and the pathogen or decreases the density of competent hosts. Additionally, if the less  
69 competent hosts are also those that tend to be extirpated from species-depleted communities,

70 biodiversity loss should result in enhanced transmission. The idea that the dilution effect may  
71 produce a beneficial impact of biodiversity conservation on public health in most disease  
72 systems, has triggered important interest as well as strong criticism (Civitello et al., 2015;  
73 Halsey, 2019; Keesing et al., 2010; Lafferty & Wood, 2013; Levi et al., 2016; Randolph &  
74 Dobson, 2012; Salkeld, Padgett, & Jones, 2013; Wood et al., 2014). Advances in disease  
75 ecology have allowed a more detailed understanding of biodiversity-disease relationships, and  
76 the field has progressed beyond initial debates about the generality of the dilution effect.  
77 Studies have highlighted the importance of accounting for different factors such as the  
78 geographical scale, the nature and extent of biodiversity changes, the transmission mode, and  
79 the taxa involved in the disease system under consideration (Cohen et al., 2016; Faust et al.,  
80 2017; García-Peña et al., 2016; Gibb et al., 2018; Halliday & Rohr, 2019; Halliday, Rohr, &  
81 Laine, 2020; Johnson, Calhoun Dana M., Riepe Tawni, McDevitt-Galles Travis, &  
82 Koprivnikar Janet, 2019; Johnson, de Roode, & Fenton, 2015; Keesing & Ostfeld, 2021;  
83 Morand, Jittapalapong, Suputtamongkol, Abdullah, & Huan, 2014; Rohr et al., 2020;  
84 Weinstein et al., 2017; Wood & Lafferty, 2013; H. S. Young, Parker, Gilbert, Sofia Guerra, &  
85 Nunn, 2017). It has been shown that changes in biodiversity can either amplify or dilute  
86 pathogen transmission, through multiple mechanisms which can sometimes occur within the  
87 same system (Clay, Lehmer, St. Jeor, & Dearing, 2009; Faust et al., 2017; Huang, van  
88 Langevelde, Prins, & de Boer, 2015; Luis, Kuenzi, & Mills, 2018; Miller & Huppert, 2013;  
89 Ogden & Tsao, 2009; Roche & Guégan, 2011; Rohr et al., 2015; Swei, Ostfeld, Lane, & Briggs,  
90 2011; Wood, Summerside, & Johnson, 2020). In the case of vector-borne diseases, it has been  
91 stressed that the ecology and feeding habits of arthropod vectors must also be considered  
92 (Carlson, Dyer, Omlin, & Beier, 2009; Hamer et al., 2011; Laporta, Prado, Kraenkel, Coutinho,

93 & Sallum, 2013; Loss et al., 2009; McGregor et al., 2018; Miller & Huppert, 2013; Ogden &  
94 Tsao, 2009; Park, Cleveland, Dallas, & Corn, 2016; Randolph & Dobson, 2012; Roche &  
95 Guégan, 2011; Roche, Rohani, Dobson, & Guégan, 2013; Swei et al., 2011; Titcomb et al.,  
96 2017; Vinson & Park, 2019).

97 Overall, there is a need for more field studies on various systems to further disentangle the  
98 mechanisms underlying biodiversity-disease relationships and to inform epidemiological  
99 predictions. However, conducting such studies can be highly challenging, since it requires  
100 generating data on wild vertebrate and arthropod fauna as well as on circulating pathogens in  
101 numerous study sites and, often, in difficult environmental contexts (*e.g.*, tropical regions).  
102 Here, we used recently developed molecular tools to explore the effects of mammal diversity  
103 on the transmission of Amazonian leishmaniasis. Leishmaniasis are a group of human vector-  
104 borne diseases endemic to different tropical and subtropical regions, caused by parasites of the  
105 genus *Leishmania* (Kinetoplastida: Trypanosomatidae) and transmitted by hematophagous  
106 phlebotomine sand flies (Psychodidae) (reviewed in Bañuls, Hide, & Prugnolle, 2007). In  
107 Amazonian ecosystems, several zoonotic *Leishmania* species typically coexist, with distinct  
108 sylvatic transmission cycles involving different sand fly vector species and wild mammal  
109 reservoir hosts (Ralph Lainson & Shaw, 2010; Rotureau, 2006). Amazonian leishmaniasis  
110 represent a significant public health burden in a biodiversity-rich region threatened by human  
111 activities (Chavy et al., 2019; Pan American Health Organization, 2019; Rangel, Costa, &  
112 Carvalho, 2014). However, the mechanisms through which biodiversity may impact the  
113 transmission of these wild zoonoses remain largely unexplored.

## 114 **Material and methods**

### 115 *Sampling*

116 Sampling was performed between 2015 and 2017 in 19 different forest sites in French Guiana  
117 (Figure 1, Table 1, Data S1). Sites were separated by at least ~5-km distance and chosen to  
118 represent sylvatic environments with variable levels of human-induced disturbance, ranging  
119 from remote and protected areas to forest patches in the vicinity of urbanized zones. When  
120 several sites were sampled within a same area (i.e., Saint-Georges, Belizon, Kaw or Counami),  
121 those were specifically chosen to represent contrasting situations with respect to hunting  
122 pressure (e.g., closer to or further from the nearest city or accessible road, inside or outside of  
123 a protected area). Human-induced disturbance was measured using the average Human  
124 FootPrint index [HFP; (de Thoisy et al., 2010); updated in 2012; Table 1, Data S1]. In each  
125 site, sand flies were collected using US Centers for Disease Control and Prevention (CDC)  
126 miniature light traps set up across a *ca.* 1 ha plot. Traps were separated by *ca.* 50-m distance  
127 from each other and left for up to six consecutive nights. Each morning, the contents of each  
128 trap was collected. Sand fly females were sorted using a stereo microscope and kept in pools  
129 (corresponding to each trap-night) in microcentrifuge tubes with 95% ethanol for later  
130 molecular analyses. A maximum of 50 individuals was included in a pool, and several pools  
131 were made when more than 50 specimens were caught in a given trap (with a maximum of  
132 four pools per trap, *i.e.*, 200 individuals). The total number of sand flies caught in each trap  
133 was systematically recorded, unless the contents of the trap was importantly damaged or  
134 partially lost (mainly due to rainy conditions or manipulation during collection). Visibly blood-  
135 fed dipterans, including sand flies, mosquitoes (Culicidae), and biting midges  
136 (Ceratopogonidae), were kept individually in microcentrifuge tubes with 95% ethanol for

137 molecular analyses. Additional blood-fed dipterans resting during the day along tree trunks  
138 were collected using a Prokopack aspirator (John W. Hock co., Gainesville, FL, USA) and  
139 conserved in the same way.

#### 140 ***Laboratory***

141 We analyzed sand fly pools to identify their species composition using a previously developed  
142 DNA metabarcoding protocol (Kocher, Gantier, et al., 2017). *Leishmania* DNA detection and  
143 identification was performed on the same pools using high-throughput sequencing of kDNA  
144 minicircle amplicons (Kocher, Valière, Bañuls, & Murienne, 2018). For each blood-fed  
145 specimen, the dipteran species and blood meal source were identified individually as  
146 previously described (Kocher, de Thoisy, Catzeflis, Valière, et al., 2017). Sand fly pools were  
147 homogenized using a *Qiagen TissueLyser 2* (Qiagen, Valencia, CA, USA), and DNA was  
148 extracted with the *Qiagen DNeasy Blood and Tissue kit*. For individual blood-fed specimens,  
149 a modified *Chelex* (Bio-Rad, Hercules, CA, USA) protocol was used for DNA extraction  
150 (Casquet, Thebaud, & Gillespie, 2012). The Ins16S\_1 [F: TRRGACGAGAAGACCCTATA;  
151 R: TCTTAATCCAACATCGAGGTC; (Clarke, Soubrier, Weyrich, & Cooper, 2014)], 12S-  
152 V5 [F: TAGAACAGGCTCCTCTAG; R: TTAGATACCCCACTATGC; (Riaz et al., 2011)]  
153 and leishmini [F: 5'-GGKAGGGGCGTTCTGC-3'; R: 5'-STATWTTACACCAACCCC-3';  
154 Kocher, Valière, Bañuls, & Murienne, 2018] PCR primers were used to amplify short  
155 fragments of dipteran, vertebrate and *Leishmania* DNA, respectively. Tags of eight base pairs  
156 with at least five differences between them were added at the 5' end of each primer to enable  
157 multiplexing of PCR products for subsequent sequencing (Binladen et al., 2007). A Latin  
158 square design was used for PCR multiplexing to allow for the detection and filtering of  
159 mistagged sequencing read (Esling, Lejzerowicz, & Pawlowski, 2015). For sand fly



160 metabarcoding, two PCR replicates were performed. PCR products were pooled according to  
161 the multiplexing design and used for sequencing library preparation and high-throughput  
162 sequencing on Illumina HiSeq or MiSeq platforms at the GeT-PlaGe core facilities of Genotoul  
163 (Toulouse, France).

### 164 ***Bioinformatics***

165 Bioinformatic analyses were performed using the *OBITools 1.2.9* package (Boyer et al., 2016)  
166 and *R 4.0.3* (R Core team, 2020). Paired-end reads were merged with *illumina-paired-end* and  
167 demultiplexed based on PCR primer tags using *ngsfilter*. Reads were dereplicated using  
168 *obiuniq* and sequences supported by less than ten reads in a given sample were discarded using  
169 *obigrep*. Taxonomic assignments were performed using *ecotag*, with customized reference  
170 DNA sequence datasets for each studied taxonomic group. For dipteran identifications, we  
171 used previously published reference datasets for neotropical sand flies (Kocher, Gantier, et al.,  
172 2017) and mosquitoes (Talaga et al., 2017), to which we added mosquito reference sequences  
173 corresponding to the targeted 16S region, which we extracted from NCBI GenBank using  
174 *ecoPCR*. For vertebrate identifications, we used a previously published reference dataset for  
175 Amazonian mammals (Kocher, de Thoisy, Catzeflis, Huguin, et al., 2017), completed with  
176 vertebrate reference sequences corresponding to the targeted 12S region extracted from  
177 GenBank. For *Leishmania* identifications, we used a previously published dataset of kDNA  
178 minicircle reference sequences (Kocher et al., 2018). *ecotag* employs a Lower Common  
179 Ancestor algorithm that allows to perform taxonomic assignments based on the percentage of  
180 identity with multiple matches in a reference dataset. In other words, a sequence matching  
181 similarly to several members of a given taxon will be assigned to the corresponding taxon. For  
182 dipteran and vertebrate identifications, we considered taxonomic assignments at the genus

183 level at best if the percentage of identity with the closest match was lower than 97%, in order  
184 to avoid biases due to reference dataset incompleteness (*i.e.* artifactual species-level  
185 identifications in cases where only one species was represented in the dataset for a given  
186 genus). We then performed *de novo* sequence clustering using *sumacrust 1.0.31* (Mercier,  
187 Boyer, Bonin, & Coissac, 2013) with a 97% threshold. We defined molecular taxonomic units  
188 (MOTU) based on *ecotag* results in case of species-level identifications, and based on *de novo*  
189 clustering otherwise, in order to identify putative species within upper-level taxa. Vertebrate  
190 identifications were adjusted when only a subset of the matched species was known to be  
191 present in French Guiana. Because no reference sequence was available for local biting midge  
192 species, we defined MOTUs within the Ceratopogonidae family based on *de novo* clustering  
193 only.

194 For each sequencing library, we used the number of sequencing reads found with non-used  
195 primer tag combinations to perform MOTU-based filtering of mistagged reads, as suggested  
196 previously (Esling et al., 2015). Additionally, MOTUs supported by less than 100 reads in a  
197 given sample were filtered out. For sand fly metabarcoding, we further filtered MOTUs that  
198 (i) were not identified as Phlebotominae, (ii) were not recovered in two PCR replicates, and  
199 (iii) were supported by least 2% of the sequencing reads in a given sample (a maximum of 50  
200 sand flies were included in each analyzed pool). For individual blood-fed specimens, the most  
201 supported dipteran and vertebrate MOTUs were retained (*i.e.* we did not consider the  
202 possibility of multiple blood meal sources, as a conservative measure). In a few cases, the  
203 resulting dipteran identification did not match the expected dipteran group (sand fly, mosquito,  
204 or biting midge) and was therefore discarded. For blood meal identifications, results were  
205 discarded if the majority sequence was identified as human or other potential laboratory

206 contaminants which were not expected in our study sites, as well as sequences assigned to  
 207 above-order taxonomic levels which likely represented molecular or sequencing artifacts. For  
 208 *Leishmania* detection, only species-level identifications were considered, and the majority  
 209 *Leishmania* species was retained in each positive sample.

210 ***Estimating mammal diversity from individual dipteran blood meals***

211 iDNA has recently emerged as a promising tool to perform vertebrate inventories at lower cost  
 212 and effort (Calvignac-Spencer et al., 2013; Kocher, de Thoisy, Catzeflis, Valière, et al., 2017;  
 213 Schnell et al., 2015). Here, we used a probabilistic approach to generate estimates of host  
 214 community composition and diversity from our iDNA data (*i.e.* blood meal identifications of  
 215 individual dipteran specimens) while accounting for invertebrate host preferences. The model  
 216 is represented by a Bayesian network in Figure S1. We assume that the probability  $p_{sij}$  that a  
 217 blood-fed invertebrate of species  $i$  has fed on a host of species  $h$  in site  $s$  depends on  $r_{sh}$ , the  
 218 relative abundance of  $h$  in  $s$ , and on  $\alpha_{sh}$ , the relative preference of  $i$  for  $h$ . We further assume  
 219 that the relative probability of insect  $i$  feeding on host  $h_j$  rather than on host  $h_k$  in site  $s$  is given  
 220 by:

221 
$$\frac{p_{sij}}{p_{sik}} = \frac{r_{sh_j} \alpha_{ih_j}}{r_{sh_k} \alpha_{ih_k}} \quad (1)$$

222 From (1) it follows that the probability  $p'_{sij}$  that an invertebrate of species  $i$  has fed on host  $h$   
 223 in site  $s$ , given that it has fed on one of the  $H$  host species identified in our dataset, is:

224 
$$p'_{sij} = \frac{r_{sh} \alpha_{ih}}{\sum_{k=1}^H r_{sk} \alpha_{ik}} \quad (2)$$

225 Denote  $\{N\}_{si} = \{n_{si1}, \dots, n_{siH}\}$  the data vector of the number of occurrences of invertebrate  $i$   
 226 having fed on each of the  $H$  host species in site  $s$ . The probability of this data is given by the  
 227 multinomial distribution with parameters  $\{p'\}_{si} = \{p'_{si1}, \dots, p'_{siH}\}$  and  $Y_{si} = \sum_{k=1}^H n_{sik}$ :

$$228 \quad P(\{N\}_{si} | \{p'\}_{si}, Y_{si}) = \text{Multinomial}(\{N\}_{si} | \{p'\}_{si}, Y_{si}) \quad (3)$$

229 Given  $S$  sites and  $I$  invertebrate species across all sites, we denote  $\{N\} = \{\{N\}_{si}\}$  the whole  
 230 dataset,  $\{p'\} = \{\{p'\}_{si}\}$  the set of probability vectors and  $\{Y\} = \{Y_{si}\}$  the set of sample sizes,  
 231 with  $i = 1, 2 \dots I$  and  $s = 1, 2, \dots, S$ . The likelihood of the full dataset is then given by:

$$232 \quad P(\{N\} | \{p'\}, \{Y\}) = \prod_{s=1}^S \prod_{i=1}^I \text{Multinomial}(\{N\}_{si} | \{p'\}_{si}, Y_{si}) \quad (4)$$

233 The model was implemented in a Bayesian framework in *Stan* (Carpenter et al., 2017) through  
 234 its R interface *rstan*, in order to sample the posterior probability density of the parameters using  
 235 Monte Carlo Markov chain (MCMC; see Supporting Information and Figure S2 for details  
 236 about the choice of priors and simulation results, as well as the *Stan* code). Joint samples of  
 237 mammal's relative abundances in each site were used to derive the Shannon index of diversity  
 238 ( $-\sum_h r_{sh} \log(r_{sh})$ ) and the overall proportion of species known as *Leishmania* reservoirs.  
 239 Closely related dipteran species appeared to feed on similar ranges of hosts (*e.g.* sand flies of  
 240 the *Nyssomyia* genus feeding mostly on arboreal mammals, sand flies of the *Psychodopygus*  
 241 genus feeding mostly on armadillos, mosquitoes of the *Culex* genus feeding on a wide range  
 242 of vertebrates or observed ceratopogonids feeding mostly on amphibians; Table 2; Figure S3).  
 243 Therefore, we estimated feeding preferences at the genus level (or at the family level in the  
 244 case of ceratopogonids), in order to increase statistical power. Additionally, we assumed that  
 245 dipterans had identical preferences for vertebrates of the same order which exhibit similar

246 morphological and ecological features (with the exception of rodents which were separated  
247 into terrestrial and arboreal rodents; Figure S3). Sites in which less than five dipteran blood  
248 meals could be identified were not retained for the analysis. The posterior distribution of the  
249 parameters was sampled using MCMC with 3 chains of 40,000 iterations, including 4,000  
250 iterations for warmup. Convergence and mixing were assessed using trace plots and ESS  
251 values, which were >200 for all parameters (Figures S4 and S5). The prior distribution of the  
252 parameters was obtained by running the same MCMC sampling scheme as for the posterior,  
253 with the likelihood fixed to a constant. The comparison of posterior and prior densities allowed  
254 us to assess to what extent the posterior was driven by the data (Figures S4 and S5). The mean  
255 posterior estimates of parameters were computed and used as variables for the generalized  
256 linear models (GLM) described in the next section (we initially attempted to jointly estimate  
257 mammal relative abundances and GLM's parameter but this led to MCMC mixing issues).

### 258 *Effect of mammal diversity on Leishmania transmission*

259 In addition to the measures of vertebrate communities obtained from individual blood meal  
260 identifications, sand fly counts in traps and molecular analyses of sand fly pools were used to  
261 estimate the abundance and diversity of sand flies as well as the *Leishmania* prevalence rate in  
262 sand flies. All of these variables were used in a series of Bayesian GLMs to explore the impact  
263 of mammal diversity changes on the disease system (Figure 1, Supporting Information, Data  
264 S2). We assessed the relationship between human-induced disturbance and mammal diversity  
265 using a regression of the Shannon index of mammals on the HFP. We assessed whether  
266 changes of mammal diversity led to predictable changes of the mammal community  
267 competence for *Leishmania* parasites using a regression of the relative abundance of  
268 *Leishmania* reservoirs on the Shannon index of mammals. Furthermore, we assessed the effect

269 of changes of mammal diversity on sand fly density and diversity by regressing the number of  
270 sand flies collected in each trap and the Shannon index of sand flies (estimated from  
271 metabarcoding results with the *R* package *iNext*; (Hsieh, Ma, & Chao, 2016)) on the Shannon  
272 index of mammals. We then assessed the effect of mammal community competence, sand fly  
273 density and sand fly diversity on *Leishmania* transmission using a regression of *Leishmania*  
274 prevalence rates in sand flies on the relative abundance of *Leishmania* reservoirs, the expected  
275 number of sand flies collected per trap and the Shannon index of sand flies (used as predictor  
276 variables in the same regression). We checked for spatial autocorrelation of the different  
277 variables with the Moran's *I* autocorrelation index as implemented in the *R* package *ape*  
278 (Paradis, Claude, & Strimmer, 2004), using the inverse of pairwise geographical distances  
279 between sites as the weight matrix. For all regressions, predictor variables were standardized,  
280 and weakly informative normal priors were used for regression coefficients. Prior distributions  
281 of slope coefficients were centered around 0, while that of regression intercepts were centered  
282 around the mid-range value of the corresponding dependent variable. The posterior  
283 distributions of slope coefficients were used to assess the significance of inferred relationships.  
284 We evaluated the cumulative effect of mammal diversity changes on *Leishmania* transmission  
285 by generating posterior predictions of the *Leishmania* prevalence rate and of the expected  
286 number of infected sand flies per trap for different values of the Shannon index of mammals.  
287 We visualized sampled mean prediction curves against the Shannon index of mammals (across  
288 the range of values observed in our study), and used the distribution of their average slope to  
289 measure the predicted effects.

## 290 **Results**

### 291 *Sampling and molecular analyses*

292 In total, we collected 18,508 sand fly females, which were gathered in 666 pools used for sand  
293 fly metabarcoding and *Leishmania* detection (Table 1, Data S1). After DNA extraction and  
294 amplification, high-throughput sequencing and bioinformatic filtering, 600 (90.1%) sand fly  
295 pools could be characterized with metabarcoding, and *Leishmania* DNA was detected in 175  
296 (26.3%) of them. We further collected 855 blood-fed dipterans, including 715 sand flies, 123  
297 mosquitoes and 17 biting midges that were analyzed individually (Table S1). Dipteran  
298 identification was successful in 91.7% of the individuals (although not necessarily at the  
299 species level), and their blood meal content was identified in 75.9% of the cases. Both dipteran  
300 and blood meal identifications were successful for 602 (70.4%) individuals. Three sites in  
301 which less than five blood-fed dipterans could be identified were not retained for statistical  
302 analyses (Figure 1, Table 1).

### 303 *Sand fly, vertebrate and Leishmania identifications*

304 In total, we identified 63 sand fly MOTUs in sand fly pools (Table S1, Data S1), which is fairly  
305 consistent with the known sand fly species richness of French Guiana (about 80 species  
306 recorded so far). 34 of these were identified at the species level, including seven known or  
307 suspected *Leishmania* vector species in the region: *Bichromomyia flaviscutellata*, *Nyssomyia*  
308 *umbratilis*, *Psychodopygus ayrozai*, *Ps. panamensis*, *Ps. squamiventris maripaensis*,  
309 *Trichophoromyia ubiquitalis*, *Viannamyia furcata* (Rotureau, 2006). The most abundant  
310 species were *Ps. squamiventris maripaensis*, *Th. ininii*, *Trichopygomyia trichopyga* and *Th.*  
311 *ubiquitalis* (25.2%, 18.3%, 18.0% and 10.9% of the estimated number of individuals,

312 respectively). Five *Leishmania* species were detected in sand fly pools: *L. lainsoni*, *L.*  
313 *amazonensis*, *L. naiffi*, *L. braziliensis*, *L. guyanensis*; the most frequent being *L. lainsoni* and  
314 *L. naiffi* (48.6% and 43.4% of the positive samples respectively). Blood-fed specimens  
315 belonged to 51 dipteran MOTUs, and blood meal analyses revealed a total of 52 vertebrate  
316 MOTUs (Table 2, Figure S3, Data S1), including 28 mammals, among which 11 were  
317 recognized *Leishmania* reservoir hosts in the region: *Didelphis marsupialis*, *Philander*  
318 *opossum*, *Metachirus nudicaudatus*, *Choloepus didactylus*, *Dasypus novemcinctus*, *Tamandua*  
319 *tetradactyla*, *Coendou melanurus*, *C. prehensilis*, *Dasyprocta leporina*, *Proechymis cuvieri*  
320 and *P. guyannensis* (Rotureau, 2006). Our results revealed contrasting host preferences across  
321 sand fly species, and were consistent with existing knowledge [e.g. *Nyssomyia* spp. feeding  
322 mostly on sloth and other arboreal mammals (Christensen, Arias, de Vasquez, & de Freitas,  
323 1982), *B. flaviscutellata* feeding mostly on terrestrial rodents (R. Lainson & Shaw, 1968), and  
324 *Psychodopygus* spp. feeding mostly on armadillos (Le Pont, 1990); Table 2; Figure S3].

### 325 ***Statistical analyses***

326 We estimated mammal diversity (Shannon index) and the relative abundance of *Leishmania*  
327 reservoirs in each site based on dipteran blood meal identifications using a probabilistic  
328 approach (Figure S1, S2). We then used these estimates in a series of Bayesian regressions to  
329 investigate the effect of mammal diversity on the transmission of *Leishmania* parasites.  
330 MCMC trace plots and ESS values indicated convergence and correct sampling of the posterior  
331 distribution for all parameters (Figures S4, S5, S6). Mean estimates of the Shannon index of  
332 mammals across our study sites ranged from 0.46 to 2.7 (Table 1), and these correlated  
333 negatively with the human footprint index (HFP; (de Thoisy et al., 2010); mean effect: -0.34,  
334 90%CI [-0.61,-0.07]; Figure 1B). Spatial autocorrelation was not detected for the HFP



335 (expected/observed Moran's  $I$ : -0.067/0.056; p-value: 0.3), nor for the Shannon index of  
336 mammals (expected/observed Moran's  $I$ : -0.067/-0.13; p-value: 0.6). The Shannon index of  
337 mammals correlated negatively with the relative abundance of *Leishmania* reservoirs (-0.68,  
338 90%CI [-0.99,-0.38]; Figure 2A), positively with sand fly density (0.4, 90%CI [0.13,0.67];  
339 Figure 2B), and negatively (but weakly) with the Shannon index of sand flies (-0.19, 90%CI [-  
340 0.38,0]; Figure 4SA). Furthermore, we estimated that the prevalence rate of *Leishmania* in sand  
341 flies correlated positively with the relative abundance of *Leishmania* reservoirs (0.43, 90%CI  
342 [0.27,0.59]; Figure 2C), but did not correlate with sand fly density or diversity (0.03, 90%CI  
343 [-0.12,0.17] and 0.08, 90%CI [-0.07,0.23], respectively; Figure S7B,C). Finally, posterior  
344 predictions indicated that the Shannon index of mammals correlated negatively with the  
345 *Leishmania* prevalence rate in sand flies but did not correlate with the density of infected sand  
346 flies (Figure 2D,E).

## 347 **Discussion**

348 With this work, we show that wild vector-borne disease systems can be efficiently studied  
349 using DNA metabarcoding of arthropod vectors, which allows measuring arthropod, vertebrate  
350 and parasite communities altogether. Dipteran blood meal analyses allowed to identify a large  
351 variety of vertebrates across our study sites, confirming the potential of invertebrate-derived  
352 DNA (iDNA) approaches for biodiversity monitoring (Calvignac-Spencer et al., 2013; Kocher,  
353 de Thoisy, Catzeflis, Valière, et al., 2017; Schnell et al., 2015). Based on blood meal  
354 identifications, we estimated the Shannon index of mammals, which correlated negatively with  
355 the human footprint index (Figure 1B), as expected (de Thoisy et al., 2010). This suggests that  
356 the variation of mammal diversity across our study sites was, at least partly, linked to human-  
357 induced disturbance. We then explored the effect of mammal diversity on *Leishmania* disease

358 systems and transmission. In particular, we assessed the occurrence of a dilution effect  
359 potentially leading to reduced transmission with higher mammal diversity. In the case of  
360 vector-borne diseases, a dilution effect can be expected only if arthropod vectors feed on  
361 various host species, including some that are poorly competent for pathogen transmission. This  
362 is the case for Amazonian leishmaniasis, since sand fly vector species from which blood meals  
363 could be analyzed were observed to have fed on different hosts, including some that are not  
364 known as *Leishmania* reservoirs (Table 2, Figure S3). For example, while most *Ny. umbratilis*  
365 individuals had fed on two-toed sloths (main vector and reservoir of *L. guyanensis*,  
366 respectively), and most *Psychodopygus* spp. had fed on nine-banded armadillos (main vector  
367 and reservoirs of *L. naiffi*), a significant proportion of them had fed, respectively, on different  
368 primates and large terrestrial mammals which are not known to act as *Leishmania* reservoirs.  
369 Therefore, *Leishmania* parasites may indeed end up in “diluting” hosts which could contribute  
370 to reduced transmission of these disease agents. This was further supported by the observation  
371 of a positive correlation between the relative abundance of *Leishmania* reservoir hosts and  
372 *Leishmania* prevalence rate in sand flies (Figure 2C).

373 Another important assumption of the dilution effect hypothesis is that species contributing the  
374 most to pathogen transmission dominate in disturbed and less diverse communities. Ecological  
375 and evolutionary hypotheses have suggested the existence of such a positive relationship  
376 between host competence for pathogens and resilience to disturbance, leading to a general  
377 increase of the overall community competence with biodiversity loss (Johnson, Ostfeld, &  
378 Keesing, 2015; Keesing et al., 2010; Ostfeld & Keesing, 2012). Species that are resilient to  
379 changing environments are frequently characterized by fast life history strategies, including  
380 low investment in adaptive immunity and high reproductive rate, yielding an important influx

381 of susceptible individuals in the population. In addition, pathogens may adapt predominantly  
382 to resilient host species, because these hosts are generally widespread, mobile and abundant,  
383 therefore constituting the most frequently encountered resource. Empirical evidence has  
384 supported these ideas (García-Peña et al., 2016; Han, Schmidt, Bowden, & Drake, 2015;  
385 Johnson et al., 2019; Johnson, Preston, Hoverman, & Richgels, 2013; Johnson et al., 2012),  
386 although it seems that the situation may vary depending on the taxa under consideration (Gibb  
387 et al., 2018; H. Young, Griffin, Wood, & Nunn, 2013). Here, we show that mammal species  
388 known as *Leishmania* reservoirs indeed dominate less diverse mammal communities, with a  
389 233.6% (90%CI: [293.0%,190.8%]) increase of their relative abundance along the range of  
390 estimated mammal's Shannon index (Figure 2A). In sum, these results suggest a predictable  
391 effect of local mammal diversity changes on the overall host community competence for  
392 *Leishmania* parasites, contributing to an increase of vectorial transmission with decreasing  
393 mammal diversity, in accordance with the dilution effect hypothesis.

394 However, arthropod vector ecology should also be accounted for when investigating the impact  
395 of biodiversity changes on a vector-borne disease. In particular, higher vector density should  
396 be associated with more frequent host-pathogen contacts and increase transmission (Smith et  
397 al., 2012). Additionally, when several arthropod species can act as vectors for a given  
398 pathogen, higher arthropod diversity can result in higher pathogen transmission through an  
399 overall increase of vector abundance, or due to functional complementarity between vector  
400 species (Park et al., 2016; Roche et al., 2013). Given that vertebrates constitute trophic  
401 resources for blood-feeding arthropods, it can be expected that arthropod communities are  
402 partially driven by the abundance and composition of local vertebrate fauna. This might,  
403 however, depend on the ecology of the considered arthropod species. For instance, highly

404 mobile mosquitoes might be little affected by changes in host density, which is a classical  
405 assumption in epidemiological models (Dobson, 2004), while ticks may be more sensitive to  
406 the presence of suitable hosts in their immediate environment (Ogden & Tsao, 2009; Randolph  
407 & Dobson, 2012; Swei et al., 2011; Titcomb et al., 2017). Little is known in this regard for  
408 sand flies, which, despite being flying insects, have relatively small flight ranges (Casanova,  
409 Costa, & Natal, 2005; Morrison, Ferro, Morales, Tesh, & Wilson, 1993). Here, we observed a  
410 positive relationship between mammal diversity and sand fly density (Figure 2B), which  
411 suggests that sand flies may indeed be affected by changes in local mammal communities.  
412 Such a relationship could be mediated by a correlation between mammal diversity and overall  
413 mammal biomass, resulting in greater availability of blood meal resources for sand flies in  
414 more diverse ecosystems. However, our data does not provide information regarding the  
415 absolute abundance of mammals, and it is difficult draw conclusions about the causality of  
416 such a relationship since some environmental factors could affect both mammal and arthropod  
417 species communities. On the other hand, our results point to a negative correlation between  
418 mammal and sand fly diversity (although weakly significant; effect 90%CI: [-0.38%,0.00%];  
419 Figure S7A), which suggest that different factors might shape mammal and sand fly  
420 communities. This further highlights the need for more research to understand the factors  
421 shaping sand fly assemblages in sylvatic systems and their potential consequences for  
422 *Leishmania* transmission.

423 Our results did not reveal an effect of sand fly density or diversity on the prevalence rate of  
424 *Leishmania* parasites in sand flies (Figure S7B, C). Thus, the proportion of *Leishmania*  
425 reservoir hosts in mammal communities appeared as the main driver of *Leishmania*  
426 transmission, resulting in a negative effect of mammal diversity on *Leishmania* prevalence

427 rate, through host dilution (Figure 2D). However, the prevalence of a pathogen in vectors is  
428 not necessarily a relevant measure of disease transmission, and the density of infected vectors  
429 should rather be considered for this matter. For a given prevalence rate, higher vector density  
430 should be associated with a higher density of infected vectors. Therefore, the positive  
431 relationship observed between mammal diversity and sand fly density across our study sites  
432 might act on leishmaniasis transmission in opposition to the dilution effect, which it may  
433 attenuate, cancel, or even reverse. This shows that the alteration of mammal diversity is  
434 associated with changes in the ecosystem that independently impact *Leishmania* transmission  
435 in contrasting ways. Posterior predictions of the density of infected sand flies indicate a weak  
436 overall impact of mammal diversity changes on *Leishmania* transmission (Figure 2E).  
437 Therefore, it seems that the observed dilution and amplification effects compensate for each  
438 other in the system studied here. However, one or the other could predominate in other contexts  
439 depending on characteristics of the considered system and environmental conditions that need  
440 to be further determined. This constitutes additional evidence that biodiversity changes may  
441 impact vector-borne pathogen transmission through concurrent mechanisms, and further  
442 stresses the importance of better accounting for arthropod vector ecology in biodiversity-  
443 disease research.

#### 444 **Acknowledgements**

445 This work was supported by “Investissements d’Avenir” and other grants managed by the  
446 French National Research Agency (LABEX CEBA: ANR-10-LABX-25-01; LABEX TULIP:  
447 ANR-10-LABX-41, AnaEE-France; ANR-11-INBS-0001; LABEX DRIIHM: ANR-11-  
448 LABX-0010-DRIIHM), as well as the RESERVOIRS projects (via ERDF and CNES funds)  
449 and the Inkermann fund hosted at the Fondation de France. J.C. is supported by Marie-Curie

450 Actions People H2020, MSCAIF-EF-ST-708207. J.-F.G. is supported by a US NSF-NIH  
451 Ecology of infectious diseases award (#191145). We thank the Nouragues research field station  
452 and the Trésor natural reserve for logistical support, F. Catzefflis for his involvement in the  
453 creation of the project, S. Talaga for his contribution to sand fly sampling, B. Roche, J.B. Ferdy  
454 and T. Vergne for their advices on data analyses, F. Boyer and the LECA laboratory (UMR  
455 5553) for their help in bioinformatics, the LIVH and Parasitology team of the Pasteur Institute  
456 of French Guiana for their support during laboratory work, and M. Michel for English  
457 proofreading.

## 458 **References**

- 459 Bañuls, A.-L., Hide, M., & Prugnonle, F. (2007). *Leishmania* and the leishmaniasis: A parasite genetic update and  
460 advances in taxonomy, epidemiology and pathogenicity in humans. *Advances in Parasitology*, *64*, 1–109.
- 461 Binladen, J., Gilbert, M. T. P., Bollback, J. P., Panitz, F., Bendixen, C., Nielsen, R., & Willerslev, E. (2007). The use  
462 of coded PCR primers enables high-throughput sequencing of multiple homolog amplification products by  
463 454 parallel sequencing. *PLOS One*, *2*(2), e197. doi: 10.1371/journal.pone.0000197
- 464 Boyer, F., Mercier, C., Bonin, A., Le Bras, Y., Taberlet, P., & Coissac, E. (2016). obitools: A unix-inspired software  
465 package for DNA metabarcoding. *Molecular Ecology Resources*, *16*(1), 176–182.
- 466 Calvignac-Spencer, S., Merkel, K., Kutzner, N., Kühl, H., Boesch, C., Kappeler, P. M., ... Leendertz, F. H. (2013).  
467 Carrion fly-derived DNA as a tool for comprehensive and cost-effective assessment of mammalian  
468 biodiversity. *Molecular Ecology*, *22*(4), 915–924. doi: 10.1111/mec.12183
- 469 Cardinale, B. J., Duffy, J. E., Gonzalez, A., Hooper, D. U., Perrings, C., Venail, P., ... Naeem, S. (2012). Biodiversity  
470 loss and its impact on humanity. *Nature*, *486*(7401), 59–67. doi: 10.1038/nature11148
- 471 Carlson, J. C., Dyer, L. A., Omlin, F. X., & Beier, J. C. (2009). Diversity cascades and malaria vectors. *Journal of*  
472 *Medical Entomology*, *46*(3), 460.
- 473 Carpenter, B., Gelman, A., Hoffman, M., Lee, D., Goodrich, B., Betancourt, M., ... Riddell, A. (2017). Stan: A  
474 probabilistic programming language. *J Stat Softw*, *76*(1), 1–32. (a). doi: 10.18637/jss.v076.i01
- 475 Casanova, C., Costa, A. I., & Natal, D. (2005). Dispersal pattern of the sand fly *Lutzomyia neivai* (Diptera:  
476 Psychodidae) in a cutaneous leishmaniasis endemic rural area in Southeastern Brazil. *Memórias Do Instituto*  
477 *Oswaldo Cruz*, *100*(7), 719–724. doi: 10.1590/S0074-02762005000700006
- 478 Casquet, J., Thebaud, C., & Gillespie, R. G. (2012). Chelex without boiling, a rapid and easy technique to obtain stable  
479 amplifiable DNA from small amounts of ethanol-stored spiders. *Molecular Ecology Resources*, *12*(1), 136–  
480 141. doi: 10.1111/j.1755-0998.2011.03073.x
- 481 Chavy, A., Nava, A. F. D., Luz, S. L. B., Ramírez, J. D., Herrera, G., Santos, T. V. dos, ... Thoisy, B. de. (2019).  
482 Ecological niche modelling for predicting the risk of cutaneous leishmaniasis in the Neotropical moist forest  
483 biome. *PLOS Neglected Tropical Diseases*, *13*(8), e0007629. doi: 10.1371/journal.pntd.0007629

- 484 Christensen, H. A., Arias, J. R., de Vasquez, A. M., & de Freitas, R. A. (1982). Hosts of sandfly vectors of *Leishmania*  
485 *braziliensis guyanensis* in the central Amazon of Brazil. *The American Journal of Tropical Medicine and*  
486 *Hygiene*, 31(2), 239–242.
- 487 Civitello, D. J., Cohen, J., Fatima, H., Halstead, N. T., Liriano, J., McMahon, T. A., ... Rohr, J. R. (2015). Biodiversity  
488 inhibits parasites: Broad evidence for the dilution effect. *Proceedings of the National Academy of Sciences*,  
489 112(28), 8667–8671. doi: 10.1073/pnas.1506279112
- 490 Clarke, L. J., Soubrier, J., Weyrich, L. S., & Cooper, A. (2014). Environmental metabarcodes for insects: *In silico*  
491 PCR reveals potential for taxonomic bias. *Molecular Ecology Resources*, 14(6), 1160–1170. doi:  
492 10.1111/1755-0998.12265
- 493 Clay, C. A., Lehmer, E. M., Jeor, S. St., & Dearing, M. D. (2009). Sin Nombre Virus and Rodent Species Diversity:  
494 A Test of the Dilution and Amplification Hypotheses. *PLoS ONE*, 4(7), e6467. doi:  
495 10.1371/journal.pone.0006467
- 496 Clay, C. A., Lehmer, E. M., St. Jeor, S., & Dearing, M. D. (2009). Testing Mechanisms of the Dilution Effect: Deer  
497 Mice Encounter Rates, Sin Nombre Virus Prevalence and Species Diversity. *EcoHealth*, 6(2), 250–259. doi:  
498 10.1007/s10393-009-0240-2
- 499 Cohen, J. M., Civitello, D. J., Brace, A. J., Feichtinger, E. M., Ortega, C. N., Richardson, J. C., ... Rohr, J. R. (2016).  
500 Spatial scale modulates the strength of ecological processes driving disease distributions. *Proceedings of the*  
501 *National Academy of Sciences*, 133(24), E3359–E3364. doi: 10.1073/pnas.1521657113
- 502 de Thoisy, B., Richard-Hansen, C., Goguillon, B., Joubert, P., Obstancias, J., Winterton, P., & Brosse, S. (2010).  
503 Rapid evaluation of threats to biodiversity: Human footprint score and large vertebrate species responses in  
504 French Guiana. *Biodiversity and Conservation*, 19(6), 1567–1584. doi: 10.1007/s10531-010-9787-z
- 505 Derne, B. T., Fearnley, E. J., Lau, C. L., Paynter, S., & Weinstein, P. (2011). Biodiversity and leptospirosis risk: A  
506 case of pathogen regulation? *Medical Hypotheses*, 77(3), 339–344. doi: 10.1016/j.mehy.2011.05.009
- 507 Dobson, A. (2004). Population Dynamics of Pathogens with Multiple Host Species. *The American Naturalist*, 164(s5),  
508 64–78. doi: 10.1086/424681
- 509 Esling, P., Lejzerowicz, F., & Pawlowski, J. (2015). Accurate multiplexing and filtering for high-throughput  
510 amplicon-sequencing. *Nucleic Acids Research*, 43(5), 2513–2524. doi: 10.1093/nar/gkv107
- 511 Ezenwa, V. O., Godsey, M. S., King, R. J., & Guphill, S. C. (2006). Avian diversity and West Nile virus: Testing  
512 associations between biodiversity and infectious disease risk. *Proceedings of the Royal Society B: Biological*  
513 *Sciences*, 273(1582), 109–117. doi: 10.1098/rspb.2005.3284
- 514 Faust, C. L., Dobson, A. P., Gottdenker, N., Bloomfield, L. S. P., McCallum, H. I., Gillespie, T. R., ... Plowright, R.  
515 K. (2017). Null expectations for disease dynamics in shrinking habitat: Dilution or amplification? *Phil. Trans.*  
516 *R. Soc. B*, 372(1722), 20160173. doi: 10.1098/rstb.2016.0173
- 517 García-Peña, G. E., Garchitorena, A., Carolan, K., Canard, E., Prieur-Richard, A.-H., Suzán, G., ... Guégan, J.-F.  
518 (2016). Niche-based host extinction increases prevalence of an environmentally acquired pathogen. *Oikos*,  
519 125(10), 1508–1515. doi: 10.1111/oik.02700
- 520 Gibb, R., Redding, D. W., Chin, K. Q., Blackburn, T. M., Newbold, T., & Jones, K. E. (2018). Effects of land use on  
521 zoonotic host communities: A global correlative analysis. *The Lancet Planetary Health*, 2, S2. doi:  
522 10.1016/S2542-5196(18)30087-1
- 523 Gilbert, L., Norman, R., Laurenson, K. M., Reid, H. W., & Hudson, P. J. (2001). Disease persistence and apparent  
524 competition in a three-host community: An empirical and analytical study of large-scale, wild populations.  
525 *Journal of Animal Ecology*, 70(6), 1053–1061. doi: 10.1046/j.0021-8790.2001.00558.x
- 526 Gottdenker, N. L., Chaves, L. F., Calzada, J. E., Saldaña, A., & Carroll, C. R. (2012). Host Life History Strategy,  
527 Species Diversity, and Habitat Influence *Trypanosoma cruzi* Vector Infection in Changing Landscapes.  
528 *PLoS Negl Trop Dis*, 6(11), e1884. doi: 10.1371/journal.pntd.0001884

- 529 Halliday, F. W., & Rohr, J. R. (2019). Measuring the shape of the biodiversity-disease relationship across systems  
530 reveals new findings and key gaps. *Nature Communications*, *10*(1), 1–10. doi: 10.1038/s41467-019-13049-  
531 w
- 532 Halliday, F. W., Rohr, J. R., & Laine, A.-L. (2020). Biodiversity loss underlies the dilution effect of biodiversity.  
533 *Ecology Letters*, *23*(11), 1611–1622. doi: 10.1111/ele.13590
- 534 Halsey, S. (2019). Defuse the dilution effect debate. *Nature Ecology & Evolution*, *3*, 145–146. doi: 10.1038/s41559-  
535 018-0764-3
- 536 Hamer, G. L., Chaves, L. F., Anderson, T. K., Kitron, U. D., Brawn, J. D., Ruiz, M. O., ... Goldberg, T. L. (2011).  
537 Fine-scale variation in vector host use and force of infection drive localized patterns of west nile virus  
538 transmission. *PLOS One*, *6*(8), e23767. doi: 10.1371/journal.pone.0023767
- 539 Han, B. A., Schmidt, J. P., Bowden, S. E., & Drake, J. M. (2015). Rodent reservoirs of future zoonotic diseases.  
540 *Proceedings of the National Academy of Sciences*, *112*(22), 7039–7044. doi: 10.1073/pnas.1501598112
- 541 Hsieh, T. C., Ma, K. H., & Chao, A. (2016). iNEXT: An R package for rarefaction and extrapolation of species  
542 diversity (Hill numbers). *Methods in Ecology and Evolution*, *7*(12), 1451–1456. doi: 10.1111/2041-  
543 210X.12613
- 544 Huang, Z. Y. X., van Langevelde, F., Prins, H. H. T., & de Boer, W. F. (2015). Dilution versus facilitation: Impact of  
545 connectivity on disease risk in metapopulations. *Journal of Theoretical Biology*, *376*, 66–73. doi:  
546 10.1016/j.jtbi.2015.04.005
- 547 Johnson, P. T. J., Calhoun Dana M., Riepe Tawni, McDevitt-Galles Travis, & Koprivnikar Janet. (2019). Community  
548 disassembly and disease: Realistic—but not randomized—biodiversity losses enhance parasite transmission.  
549 *Proceedings of the Royal Society B: Biological Sciences*, *286*(1902), 20190260. doi: 10.1098/rspb.2019.0260
- 550 Johnson, P. T. J., de Roode, J. C., & Fenton, A. (2015). Why infectious disease research needs community ecology.  
551 *Science*, *349*(6252), 1259504–1259504. doi: 10.1126/science.1259504
- 552 Johnson, P. T. J., Hartson, R. B., Larson, D. J., & Sutherland, D. R. (2008). Diversity and disease: Community  
553 structure drives parasite transmission and host fitness. *Ecology Letters*, *11*(10), 1017–1026. doi:  
554 10.1111/j.1461-0248.2008.01212.x
- 555 Johnson, P. T. J., Lund, P. J., Hartson, R. B., & Yoshino, T. P. (2009). Community diversity reduces *Schistosoma*  
556 *mansonii* transmission, host pathology and human infection risk. *Proceedings of the Royal Society B:*  
557 *Biological Sciences*, *276*(1662), 1657–1663. doi: 10.1098/rspb.2008.1718
- 558 Johnson, P. T. J., Ostfeld, R. S., & Keesing, F. (2015). Frontiers in research on biodiversity and disease. *Ecology*  
559 *Letters*, *18*(10), 1119–1133. doi: 10.1111/ele.12479
- 560 Johnson, P. T. J., Preston, D. L., Hoverman, J. T., & Richgels, K. L. D. (2013). Biodiversity decreases disease through  
561 predictable changes in host community competence. *Nature*, *494*(7436), 230–233. doi: 10.1038/nature11883
- 562 Johnson, P. T. J., Rohr, J. R., Hoverman, J. T., Kellermanns, E., Bowerman, J., & Lunde, K. B. (2012). Living fast  
563 and dying of infection: Host life history drives interspecific variation in infection and disease risk: Living  
564 fast and dying of infection. *Ecology Letters*, *15*(3), 235–242. doi: 10.1111/j.1461-0248.2011.01730.x
- 565 Keesing, F., Belden, L. K., Daszak, P., Dobson, A., Harvell, C. D., Holt, R. D., ... Ostfeld, R. S. (2010). Impacts of  
566 biodiversity on the emergence and transmission of infectious diseases. *Nature*, *468*(7324), 647–652. doi:  
567 10.1038/nature09575
- 568 Keesing, F., & Ostfeld, R. S. (2021). Impacts of biodiversity and biodiversity loss on zoonotic diseases. *Proceedings*  
569 *of the National Academy of Sciences*, *118*(17). doi: 10.1073/pnas.2023540118
- 570 Kocher, A., de Thoisy, B., Catzeflis, F., Huguin, M., Valiere, S., Zinger, L., ... Murienne, J. (2017). Evaluation of  
571 short mitochondrial metabarcodes for the identification of Amazonian mammals. *Methods in Ecology and*  
572 *Evolution*, *8*, 1276–1283. doi: 10.1111/2041-210X.12729
- 573 Kocher, A., de Thoisy, B., Catzeflis, F., Valière, S., Bañuls, A.-L., & Murienne, J. (2017). iDNA screening: Disease  
574 vectors as vertebrate samplers. *Molecular Ecology*, *26*(22), 6478–6486. doi: 10.1111/mec.14362



- 575 Kocher, A., Gantier, J.-C., Gaborit, P., Zinger, L., Holota, H., Valiere, S., ... Murienne, J. (2017). Vector soup: High-  
576 throughput identification of Neotropical phlebotomine sand flies using metabarcoding. *Molecular Ecology*  
577 *Resources*, 17(2), 172–182. doi: 10.1111/1755-0998.12556
- 578 Kocher, A., Valière, S., Bañuls, A.-L., & Murienne, J. (2018). High-throughput sequencing of kDNA amplicons for  
579 the analysis of *Leishmania* minicircles and identification of Neotropical species. *Parasitology*, 145(5), 585–  
580 594. doi: 10.1017/S0031182017002013
- 581 Lafferty, K. D., & Wood, C. L. (2013). It's a myth that protection against disease is a strong and general service of  
582 biodiversity conservation: Response to Ostfeld and Keesing. *Trends in Ecology & Evolution*, 28(9), 503–  
583 504. doi: 10.1016/j.tree.2013.06.012
- 584 Lainson, R., & Shaw, J. J. (1968). Leishmaniasis in Brazil: I. Observations on enzootic rodent leishmaniasis—  
585 incrimination of *Lutzomyia flaviscutellata* as the vector in the lower amazonian basin. *Transactions of the*  
586 *Royal Society of Tropical Medicine and Hygiene*, 62(3), 396–405.
- 587 Lainson, Ralph, & Shaw, J. J. (2010). New world leishmaniasis. In B. W. J. Mahy & S. P. Meulen (Eds.), *Topley &*  
588 *Wilson's Microbiology and Microbial Infections*. Wiley. Retrieved from  
589 <http://onlinelibrary.wiley.com/doi/10.1002/9780470688618.taw0182/full>
- 590 Laporta, G. Z., Prado, P. I. K. L. de, Kraenkel, R. A., Coutinho, R. M., & Sallum, M. A. M. (2013). Biodiversity Can  
591 Help Prevent Malaria Outbreaks in Tropical Forests. *PLoS Neglected Tropical Diseases*, 7(3), e2139. doi:  
592 10.1371/journal.pntd.0002139
- 593 Le Pont, F. (1990). Attraction of the armadillo (*Dasypus novemcinctus*) and guinea pigs for phlebotomines in French  
594 Guiana. *Bulletin de La Societe de Pathologie Exotique*, 83(5), 671–676.
- 595 Levi, T., Massey, A. L., Holt, R. D., Keesing, F., Ostfeld, R. S., & Peres, C. A. (2016). Does biodiversity protect  
596 humans against infectious disease? Comment. *Ecology*, 97(2), 536–542.
- 597 LoGiudice, K., Ostfeld, R. S., Schmidt, K. A., & Keesing, F. (2003). The ecology of infectious disease: Effects of  
598 host diversity and community composition on Lyme disease risk. *Proceedings of the National Academy of*  
599 *Sciences*, 100(2), 567–571.
- 600 Loss, S. R., Hamer, G. L., Walker, E. D., Ruiz, M. O., Goldberg, T. L., Kitron, U. D., & Brawn, J. D. (2009). Avian  
601 host community structure and prevalence of West Nile virus in Chicago, Illinois. *Oecologia*, 159(2), 415–  
602 424. doi: 10.1007/s00442-008-1224-6
- 603 Luis, A. D., Kuenzi, A. J., & Mills, J. N. (2018). Species diversity concurrently dilutes and amplifies transmission in  
604 a zoonotic host–pathogen system through competing mechanisms. *Proceedings of the National Academy of*  
605 *Sciences*, 115(31), 7979–7984. doi: 10.1073/pnas.1807106115
- 606 McGregor, B. L., Stenn, T., Saylor, K. A., Blosser, E. M., Blackburn, J. K., Wisely, S. M., & Burkett-Cadena, N. D.  
607 (2018). Host use patterns of *Culicoides* spp. Biting midges at a big game preserve in Florida, U.S.A., and  
608 implications for the transmission of orbiviruses. *Medical and Veterinary Entomology*, 33(1), 110–120. doi:  
609 10.1111/mve.12331
- 610 Mercier, C., Boyer, F., Bonin, A., & Coissac, E. (2013). SUMATRA and SUMACLUSt: Fast and exact comparison  
611 and clustering of sequences. *Programs and Abstracts of the SeqBio 2013 Workshop*, 27–29. Retrieved from  
612 <http://www.gdr-bim.cnrs.fr/seqbio2013/wp-content/uploads/2013/12/seqbio2013-actes.pdf#page=28>
- 613 Mihaljevic, J. R., Joseph, M. B., Orlofske, S. A., & Paull, S. H. (2014). The Scaling of Host Density with Richness  
614 Affects the Direction, Shape, and Detectability of Diversity-Disease Relationships. *PLOS ONE*, 9(5), e97812.  
615 doi: 10.1371/journal.pone.0097812
- 616 Miller, E., & Huppert, A. (2013). The Effects of Host Diversity on Vector-Borne Disease: The Conditions under  
617 Which Diversity Will Amplify or Dilute the Disease Risk. *PLOS One*, 8(11), e80279. doi:  
618 10.1371/journal.pone.0080279
- 619 Morand, S., Jittapalpong, S., Suputtamongkol, Y., Abdullah, M. T., & Huan, T. B. (2014). Infectious Diseases and  
620 Their Outbreaks in Asia-Pacific: Biodiversity and Its Regulation Loss Matter. *PLOS One*, 9(2), e90032. doi:  
621 10.1371/journal.pone.0090032

- 622 Morrison, A. C., Ferro, C., Morales, A., Tesh, R. B., & Wilson, M. L. (1993). Dispersal of the sand fly *Lutzomyia*  
623 *longipalpis* (Diptera: Psychodidae) at an endemic focus of visceral leishmaniasis in Colombia. *Journal of*  
624 *Medical Entomology*, *30*(2), 427–435.
- 625 Norman, R., Bowers, R. G., Begon, M., & Hudson, P. J. (1999). Persistence of Tick-borne Virus in the Presence of  
626 Multiple Host Species: Tick Reservoirs and Parasite Mediated Competition. *Journal of Theoretical Biology*,  
627 *200*(1), 111–118. doi: 10.1006/jtbi.1999.0982
- 628 Ogden, N. H., & Tsao, J. I. (2009). Biodiversity and Lyme disease: Dilution or amplification? *Epidemics*, *1*(3), 196–  
629 206. doi: 10.1016/j.epidem.2009.06.002
- 630 Oliver, T. H., Heard, M. S., Isaac, N. J. B., Roy, D. B., Procter, D., Eigenbrod, F., ... Bullock, J. M. (2015).  
631 Biodiversity and Resilience of Ecosystem Functions. *Trends in Ecology & Evolution*, *30*(11), 673–684. doi:  
632 10.1016/j.tree.2015.08.009
- 633 Ostfeld, R. S., & Keesing, F. (2000). Biodiversity and disease risk: The case of Lyme disease. *Conservation Biology*,  
634 *14*(3), 722–728.
- 635 Ostfeld, R. S., & Keesing, F. (2012). Effects of Host Diversity on Infectious Disease. *Annual Review of Ecology,*  
636 *Evolution, and Systematics*, *43*(1), 157–182. doi: 10.1146/annurev-ecolsys-102710-145022
- 637 Ostfeld, R. S., & LoGiudice, K. (2003). Community disassembly, biodiversity loss, and the erosion of an ecosystem  
638 service. *Ecology*, *84*(6), 1421–1427.
- 639 Pan American Health Organization. (2019). *Leishmaniasis: Epidemiological Report in the Americas*. Washington.  
640 Retrieved from <https://www.paho.org/en/topics/leishmaniasis>
- 641 Paradis, E., Claude, J., & Strimmer, K. (2004). APE: Analyses of Phylogenetics and Evolution in R language.  
642 *Bioinformatics (Oxford, England)*, *20*(2), 289–290.
- 643 Park, A. W., Cleveland, C. A., Dallas, T. A., & Corn, J. L. (2016). Vector species richness increases haemorrhagic  
644 disease prevalence through functional diversity modulating the duration of seasonal transmission.  
645 *Parasitology*, *143*(7), 874–879. doi: 10.1017/S0031182015000578
- 646 R Core team. (2020). *R: A language and environment for statistical computing*. Vienna, Austria: R Foundation for  
647 Statistical Computing. Retrieved from <https://www.R-project.org>
- 648 Randolph, S. E., & Dobson, A. D. M. (2012). Pangloss revisited: A critique of the dilution effect and the biodiversity-  
649 buffers-disease paradigm. *Parasitology*, *139*(07), 847–863. doi: 10.1017/S0031182012000200
- 650 Rangel, E. F., Costa, S. M., & Carvalho, B. M. (2014). Environmental changes and the geographic spreading of  
651 American cutaneous leishmaniasis in Brazil. In D. M. Claborn (Ed.), *Leishmaniasis—trends in epidemiology,*  
652 *diagnosis and treatment*. (pp. 1–25). Rijeka: InTech.
- 653 Riaz, T., Shehzad, W., Viari, A., Pompanon, F., Taberlet, P., & Coissac, E. (2011). ecoPrimers: Inference of new  
654 DNA barcode markers from whole genome sequence analysis. *Nucleic Acids Research*, *39*(21), 1–11. doi:  
655 10.1093/nar/gkr732
- 656 Roche, B., Dobson, A. P., Guégan, J.-F., & Rohani, P. (2012). Linking community and disease ecology: The impact  
657 of biodiversity on pathogen transmission. *Philosophical Transactions of the Royal Society B: Biological*  
658 *Sciences*, *367*(1604), 2807–2813. doi: 10.1098/rstb.2011.0364
- 659 Roche, B., & Guégan, J.-F. (2011). Ecosystem dynamics, biological diversity and emerging infectious diseases.  
660 *Comptes Rendus Biologies*, *334*(5–6), 385–392. doi: 10.1016/j.crv.2011.02.008
- 661 Roche, B., Rohani, P., Dobson, A. P., & Guégan, J.-F. (2013). The Impact of Community Organization on Vector-  
662 Borne Pathogens. *The American Naturalist*, *181*(1), 1–11. doi: 10.1086/668591
- 663 Rohr, J. R., Civitello, D. J., Crumrine, P. W., Halstead, N. T., Miller, A. D., Schotthoefer, A. M., ... Beasley, V. R.  
664 (2015). Predator diversity, intraguild predation, and indirect effects drive parasite transmission. *Proceedings*  
665 *of the National Academy of Sciences*, *112*(10), 3008–3013. doi: 10.1073/pnas.1415971112

- 666 Rohr, J. R., Civitello, D. J., Halliday, F. W., Hudson, P. J., Lafferty, K. D., Wood, C. L., & Mordecai, E. A. (2020).  
667 Towards common ground in the biodiversity–disease debate. *Nature Ecology & Evolution*, 4(1), 24–33. doi:  
668 10.1038/s41559-019-1060-6
- 669 Rotureau, B. (2006). Ecology of the *Leishmania* species in the Guianan ecoregion complex. *The American Journal of*  
670 *Tropical Medicine and Hygiene*, 74(1), 81–96.
- 671 Rudolf, V. H. W., & Antonovics, J. (2005). Species Coexistence and Pathogens with Frequency-Dependent  
672 Transmission. *The American Naturalist*, 166(1), 112–118. doi: 10.1086/430674
- 673 Salkeld, D. J., Padgett, K. A., & Jones, J. H. (2013). A meta-analysis suggesting that the relationship between  
674 biodiversity and risk of zoonotic pathogen transmission is idiosyncratic. *Ecology Letters*, 16(5), 679–686.  
675 doi: 10.1111/ele.12101
- 676 Schnell, I. B., Sollmann, R., Calvignac-Spencer, S., Siddall, M. E., Yu, D. W., Wilting, A., & Gilbert, M. Thomas. P.  
677 (2015). IDNA from terrestrial haematophagous leeches as a wildlife surveying and monitoring tool –  
678 prospects, pitfalls and avenues to be developed. *Frontiers in Zoology*, 12, 24. doi: 10.1186/s12983-015-0115-  
679 z
- 680 Smith, D. L., Battle, K. E., Hay, S. I., Barker, C. M., Scott, T. W., & McKenzie, F. E. (2012). Ross, Macdonald, and  
681 a Theory for the Dynamics and Control of Mosquito-Transmitted Pathogens. *PLOS Pathogens*, 8(4),  
682 e1002588. doi: 10.1371/journal.ppat.1002588
- 683 Suzán, G., Marcé, E., Giermakowski, J. T., Mills, J. N., Ceballos, G., Ostfeld, R. S., ... Yates, T. L. (2009).  
684 Experimental Evidence for Reduced Rodent Diversity Causing Increased Hantavirus Prevalence. *PLoS ONE*,  
685 4(5), e5461. doi: 10.1371/journal.pone.0005461
- 686 Swei, A., Ostfeld, R. S., Lane, R. S., & Briggs, C. J. (2011). Impact of the experimental removal of lizards on Lyme  
687 disease risk. *Proceedings of the Royal Society of London B: Biological Sciences*, 278(1720), 2970–2978. doi:  
688 10.1098/rspb.2010.2402
- 689 Talaga, S., Leroy, C., Guidez, A., Dusfour, I., Girod, R., Dejean, A., & Murienne, J. (2017). DNA reference libraries  
690 of French Guianese mosquitoes for barcoding and metabarcoding. *PLOS One*, 12(6), e0176993. doi:  
691 10.1371/journal.pone.0176993
- 692 Telfer, S., Bown, K. J., Sekules, R., Begon, M., Hayden, T., & Birtles, R. (2005). Disruption of a host-parasite system  
693 following the introduction of an exotic host species. *Parasitology*, 130(06), 661–668. doi:  
694 10.1017/S0031182005007250
- 695 Titcomb, G., Allan, B. F., Ainsworth, T., Henson, L., Hedlund, T., Pringle, R. M., ... Young, H. S. (2017). Interacting  
696 effects of wildlife loss and climate on ticks and tick-borne disease. *Proc. R. Soc. B*, 284(1862), 20170475.  
697 doi: 10.1098/rspb.2017.0475
- 698 Van Buskirk, J., & Ostfeld, R. S. (1995). Controlling Lyme Disease by Modifying the Density and Species  
699 Composition of Tick Hosts. *Ecological Applications*, 5(4), 1133–1140. doi: 10.2307/2269360
- 700 Vinson, J. E., & Park, A. W. (2019). Vector-borne parasite invasion in communities across space and time.  
701 *Proceedings of the Royal Society B: Biological Sciences*, 286(1917), 20192614. doi: 10.1098/rspb.2019.2614
- 702 Weinstein, S., Titcomb, G., Agwanda, B., Riginos, C., & Young, H. (2017). Parasite responses to large mammal loss  
703 in an African savanna. *Ecology*, 98(7), 1839–1848. doi: 10.1002/ecy.1858
- 704 Wood, C. L., & Lafferty, K. D. (2013). Biodiversity and disease: A synthesis of ecological perspectives on Lyme  
705 disease transmission. *Trends in Ecology & Evolution*, 28(4), 239–247. doi: 10.1016/j.tree.2012.10.011
- 706 Wood, C. L., Lafferty, K. D., DeLeo, G., Young, H. S., Hudson, P. J., & Kuris, A. M. (2014). Does biodiversity protect  
707 humans against infectious disease? *Ecology*, 95(4), 817–832.
- 708 Wood, C. L., Summerside, M., & Johnson, P. T. J. (2020). How host diversity and abundance affect parasite infections:  
709 Results from a whole-ecosystem manipulation of bird activity. *Biological Conservation*, 248, 108683. doi:  
710 10.1016/j.biocon.2020.108683

- 711 Young, H., Griffin, R. H., Wood, C. L., & Nunn, C. L. (2013). Does habitat disturbance increase infectious disease  
712 risk for primates? *Ecology Letters*, *16*(5), 656–663. doi: 10.1111/ele.12094
- 713 Young, H. S., Parker, I. M., Gilbert, G. S., Sofia Guerra, A., & Nunn, C. L. (2017). Introduced Species, Disease  
714 Ecology, and Biodiversity–Disease Relationships. *Trends in Ecology & Evolution*, *32*(1), 41–54. doi:  
715 10.1016/j.tree.2016.09.008

## 716 **Data Accessibility and Benefit-Sharing**

717 Sample metadata: Data and code used for statistical analyses are available as supplementary  
718 material.

719 Genetic data: Sequencing data has been deposited at the Dryad database (doi:  
720 10.5061/dryad.44j0zpcfp).

721 Benefits generated: Benefits from this research accrue from the sharing of our data and results  
722 on public databases as described above.

## 723 **Author Contributions**


724 JM, AK, BdT, ALB and JFG designed the study. AK, JM, JCG, AC, BdT, ALB, MG, GP, RG,  
725 ID and PMF conducted the field work. AK and SM performed the laboratory work. AK and  
726 JC analyzed the data. AK wrote the initial version of the manuscript, which was edited by JM,  
727 JC, BdT and JFG, and all authors contributed to its improvement.

728

729

730

## Figures



731

732

733

734


735

736

737

738

**Figure 1: (A) Location of study sites (French Guiana). (B) Regression of the Shannon index of mammals on human-induced disturbance. Shannon indices of mammals were estimated through dipteran blood meals in each site. The level of human-induced disturbance was measured using the human footprint index (de Thoisy et al., 2010). The mean prediction curve is depicted with 90% CI. In the inset figure, the posterior density of the regression's slope coefficient is represented (dotted lines are positioned at 5% and 95% quantiles; the red line is positioned at  $x=0$ ).**



739

740

741

742

743

744

**Figure 2: Effect of mammal diversity changes on *Leishmania* transmission, assessed by a series of Bayesian regressions and posterior simulations. The most significant relationships are summarized in the top-panel diagram (but see Figure S7 for the complete model). Direct and indirect relationships are represented with solid and dashed arrows, respectively. Positive, negative, and non-significant effects are represented by red, blue and black arrows, respectively. Light grey arrows represent deterministic relationships. For each**

745 studied relationship, regression plots are presented and referenced with the corresponding letter in the  
746 lower panels. (A,B,C) Regression curves are depicted with 90% CI, and the posterior density of regression's  
747 slopes are represented in inset plots (dotted lines are positioned at 5% and 95% quantiles; the red line is  
748 positioned at  $x=0$ ). (D,E) The indirect (cumulative) effect of mammal diversity on *Leishmania* transmission  
749 was assessed by using posterior samples of model parameters to predict the *Leishmania* prevalence rate  
750 and the expected number of infected sand flies per trap. Sampled mean prediction curves are plotted (a  
751 subset of 200 curves, in order to facilitate visualization). Curves indicating an overall increase or decrease  
752 across the range of predictor values are depicted in red and blue, respectively. Inner plots represent the  
753 posterior density of the average slope of mean prediction curves.

754

**Table 1: Details about the study sites, including sampling size and estimates of the different variables used to describe the ecological system.**

Site	Nb. sand flies	Nb. blood-fed dipt. (success. ident.)	HFP	Mammal diversity (Shannon index) <sup>†</sup>	Prop. <i>Leish.</i> reservoirs <sup>†</sup>	Sand fly density (indiv. per trap) <sup>†</sup>	Sand fly diversity (Shannon index) <sup>†</sup>	<i>Leishmania</i> prev. rate (%) <sup>†</sup>
Belizon: A	1,028	17 (13)	36.0	2.22 [1.53,2.83]	0.42 [0.19,0.67]	44.35 [42.27,46.45]	2.18 [2.16,2.16]	1.05 [0.8,1.33]
Belizon: B	1,176	67 (38)	30.1	1.68 [1.16,2.18]	0.66 [0.38,0.89]	55.75 [53.31,58.25]	2.07 [2.04,2.04]	1.64 [1.37,1.92]
Belizon: C	1,298	22 (10)	17.0	1.95 [1.23,2.66]	0.71 [0.43,0.95]	58.69 [56.17,61.26]	1.72 [1.71,1.71]	1.71 [1.46,1.98]
Camp du tigre: Layon BdT	450	43 (30)	40.9	0.46 [0.13,0.95]	0.98 [0.91,1]	11.87 [10.97,12.79]	2.38 [2.35,2.35]	3.13 [2.18,4.22]
Counami: Carbet	349	60 (40)	21.0	2.27 [1.69,2.79]	0.65 [0.43,0.87]	15.18 [14.11,16.3]	1.45 [1.4,1.4]	1.4 [1.09,1.77]
Counami: Crossing	719	35 (32)	17.0	1.1 [0.49,1.85]	0.87 [0.58,1]	38.17 [36.06,40.33]	1.48 [1.46,1.46]	2.2 [1.7,2.78]
Counami: Patagai	572	13 (6)	17.0	1.57 [0.69,2.45]	0.49 [0.13,0.87]	19.43 [18.16,20.7]	1.69 [1.67,1.67]	1.08 [0.83,1.35]
Counami: T1	376	13 (8)	9.7	1.3 [0.36,2.2]	0.3 [0.04,0.68]	12.6 [11.58,13.65]	2.52 [2.47,2.47]	0.89 [0.54,1.31]
Counami: T2	817	97 (74)	4.0	2.09 [1.52,2.61]	0.69 [0.49,0.87]	30.7 [29.06,32.39]	1.9 [1.88,1.88]	1.64 [1.38,1.93]
Kaw: NR Tresor	2,964	50 (31)	8.0	2.32 [1.75,2.83]	0.61 [0.41,0.82]	109.27 [106.04,112.56]	1.68 [1.68,1.68]	1.46 [1.15,1.79]
NR Nouragues: Museum plot	2,676	196 (146)	4.0	2.41 [2.08,2.7]	0.18 [0.09,0.29]	142.16 [138.52,145.78]	1.93 [1.93,1.93]	0.68 [0.47,0.93]
NR Trinite: Layon D	795	14 (9)	0.0	2.27 [1.59,2.87]	0.48 [0.27,0.72]	26.56 [25.05,28.13]	1.56 [1.55,1.55]	1.04 [0.81,1.3]
Saint-Georges: Bridge	1,647	100 (70)	32.4	1.75 [1.4,2.11]	0.53 [0.31,0.76]	54.83 [52.06,57.66]	0.94 [0.93,0.93]	1.06 [0.76,1.4]
Saint-Georges: SG30B	182	38 (36)	22.7	1.03 [0.37,1.75]	0.89 [0.66,1]	15.25 [13.49,17.05]	2.39 [2.32,2.32]	2.67 [1.93,3.52]
Saint-Georges: SG6A	1,606	38 (27)	26.5	1.98 [1.42,2.52]	0.67 [0.48,0.84]	89.1 [85.42,92.83]	1.2 [1.19,1.19]	1.49 [1.13,1.9]
Saint-Georges: Track ONF	749	35 (21)	14.7	2.66 [2.19,2.99]	0.35 [0.19,0.52]	42.64 [40.22,45.05]	1.4 [1.39,1.39]	0.8 [0.59,1.03]

<sup>†</sup>Mean estimate (90% HPD)



**Table 2: List of vertebrates identified in blood meals of each dipteran species or MOTU**

Species/MOTU	Nb.	Vertebrates identified
Ceratopogonidae MOTU 1	3	<i>Rhinella</i> cf. <i>margaritifera</i> (67%); Hylloidea MOTU 1 (33%)
Ceratopogonidae MOTU 2	2	<i>Osteocephalus</i> MOTU 1 (50%); <i>R. cf. margaritifera</i> (50%)
Ceratopogonidae MOTU 3	1	<i>Choloepus didactylus</i> (100%)
Ceratopogonidae MOTU 4	1	<i>Osteocephalus</i> MOTU 1 (100%)
Ceratopogonidae MOTU 5	2	<i>Osteocephalus</i> MOTU 1 (100%)
Aedini MOTU 1	1	<i>Adenomera andreae</i> (100%)
<i>Culex imitator</i>	1	Squamata MOTU 1 (100%)
<i>Culex</i> MOTU 1	58	<i>Proechimys cuvieri</i> (36%); <i>Thamnophilus nigrocinereus</i> (28%); <i>Cuniculus paca</i> (7%); <i>Dasyprocta leporina</i> (3%); <i>Didelphis marsupialis</i> (3%); <i>Mazama nemorivaga</i> (3%); <i>Proechimys guyannensis</i> (3%); <i>Thamnophilus</i> MOTU 1 (3%); <i>Bradypus tridactylus</i> (2%); <i>Chelonoidis denticulatus</i> (2%); <i>Dasybus novemcinctus</i> (2%); <i>Echimyus chrysurus</i> (2%); <i>Metachirus nudicaudatus</i> (2%); <i>Oecomys rutilus</i> (2%); <i>Tapirus terrestris</i> (2%)
<i>Culex</i> MOTU 2	14	Squamata MOTU 1 (14%); <i>T. nigrocinereus</i> (14%); Accipitrinae MOTU 1 (7%); <i>Anolis fuscoauratus</i> (7%); <i>Bothrops atrox</i> (7%); <i>D. leporina</i> (7%); <i>D. novemcinctus</i> (7%); <i>Gonatodes humeralis</i> (7%); <i>Hypsiobas</i> MOTU 1 (7%); <i>Philander opossum</i> (7%); <i>Polychrus marmoratus</i> (7%); <i>Thamnophilus</i> MOTU 1 (7%)
<i>Culex</i> MOTU 3	2	<i>D. marsupialis</i> (50%); <i>T. nigrocinereus</i> (50%)
<i>Culex</i> sp.stJ	1	<i>Thamnophilus</i> MOTU 2 (100%)
<i>Culex</i> sp.stK	1	<i>D. leporina</i> (100%)
<i>Culex</i> sp.stL	1	<i>Osteocephalus</i> MOTU 1 (100%)
Culicinae MOTU 1	3	<i>P. cuvieri</i> (67%); <i>Thamnophilus</i> MOTU 1 (33%)
<i>Limatus flavisetosus</i>	1	<i>Tinamus major</i> (100%)
<i>Ochlerotatus serratus</i>	5	<i>D. leporina</i> (40%); <i>Thamnophilus</i> MOTU 1 (40%); <i>M. nemorivaga</i> (20%)
<i>Psorophora ferox</i>	1	<i>Myoprocta acouchy</i> (100%)
Sabethini MOTU 1	4	<i>D. leporina</i> (25%); <i>D. novemcinctus</i> (25%); <i>G. humeralis</i> (25%); <i>P. opossum</i> (25%)
<i>Bichromomyia flaviscutellata</i>	5	<i>D. leporina</i> (40%); <i>P. guyannensis</i> (40%); <i>P. cuvieri</i> (20%)
<i>Evandromyia brachyphalla</i>	1	<i>P. cuvieri</i> (100%)
<i>Ev. infraspinoso</i>	4	<i>D. novemcinctus</i> (50%); <i>C. didactylus</i> (25%); <i>M. acouchy</i> (25%)
<i>Ev. monstrosa</i>	1	<i>D. leporina</i> (100%)
<i>Ev. sericea</i>	5	<i>Lachesis muta</i> (60%); <i>M. acouchy</i> (40%)
<i>Ev. walkeri</i>	1	<i>Isothrix sinnamariensis</i> (100%)
<i>Pintomyia serrana</i>	5	<i>Alouatta seniculus macconnelli</i> (60%); <i>C. didactylus</i> (20%); <i>Pithecia pithecia</i> (20%)
<i>Micropygomyia roretaensis</i>	1	Squamata MOTU 1 (100%)
<i>Nyssomyia sylvicola</i>	2	<i>D. leporina</i> (100%)
<i>Ny. umbratilis</i>	13	<i>C. didactylus</i> (49%); <i>Coendou melanurus</i> (15%); <i>A. macconnelli</i> (14%); <i>Ateles paniscus</i> (9%); <i>D. novemcinctus</i> (4%); <i>Coendou prehensilis</i> (3%); <i>Tamandua tetradactyla</i> (2%); <i>Cebus olivaceus</i> (1%); <i>Nasua nasua</i> (1%); <i>P. pithecia</i> (1%); <i>Psophia crepitans</i> (1%)
<i>Ny. yuilli pajoti</i>	18	<i>A. macconnelli</i> (17%); <i>C. didactylus</i> (17%); <i>A. paniscus</i> (11%); <i>C. melanurus</i> (11%); <i>D. novemcinctus</i> (11%); <i>P. crepitans</i> (11%); <i>T. tetradactyla</i> (11%); <i>C. prehensilis</i> (6%); <i>T. terrestris</i> (6%)
<i>Pi. paca</i>	1	<i>M. nemorivaga</i> (100%)
<i>Pressatia choti</i>	7	<i>D. leporina</i> (86%); <i>G. humeralis</i> (14%)
<i>Psychodopygus amazonensis</i>	32	<i>D. novemcinctus</i> (44%); <i>D. leporina</i> (16%); <i>T. terrestris</i> (16%); <i>Dasybus kappleri</i> (12%); <i>D. marsupialis</i> (6%); <i>Mazama americana</i> (3%); <i>Tayassu pecari</i> (3%)
<i>Ps. ayrozai</i>	79	<i>D. novemcinctus</i> (84%); <i>D. kappleri</i> (15%); <i>Thamnophilus</i> MOTU 1 (1%)
<i>Ps. clausi</i>	11	<i>C. paca</i> (36%); <i>M. acouchy</i> (36%); <i>D. leporina</i> (18%); <i>M. nemorivaga</i> (9%)
<i>Ps. hirsutus</i>	34	<i>D. novemcinctus</i> (59%); <i>M. americana</i> (15%); <i>T. terrestris</i> (15%); <i>D. kappleri</i> (12%)
<i>Ps.</i> MOTU 1	8	<i>D. novemcinctus</i> (50%); <i>D. leporina</i> (12%); <i>D. kappleri</i> (12%); <i>M. acouchy</i> (12%); <i>T. terrestris</i> (12%)
<i>Ps.</i> MOTU 10	1	<i>D. novemcinctus</i> (100%)
<i>Ps.</i> MOTU 2	5	<i>D. novemcinctus</i> (40%); <i>T. terrestris</i> (40%); <i>Pecari tajacu</i> (20%)
<i>Ps.</i> MOTU 3	5	<i>D. novemcinctus</i> (60%); <i>D. kappleri</i> (40%)
<i>Ps.</i> MOTU 5	3	<i>D. novemcinctus</i> (100%)
<i>Ps. panamensis</i>	3	<i>D. leporina</i> (100%)
<i>Ps. s. maripaensis</i>	12	<i>D. novemcinctus</i> (67%); <i>M. americana</i> (11%); <i>D. kappleri</i> (10%); <i>T. terrestris</i> (5%); <i>M. nemorivaga</i> (3%); <i>C. paca</i> (2%); <i>A. paniscus</i> (1%); <i>P. guyannensis</i> (1%); <i>Thamnophilus</i> MOTU 1 (1%)
<i>Sciopemyia sordellii</i>	2	<i>Chiasmocleis shudikarensis</i> (50%); Hylloidea MOTU 2 (50%)
<i>Trichophoromyia ininii</i>	1	<i>C. paca</i> (100%)
<i>Trichopygomyia trichopyga</i>	6	<i>D. novemcinctus</i> (50%); <i>C. paca</i> (33%); <i>D. kappleri</i> (17%)
<i>Viannomyia tuberculata</i>	1	<i>C. melanurus</i> (100%)