



HAL
open science

Species choice and N fertilization influence yield gains through complementarity and selection effects in cereal-legume intercrops

Rémi Mahmoud, Pierre Casadebaig, Nadine Hilgert, Lionel Alletto, Grégoire T. Freschet, Claire de Mazancourt, Noémie Gaudio

► To cite this version:

Rémi Mahmoud, Pierre Casadebaig, Nadine Hilgert, Lionel Alletto, Grégoire T. Freschet, et al.. Species choice and N fertilization influence yield gains through complementarity and selection effects in cereal-legume intercrops. *Agronomy for Sustainable Development*, 2022, 42 (2), 10.1007/s13593-022-00754-y . hal-03582634

HAL Id: hal-03582634

<https://hal.inrae.fr/hal-03582634v1>

Submitted on 10 Mar 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

1 **Species choice and N fertilization influence yield gains through complementarity and**
2 **selection effects in cereal-legume intercrops**

3 Rémi Mahmoud^{1*}, Pierre Casadebaig¹, Nadine Hilgert², Lionel Alletto¹, Grégoire T. Freschet³, Claire
4 de Mazancourt³, Noémie Gaudio¹

5
6 ¹AGIR, University of Toulouse, INRAE, Castanet-Tolosan, France

7 ²MISTEA, University of Montpellier, INRAE Institut Agro, Montpellier, France

8 ³Station d'Ecologie Théorique et Expérimentale, CNRS, 2 route du CNRS, Moulis, France

9 *Corresponding author: remi.mahmoud@inrae.fr

10

11 Key words: cereal-legume intercropping, biodiversity effect, complementarity effect, selection effect

12 **Abstract**

13 Maintaining yield when reducing inputs is one prime objective of sustainable agriculture. In this
14 context, cereal-legume intercropping is a practice that can achieve increased yield under low-input
15 conditions through the complementary use of abiotic resources and facilitation mechanisms. Many
16 management options exist to design cereal-legume intercropping systems, among which the choice of
17 the species intercropped and the level of nitrogen (N) fertilization are essential.

18 In this study, we collected the results of 35 field experiments across Europe of cereal-grain legume
19 intercrops that combined various intercropped species and N fertilization levels. We first assessed the
20 intensity of the biodiversity effect and its components in unfertilized intercrops. Then, we focused on a
21 subset of systems to analyze how N fertilization influenced biodiversity effects on three intercrops
22 (durum wheat / pea, soft wheat / pea and durum wheat / faba bean). The biodiversity effect represents
23 the gap between observed and expected yield of a mixture. The complementarity effect is the
24 performance of mixtures relative to the performance of the component monocultures. The selection
25 effect captures the extent to which a species with a high monoculture yield dominate a mixture at the
26 expense of the other intercropped species.

27 Our results confirmed an overall positive biodiversity effect under unfertilized conditions and various
28 climate conditions (0.86 ± 0.04 t.ha⁻¹). Complementarity effect was the main driver as it represented
29 76% of the biodiversity effect, confirming intercropping as a useful practice in low-input systems. N
30 fertilization lowered the complementarity effect in durum wheat / pea intercrops, did not influence these
31 effects in soft wheat / pea intercrops and increased only the selection effect in durum wheat / faba bean
32 intercrops. These results highlight the need for a sufficiently competitive legume in intercrops when N
33 fertilizers are applied in order to avoid too much disruption of plant-plant interactions.

34 **Introduction**

35 From 1960-2000, the use of fertilizers, irrigation and pesticides mitigated effects of climatic hazards,
36 soil heterogeneity and pest pressure, and had a large and positive impact on crop yield (Tilman et al.
37 2002). More recently, especially in Europe, the growing trend of reducing inputs in agricultural systems,
38 due to environmental and social concerns, and the climatic uncertainty caused by climate change have
39 increased the variability in cropping conditions compared to that of the intensive agriculture practiced
40 in the late 20th century. To reduce the negative consequences of climatic uncertainty and continue to
41 produce enough food while reducing the use of inputs (Sadras and Denison 2016), a promising avenue
42 is to favor functional complementarity of abiotic resource use and biological regulations between plants
43 by designing innovative agricultural practices and systems (Duru et al. 2015). This can be achieved by
44 selecting relevant plant phenotypes (Lynch 2019) and/or using positive biodiversity effects through
45 plant mixtures, also known as the biodiversity-ecosystem function (BEF) effects (Brooker et al. 2021).

46 Positive BEF effects on ecosystem services have been widely studied in natural communities (Cardinale
47 et al. 2012), and interest in using them in cropping systems has increased in the past several years (Gurr
48 et al. 2016; Martin-Guay et al. 2018; Brooker et al. 2021). Analyzing the diversity-productivity
49 relationship enables the effect of biodiversity on primary production of a given system to be estimated
50 and can divide it into complementarity and selection effects (Loreau and Hector 2001). The former
51 measures the effect due to niche complementarity and/or facilitation, while the latter measures the effect
52 due to the dominance of a given species that fits well with the growth environment. Thus, BEF effects
53 should be viewed as resulting from particularly positive specific interactions rather than explaining
54 underlying processes themselves (Maier 2012). As Brooker et al. (2021) highlight, a collaboration gap
55 between BEF scientists and crop scientists has led to a poor understanding of “the operation of positive
56 diversity effects in intensive agricultural systems” and thus of how to enhance them.

57 In agricultural systems, plant diversity can be promoted by a range of intercropping practices (i.e.,
58 combining at least two crop species in the same field for most of their growing periods), which may
59 improve crop yield (Li et al. 2020a). Several mechanisms can, for example, improve nitrogen (N)
60 acquisition by the intercrops, including complementary distribution of roots in soil volumes (Postma
61 and Lynch 2012), use of distinct forms of N in soils (McKane et al. 2002) and fixation of atmospheric
62 N₂ by one species in the intercrop (Jensen et al. 2020). In a context of input reduction, the use of N₂-
63 fixing legumes is particularly promising. In Europe, this has been widely demonstrated in low-input
64 cereal-legume intercrops, with an increase in total yield and cereal grain quality compared to those of
65 sole crops (Bedoussac et al. 2015). However, supplying too much N fertilizer can cause the cereal to
66 dominate the legume, which decreases positive plant-plant interactions in intercropping systems (Pelzer
67 et al. 2012). Thus, the extent to which N fertilization can be used without compromising BEF effects in
68 such systems remains unclear. More particularly, while recent meta-analyses and reviews generally
69 agree upon positive BEF effects when multiple experiments are assessed, the results of individual

70 experiments have high variability (Bedoussac et al. 2015; Gurr et al. 2016; Raseduzzaman and Jensen
71 2017; Martin-Guay et al. 2018). Few recent studies underline a positive effect on intercrops' yield, via
72 temporal niche differentiation (Yu et al. 2016; Dong et al. 2018; Li et al. 2020b).

73 In this study, using a database of 35 field experiments (Fig. 1) from five European countries, we first
74 assessed the intensity of the biodiversity effect in winter and spring cereal-grain legume intercrops
75 under unfertilized conditions. Then, focusing on a subset of three winter intercrops – durum wheat
76 (*Triticum turgidum* L.) / pea (*Pisum sativum* L.), soft wheat (*Triticum aestivum* L.) / pea and durum
77 wheat / faba bean (*Vicia faba* L.) – we tested the influence of two levels of N fertilization (moderate
78 and high) on the biodiversity effect depending on the intercropped species considered.

79



80


81 Fig. 1. Example of a field experiment of winter wheat / pea intercrops (and their corresponding sole
82 crops) conducted at the ARVALIS experimental station, near Angers, France (Photograph courtesy of
83 C. Naudin, ESA, France).

84 **Materials and methods**

85 **1. Field experiments**

86 To estimate the net biodiversity effect on intercrop productivity in a wide range of environmental
87 conditions, we collected results from 35 factorial experiments conducted in five countries (France,
88 Denmark, Italy, Germany, and the United Kingdom; Fig. 2A), as detailed hereafter.

89 We used the following criteria to include set of experiments in our database: (1) grain yield was
90 measured for both species in sole- and intercropping conditions, (2) different species and genotypes
91 were used among cereal and legumes, and (3) a given mixture was observed at least in two locations.



92

93 Fig. 2. Location and main climatic features of the experiments. Panel A displays the number of
 94 experiments conducted at each location (different years and cropping systems). Panel B displays the
 95 sum of precipitation (mm) as a function of mean temperature (°C) during the crop cycle, with spring
 96 and winter crops encoded by colors, and experiment location encoded by symbols.

97 1.1. Environmental conditions

98 Climate conditions of each experiment were characterized using the following variables retrieved from
 99 the NASA POWER [API](#): the sum of precipitation (mm) and mean temperature (°C) during the crop
 100 cycle (from sowing to harvest dates). The experiments were separated into two groups: winter crops,
 101 which had higher precipitation (280-712 mm) and lower mean temperature (6.8-11.3°C) during the crop
 102 cycle, and spring crops, which had lower precipitation (60-366 mm) and higher mean temperature (12.3-
 103 17.3°C) (Fig. 2B).

104 1.2. Agricultural management

105 All experiments included cereal-grain legume intercrops of two annual crop species and their
 106 corresponding sole crops for which grain yield ($\text{t}\cdot\text{ha}^{-1}$) was measured at harvest. Cereals and legumes
 107 were each represented by three species: barley (*Hordeum vulgare* L.), durum wheat and soft wheat for
 108 the cereals and faba bean, lentil (*Lens culinaris* L.) and pea for the legumes (Table 1). In the database,
 109 39% and 61% of the intercrops were spring or winter crops, respectively. Intercropped species were
 110 sown and harvested at the same time. The sowing dates ranged from March 11 to May 3 for spring
 111 crops and from October 25 to December 15 for winter crops. The harvest dates for all crops ranged from
 112 June 6 to August 23.

113 Table 1. Description of the 35 cereal-legume experiments analyzed in this study. The *Type* column defines if the experiment is carried on conventional (C) or
 114 organic (O) farming.

Intercropped species (cereal / legume)	Country	Year(s)	Soil water capacity (mm)	Soil texture (clay-silt-sand, %)	Type	N treatments (kg,ha ⁻¹)	Mixture design	Spatial arrangement	No genotypes (cereal / legume)	Relative density in intercrop (cereal / legume)	References
Spring barley / fababean	Denmark	2001, 2002, 2003	173	24-29-47	O	0	substitutive	within row	2-1	0.5-0.5	(Gaudio et al., 2021; Hauggaard-Nielsen et al., 2008; Knudsen et al., 2004)
		2001, 2002, 2003	119	4-9-87	O	0	substitutive	within row	2-1	0.5-0.5	
Spring barley / pea	Denmark	2001, 2002, 2003	173	24-29-47	O	0	substitutive	within row	2-2	0.5-0.5	(Gaudio et al., 2021)
		2001, 2002, 2003	119	4-9-87	O	0	substitutive	within row	2-2	0.5-0.5	
		2003	173	24-29-47	O	0	substitutive, additive	alternate row	1-1	0.5-0.5, 0.5-1	
		2002	124	6-15-79	C	0	additive	alternate row	1-1	0.33-1	
		2003	124	6-15-79	C	0-130	substitutive, additive	alternate row	1-1	0.5-0.5, 0.5-1	
	France	2003, 2004	94	21-40-39	O	0	substitutive, additive	alternate row	1-1	0.5-0.5, 0.5-1	(Gaudio et al., 2021; Hauggaard-Nielsen et al., 2008, 2009; Launay et al., 2009)
	Germany	2004	176	51-29-20	O	0	substitutive, additive	alternate row	1-1	0.5-0.5, 0.5-1	
	Italy	2003, 2004	169	22-36-42	O	0	substitutive	alternate row	1-1	0.5-0.5	
United Kingdom	2003	142	49-32-19	O	0	substitutive, additive	alternate row	1-1	0.5-0.5, 0.5-1	(Gaudio et al., 2021; Hauggaard-Nielsen et al., 2008, 2009; Launay et al., 2009)	
Spring soft wheat / lentil	France	2015	135	10-8-82	O	0	substitutive, additive	within row	2-4	0.5-1, 0.33-1, 0.3-0.7, 0.17-1	
		2016	187	18-48-34	O	0	substitutive, additive	within row	2-4	0.5-1, 0.33-1.3, 0.33-1, 0.3-0.7, 0.17-1.3, 0.17-1	

Winter durum wheat / fababean	France	2010	187	18-48-34	C	0-60-80-140	substitutive, additive	alternate-, within row	1-1	0.5-0.5, 0.67-0.5, 0.67-1, 0.33-0.5	
		2011	187	18-48-34	C	0	substitutive	alternate row	1-1	0.5-0.5	
		2011	187	18-48-34	C	0-140	substitutive	alternate-, within row	1-1	0.5-0.5	
		2012	135	10-8-82	C	0	substitutive	within row	3-4	0.5-0.5	(Kammoun, 2014)
		2013	187	18-48-34	C	0	substitutive	within row	3-4	0.5-0.5	
Winter durum wheat / pea	France	2006	187	18-48-34	C	0-100-180	substitutive	alternate row	1-1	0.5-0.5	(Bedoussac and Justes, 2010a, 2010b)
		2007	135	10-8-82	C	0-60-80-140	substitutive	alternate row	4-1	0.5-0.5	
		2012	135	10-8-82	C	0	substitutive	within row	3-5	0.5-0.5	(Kammoun, 2014)
		2013	187	18-48-34	C	0-140	substitutive	within row	3-5	0.5-0.5	
		2015	135	10-8-82	C	0	substitutive, additive	within row	1-4	0.5-0.5, 0.5-1	
Winter soft wheat / fababean	France	2018	169	22-36-42	O	0	additive	within row	8-2	0.7-0.75	
Winter soft wheat / pea	France	2010	205	11-54-35	C	0-45-90-140	substitutive, additive	within row	1-1	0.5-0.5, 0.33- 0.66, 0.7-0.5	(Pelzer et al., 2016)
		2017	205	11-54-35	C	0	substitutive, additive	within row	1-2	0.5-0.5, 0.5-1, 0.15-1, 0.05-1	
		2007	83	20-38-42	C	0-30-45	substitutive	within row	1-1	0.5-0.5	(Gaudio et al., 2021; Naudin et al., 2010, 2014)
		2008	83	20-38-42	C	0-30-45-60-90	substitutive	within row	1-1	0.5-0.5	
		2017	197	19-49-32	O	0	additive	within row	8-3	0.5-0.75, 0.5-1	
		2018	169	22-36-42	O	0	additive	within row	8-3	0.5-0.75, 0.5-1	
		2006	94	21-40-39	O	0	substitutive	within row	1-1	0.5-0.5, 0.3-0.7	
		2007	94	21-40-39	O	0-30	substitutive	within row	1-1	0.5-0.5, 0.7-0.3	
		2008	94	21-40-39	O	0-35-72	substitutive	within row	1-1	0.5-0.5, 0.7-0.3	(Gaudio et al., 2021)
		2009	94	21-40-39	O	0-40	substitutive	within row	1-1	0.5-0.5, 0.7-0.3	

116 In the database, 54% of the intercrops were grown in a substitutive design (i.e., the sum of the relative
117 sowing densities of the two species intercropped equals 1), while 46% were grown in an additive design
118 (i.e., the sum of relative sowing densities exceeds 1). A species' relative density is its sowing density
119 in the intercrop relative to that in its reference sole crop. Consequently, the database contained 199 sole
120 crop experimental units and 307 intercrop experimental units (site x year x mix of genotypes x relative
121 densities x N treatment), of which 140 were in an additive design and 167 in a substitutive design.
122 Depending on the experiment, each experimental unit was replicated 2-8 times.

123 Additional details on experimental designs and management practices are reported in the reference
124 publications of 33 of the 35 experiments (Knudsen et al. 2004; Corre-Hellou et al. 2006; Hauggaard-
125 Nielsen et al. 2008, 2009; Launay et al. 2009; Bedoussac and Justes 2010a, b; Naudin et al. 2010, 2014;
126 Pelzer et al. 2016; Tang et al. 2016; Viguier et al. 2018; Gaudio et al. 2021).

127

128 **2. Estimating the biodiversity effect on intercrop performance**

129 For each experimental unit, grain yield ($t \cdot ha^{-1}$) was measured for each species. We calculated the
130 biodiversity effect (BE, Loreau and Hector 2001) as the observed grain yield minus expected grain yield
131 in intercrops (Eq. 1):

$$132 \quad BE = (YO_C + YO_L) - (YE_C + YE_L) \quad (\text{Eq. 1})$$

133 where YO_C and YO_L are the observed yield of the cereal and legume grown in intercrop, respectively,
134 and YE_C and YE_L are the expected yield of the cereal and legume grown in intercrop, respectively.

135 Expected yield was estimated from the yield of the species in sole crop weighted by its scaled relative
136 density in intercrop (Eq. 2; Li et al. 2020a):

$$137 \quad YE_C = M_C \frac{RD_C}{RD_C + RD_L} \quad \text{and} \quad YE_L = M_L \frac{RD_L}{RD_C + RD_L} \quad (\text{Eq. 2})$$

138 where M_C and M_L are the yield of the cereal and legume in sole crop, respectively, and RD_C and RD_L
139 are the relative density of the cereal and legume in intercrop, respectively. Grain yield in sole crops and
140 intercrops is calculated as the mean from each replicate of every experimental units, within each
141 experiment.

142 As mentioned, the biodiversity effect can be divided into a selection effect (SE, Eq. 3) and a
143 complementarity effect (CE, Eq. 4) (Loreau and Hector, 2001; Li et al. 2020a):

$$144 \quad SE = \frac{1}{2} \times \left(\left(\frac{YO_C}{M_C} - \frac{RD_C}{RD_C + RD_L} \right) - \left(\frac{YO_L}{M_L} - \frac{RD_L}{RD_C + RD_L} \right) \right) \times (M_C - M_L) \quad (\text{Eq. 3})$$

145
$$CE = \frac{M_C + M_L}{2} \times \left(\frac{YO_C}{M_C} - \frac{RD_C}{RD_C + RD_L} + \frac{YO_L}{M_L} - \frac{RD_L}{RD_C + RD_L} \right) = M \times (LER - 1) \text{ (Eq. 4)}$$

146 These formulas, used to compute selection and complementarity effects, are only valid in bispecific
147 mixtures.

148 The first term of Eq. 3 calculates the difference in increase or decrease in yield between the two species
149 intercropped, while the second term calculates the difference between their sole crop yields. Thus, a
150 positive selection effect means that the species with the higher yield in sole crop has a higher relative
151 increase in yield in intercrop (i.e., benefits more from intercropping).

152 Into the equation for the complementarity effect (Eq. 4), we introduced the classic Land Equivalent
153 Ratio, which is used to calculate land-use efficiency ($LER = Y_C/M_C + Y_L/M_L$; Willey and Rao 1980).

154 Thus, the complementarity effect equals the Land Equivalent Ratio minus 1, multiplied by M, the mean
155 yield in sole crops.

156

157 **3. Experimental design, data processing and analysis**

158 The data were curated and formatted in a database. The data were ordered, reshaped and homogenized
159 using the collection of R packages *tidyverse* (Wickham et al. 2019).

160 The dataset was unbalanced (i.e., groups had different numbers of observations) because the
161 experiments collected were conducted for different purposes and examined many factors (e.g., N
162 fertilization, intercrop design) (Table 1). Thus, the influence of several of the factors on the biodiversity
163 effect and its components could not be analyzed, especially due to the lack of certain treatments in some
164 experiments and to the nesting of factors. For example, only 12 of the 35 experiments tested N
165 fertilization levels, or the species effect also included site and year effects (e.g., spring barley / faba
166 bean intercrops were grown only in Denmark, so they could not be analyzed properly). The statistical
167 analysis performed was adjusted in response to this unbalanced structure.

168 We first investigated the overall behavior of mean biodiversity, complementarity and selection effects
169 within the unfertilized cereal-legume intercrops in the 35 experiments, and the correlation between the
170 biodiversity effect and each of its components. Thus, our goal was to assess the influence of N
171 fertilization on the biodiversity effect and its components. N fertilization ranged from 0-180 kg N.ha⁻¹,
172 which we split into three levels: null, moderate (30-80 kg N.ha⁻¹) and high (> 80 kg N.ha⁻¹). A factorial
173 design was then defined between the species intercropped and these levels of N fertilization. The subset
174 of our database with a factorial design of species and N fertilization levels corresponded to three
175 intercrops: durum wheat / pea, soft wheat / pea and durum wheat / faba bean (70 experimental units,
176 among which 62 are in substitutive design, all located in France, Table 1). Durum wheat / pea and

177 durum wheat / faba bean intercrops were grown in experiments with moderate and high levels of N
178 fertilization, while soft wheat / pea intercrops were grown only with a moderate level of N fertilization.

179 The effect of N fertilization on the biodiversity effect and its components in intercrops was assessed
180 using the Bayesian approach. Bayesian inference is based on reallocating credible values for a parameter
181 (posterior distribution) given prior knowledge (prior distribution) and the adequacy of the data to the
182 model (likelihood). The Bayesian approach provides information about the probability of a hypothesis
183 being true given the data ($P(\text{hypothesis}|\text{data})$). Bayesian estimation for the difference in group means
184 (Kruschke 2018) is an alternative to the classic Student's t test to compare the means of two groups.
185 This method calculates a posterior distribution for the mean differences between the two groups and
186 derives a 95% highest density interval (HDI), which is defined as the 95% most credible values of the
187 parameter. We performed Bayesian estimation for the difference in mean values of components of the
188 biodiversity effect between N-fertilized (moderate and high) and unfertilized treatments for each of the
189 three intercrops. The null hypothesis (H_0) was defined as equal mean biodiversity effect components
190 for N-fertilized and unfertilized intercrops. We applied the following decision rule to the position of the
191 95% HDI: reject H_0 if the 95% HDI excludes 0 but do not reject H_0 if it includes 0.

192 All indicator calculations and statistical analyses were performed with R software, v. 4.0.0 (R Core
193 Team 2020). Bayesian statistical analyses were performed using the R package *BEST* (Kruschke and
194 Meredith 2020).

195 **4. Definition of references for fertilized legumes**

196 A common assumption when calculating indicators to compare the performance of intercrops to that of
197 sole crops is that N is not a limiting resource for legumes and does not influence their yield (e.g., Pelzer
198 et al. 2012). To test this hypothesis, we performed Bayesian estimation for the difference in group means
199 between N-fertilized and unfertilized legume sole crops. The database contained only three experiments
200 (i.e., 11 experimental units) in which legume sole crops were N-fertilized, because the experiments we
201 collected were designed to conform to agronomic practices of farmers, who rarely fertilize legume sole
202 crops (Magrini et al. 2016). The Bayesian estimation confirmed that N fertilization had no significant
203 influence on the yield of legume sole crops. Given this result and the lack of data on N-fertilized legume
204 sole crops, we used the unfertilized legume sole crops as a reference when calculating the biodiversity
205 effect and its components in all experimental units.



206

207 **Results and discussion**

208 **1. Distribution of the biodiversity effect and its components in unfertilized intercrops**

209 On the whole dataset, the mean (± 1 standard error) yield gain in unfertilized intercrops equaled $0.86 \pm$
210 0.04 t.ha^{-1} ($1.04 \pm 0.01 \text{ t.ha}^{-1}$ for additive designs and $0.68 \pm 0.00 \text{ t.ha}^{-1}$ for substitutive designs) for a
211 mean total intercrop yield of $3.54 \pm 0.08 \text{ t.ha}^{-1}$ (Fig. 3A). These results highlight an increase in the yield
212 of cereal-legume intercrops in most experimental units under unfertilized conditions compared to those
213 of the corresponding sole crops, which agrees with results of several studies (Pelzer et al. 2012, 2014;
214 Yu et al. 2016) and confirms the ability of intercropping to increase grain yield in low-input farming
215 systems (Bedoussac et al. 2015).

216 However, the increase in yield observed was influenced by the cropping conditions used as references
217 to calculate the biodiversity effect. The unfertilized cereal sole crops used as references had lower grain
218 yield ($3.2 \pm 0.08 \text{ t.ha}^{-1}$, all cereals pooled) than cereals grown under conventional farming conditions,
219 which are always N fertilized (i.e., a mean grain yield of 6.1 t.ha^{-1} for the cereals of interest in the five
220 European countries considered for the period covered by the experiments (Food and Agriculture
221 Organization of the United Nations; <http://faostat.fao.org/>). Thus, the low yield observed for the
222 unfertilized cereal sole crops contributed greatly to the positive biodiversity effect estimated (Garnier
223 et al. 1997).



225 Fig. 3. **(A)** Distribution of unfertilized cereal-legume intercrop yield and biodiversity effect ($\text{t}\cdot\text{ha}^{-1}$).
226 Points represent the median, broad lines represent the interquartile range, and thin lines represent the
227 [0.1, 0.9] quantile interval. **(B)** Correlation between biodiversity effect ($\text{t}\cdot\text{ha}^{-1}$) and complementarity
228 effect ($\text{t}\cdot\text{ha}^{-1}$) or selection effect ($\text{t}\cdot\text{ha}^{-1}$) in unfertilized cereal-legume intercrops. Grey zones represent
229 the 95% confidence interval for the linear regressions. Data used: whole dataset ($n = 263$)

230

231 The biodiversity effect was strongly and positively correlated with the complementarity effect ($r = 0.86$,
232 $p < 10^{-15}$), but it was not correlated with the selection effect ($r = -0.01$, $p = 0.87$) (Fig. 3B). Thus, the
233 complementarity effect was the main driver of the yield gain in unfertilized cereal-legume intercrops,
234 meaning that positive plant-plant interactions (i.e., facilitation and / or niche complementarity) rather
235 than the dominance of one of the species increased intercrop yields (Pelzer et al. 2012). However,
236 caution is needed when distinguishing complementarity causes (e.g., niche partitioning, facilitation) of
237 the resulting complementarity effect (Barry et al. 2019). To quantify the relative importance of these
238 processes, specific measurements would be needed, such as symbiotic N_2 fixation to reflect differences
239 in N use between cereals and legumes, or a lodging score to quantify mechanical facilitation (e.g.,
240 Podgórska-Lesiak and Sobkowicz 2013). As Brooker et al. (2021) highlight, explicitly distinguishing
241 facilitation and niche partitioning would help when applying new analytical and conceptual frameworks
242 to design intercrops. Nevertheless, differences in N use in cereal-legume intercrops is a well-known
243 process in which the more competitive cereal usually takes disproportionately more soil mineral N than
244 the legume, which is forced to compensate by increasing symbiotic N_2 fixation (Rodriguez et al. 2020).
245 In a low-input context, this complementarity of N use enables cereals in intercrops to have higher grain
246 yield and quality than cereals in sole crops.

247 The complementarity effect contributed 76% of the biodiversity effect when the latter was positive (i.e.,
248 in 94% of the experimental units), but it contributed only 36% when the latter was negative (i.e., in 6%
249 of the experimental units). In the few cases in which we observed a yield loss in intercrops, the relative
250 contributions of complementarity and selection were reversed: -0.05 ± 0.02 and $-0.16 \pm 0.02 \text{ t}\cdot\text{ha}^{-1}$,
251 respectively. In these cases, the total yield of intercrops were lower than those of corresponding sole
252 crops because the competition between cereals and legumes exceeded the complementarity effect (also
253 reported by Pelzer et al. (2016) for soft wheat / pea intercrops and Baxevanos et al. (2017) for oat / pea
254 intercrops).


255

256

257 **2. Influence of N fertilization on the biodiversity effect and its components**

258 The biodiversity effect and its components were altered by N fertilization, which is a key practice in
259 agricultural systems. While the biodiversity effect was positive in 100% of the unfertilized experimental
260 units of the data subset considered (i.e., factorial designs of species and N fertilization levels), the

261 percentage of experimental units with a positive biodiversity effect decreased with N fertilization (i.e.,
 262 92% and 67% of the experimental units under moderately and highly N-fertilized conditions,
 263 respectively) (Fig. 4). Overall, the total intercrop yield increased with N fertilization (4.16 ± 0.18 , 5.09
 264 ± 0.24 and 4.62 ± 0.21 t.ha⁻¹ under unfertilized, moderately and highly N-fertilized conditions
 265 respectively); specifically, mean grain yield decreased for legumes (2.23 ± 0.12 , 1.88 ± 0.19 and 1.84
 266 ± 0.16 t.ha⁻¹ under unfertilized, moderately and highly N-fertilized conditions respectively) but
 267 increased for cereals (1.93 ± 0.20 , 3.21 ± 0.23 and 2.78 ± 0.15 t.ha⁻¹ under unfertilized, moderately and
 268 highly N-fertilized conditions respectively) with N fertilization (Fig. 4). The same pattern was observed
 269 for the complementarity effect, which was positive in 96%, 83% and 56% of the experimental units
 270 under unfertilized, moderately and highly N-fertilized conditions, respectively. Conversely, the
 271 percentage of experimental units with a positive selection effect increased with N fertilization: 25%,
 272 71% and 61% of the experimental units, under unfertilized, moderately and highly N-fertilized
 273 conditions, respectively. Thus, N fertilization tends to decrease positive plant-plant interactions within
 274 cereal-legume intercrops by acting on the balance between the two intercropped species to the benefit
 275 of the cereal (Pelzer et al. 2012).
 276



277
 278 Fig. 4. Distribution of cereal-legume intercrop yield, cereal and legume yield (t.ha⁻¹) and the biodiversity
 279 effect (t.ha⁻¹) as a function of nitrogen fertilization level. Points represent the median, broad lines
 280 represent the interquartile range, and thin lines represent the [0.1, 0.9] quantile interval. Data used:
 281 Experiments with a factorial design of species and N fertilization levels (n = 82)


282
 283 The effect of N fertilization on the biodiversity effect and its components depended on the species
 284 intercropped (Fig. 5). In durum wheat / pea intercrops, even moderate N fertilization decreased the
 285 biodiversity effect significantly by 66% compared to that under unfertilized conditions. This moderate

286 N fertilization increased the selection effect significantly by 0.21 t.ha⁻¹ (99.1% of the posterior values
287 for the difference in group means between N-fertilized and unfertilized conditions were positive), while
288 the complementarity effect decreased by 0.65 t.ha⁻¹ (99.1% of the posterior values for the difference in
289 means were negative). These effects were emphasized under highly N-fertilized conditions (Fig. 5).
290 When focusing on the yield of both species intercropped, N fertilization disadvantaged the legume,
291 since pea yield decreased by a mean of 37% under N-fertilized conditions compared to that under
292 unfertilized conditions, while the opposite was observed for durum wheat, whose yield increased by a
293 mean of 94%. These results could explain the shift in complementarity and selection effects for durum
294 wheat / pea intercrops between N-fertilized and unfertilized conditions. This behavior is usually
295 highlighted in existing literature related to cereal-legume intercrops (e.g., Naudin et al. 2010). Under
296 N-fertilized conditions, selection effect increases because durum wheat has a competitive advantage
297 over the legume (Mariotti et al. 2009; Duchene et al. 2017). Our results showed, however, that choosing
298 a different cereal or legume species can change this effect.

299 When soft wheat replaced durum wheat in wheat / pea intercrops, N fertilization did not influence the
300 biodiversity effect or its components (Fig. 5). Because the cereal and legume yields tended to increase
301 slightly with N fertilization, the latter did not disrupt the balance between the two species (Table 2).
302 Based on the soil and climate conditions considered, the level of N fertilization (45 kg N.ha⁻¹) was
303 probably too low, compared to usual N fertilization rates in conventional agriculture, to increase the
304 yield of one or both species significantly, unlike that of durum wheat / pea intercrops (60-140 kg N.ha⁻¹).
305

306 Finally, in durum wheat / faba bean intercrops, N fertilization did not influence the biodiversity effect
307 or its complementarity effect, but it did increase the selection effect significantly by 0.3 t.ha⁻¹ and 0.2
308 t.ha⁻¹ under moderately and highly N-fertilized conditions, respectively (95.5% and 95.2% of posterior
309 values for the difference in group means were positive, respectively) (Fig. 5). This increase was due to
310 an increase in durum wheat yield, since faba bean yield changed little in intercrops as N fertilization
311 increased. This behavior contrasts with that of pea yield when intercropped with durum wheat: pea yield
312 decreased as N fertilization increased. Height and biomass differences between two intercropped
313 species have been shown to influence their yields (Gaudio et al. 2021). Since the faba bean is taller and
314 larger than the pea (Guinet et al. 2018), it showed greater competitive ability (but whether aboveground
315 for light capture or belowground for nutrient and water acquisition remains to be tested), which explains
316 the lack of shift in the biodiversity effect observed in durum wheat / faba bean intercrops.

317



318

319 Fig. 5. Distribution of cereal and legume yields (t.ha⁻¹) in three cereal-grain legume intercrops (durum
 320 wheat / pea, soft wheat / pea and durum wheat / faba bean) as a function of nitrogen (N) fertilization
 321 level: null, moderate (30-80 kg N.ha⁻¹) and high (> 80 kg N.ha⁻¹). For the three intercrops, posterior
 322 distributions of the difference in mean of the biodiversity effect between the two N-fertilized (moderate
 323 and high) and unfertilized (N0) treatments is illustrated (t.ha⁻¹), with dashed lines representing the null
 324 value of the posterior difference in means. Points represent the median, broad lines represent the
 325 interquartile range, and thin lines represent the [0.1, 0.9] quantile interval. Data used: Experiments with
 326 a factorial design of species and N fertilization levels (n = 82).

327

328

329

330 3. Pathway to applications

331 Because cereal-legume intercrops are used mainly to decrease the use of agricultural inputs, most are
 332 managed without synthetic inputs. In this way, our study confirmed an increase in productivity under a
 333 wide range of unfertilized cropping conditions, with a balance between the two species intercropped
 334 (i.e., no species clearly dominated), although the increase depends on the species intercropped (Cheriere
 335 et al. 2020). N fertilization can disrupt this balance, shifting positive plant-plant interactions to a
 336 dominance of the cereal at the expense of the legume (e.g., in durum wheat / pea intercrops). This shift
 337 appeared at moderate N fertilization levels and even led to lower productivity of intercrops than that of
 338 sole crops at the high N fertilization levels applied to wheat sole crops in conventional agriculture (>
 339 100 kg N.ha⁻¹).

340 It would thus be interesting to identify the level of moderate N fertilization that provides benefits from
341 positive effects of intercropping and positive plant-plant interactions, while increasing the total yield
342 by increasing the cereal yield, as farmers often perform in winter intercrops (Verret et al. 2020). Because
343 this N level is likely to differ among species, future research should focus on the interaction between N
344 fertilization and the intercrop species chosen. For instance, recent meta-analysis (Li et al. 2020b) shows
345 high advantages of N fertilization on mixtures including maize (*Zea mays* L.).

346 In our study, only one combination of species x N fertilization had a positive interaction on yield (i.e.,
347 durum wheat / faba bean intercrops): cereal yield increased and legume yield remained the same, while
348 in durum wheat / pea intercrops, legume yield decreased. Thus, our results suggest that the legume
349 chosen can be a management mechanism, with the idea that the legume should be sufficiently
350 competitive to counterbalance the increased competition from the N-fertilized cereal (Duchene et al.
351 2017). Probably, it is the balance of competition between the two components rather than
352 competitiveness of the legume that matters. However, we also observed that the cereal yield stagnated
353 if the N fertilization level was not sufficient (e.g., soft wheat / pea intercrops). Thus, the optimal N
354 fertilization level should depend on the proportion of legume biomass in the intercrop (Naudin et al.
355 2010). As highlighted by other studies, the species chosen are a relevant mechanism for controlling
356 intercrops' yield (Cheriere et al. 2020) and suitability for the cropping environment in which they grow
357 (Baxevanos et al. 2017). Finally, it is worthwhile to recall that many barriers to adoption of intercrops
358 in Europe exist, beyond the scope of this article, such as. technical and economical ones (Bonke et al.
359 2020). Different possibilities (e.g., better communication of scientific results, breeding adapted to
360 intercrops) exist to overcome these barriers (Meynard et al. 2018) and allow intercrops to be more
361 widely cultivated.

362 **Conclusion**

363 This study highlights that the complementarity between intercropped species is the main driver of the
364 positive biodiversity effect on the performance of cereal-legume intercrops under diverse cropping
365 conditions. If the biodiversity effect depended instead mainly on the selection effect (i.e., if one
366 intercropped species strongly dominated), growing the dominant species alone would be more practical
367 agronomically, which would shift the balance towards sole crops.

368 While multiple meta-analyses and reviews highlighted the overall yield gain in intercrops, analysis and
369 tools to derive specific management recommendations for farmers from this general knowledge are still
370 lacking (Brooker et al. 2021). We argue that it may be counterproductive to emphasize that biodiversity
371 has this broad beneficial effect while the specific positive interactions between pairs of species and even
372 more so, cultivars, remain to be identified (Maier 2012).

373 The key question remains how to secure complementarity while intensifying or increasing productivity.
374 When focusing on the response of complementarity processes to N fertilization, we found that behavior

375 differed depending on the species chosen. We highlighted that N fertilization does not always depress
376 complementarity processes as long as the legume species can also benefit from it. Therefore, such shifts
377 in balance need to be understood through the prism of community ecology to develop the use of
378 intercrops in a wider range of agricultural systems besides low-input agriculture.

379 **Acknowledgements**

380 The authors thank all the technical staff who helped acquire this vast dataset and our colleagues who
381 provided us with the data: Laurent Bedoussac, Guénaëlle Corre-Hellou, Henrik Hauggaard-Nielsen,
382 Erik Steen Jensen, Etienne-Pascal Journet, Eric Justes, Nathalie Moutier, Christophe Naudin, Elise
383 Pelzer, and Loïc Viguier. We also thank Michael and Michelle Corson for their helpful comments and
384 English revision. This study was supported by the French National Research Agency under the
385 Investments for the Future Program (ANR-16-CONV-0004 and ANR-20-PCPA-0006) and by the
386 INRAE AgroEcoSystem Division.

387 **Declarations**

- 388 • Funding: This study was supported by the French National Research Agency under the Investments
389 for the Future Program (ANR-16-CONV-0004) and by the INRAE AgroEcoSystem Division
- 390 • Conflicts of interest/Competing interests (include appropriate disclosures): The authors declare that
391 they have no conflict of interest
- 392 • Ethics approval (include appropriate approvals or waivers): the study was performed in accordance
393 with the ethical standards as laid down in the 1964 Declaration of Helsinki and its later amendments
394 or comparable ethical standards
- 395 • Consent to participate (include appropriate statements): not appropriate
- 396 • Consent for publication (include appropriate statements): not appropriate
- 397 • Availability of data and material (data transparency): the datasets generated during and/or analyzed
398 during the current study are available from the corresponding author on reasonable request.
- 399 • Code availability (software application or custom code): not applicable
- 400 • Authors' contributions (include appropriate statements): “Funding acquisition: PC, NH, NG; data
401 collection and formatting: NG, RM, PC; data analysis: RM, NH; writing original draft: RM, NG,
402 PC, NH; writing, review and editing: all co-authors”

403

404 **References**

- 405 Barry KE, Mommer L, van Ruijven J, et al (2019) The Future of Complementarity: Disentangling
406 Causes from Consequences. *Trends Ecol Evol* 34:167–180. <https://doi.org/10.1016/j.tree.2018.10.013>
- 407 Baxevanos D, Tsialtas IT, Vlachostergios DN, et al (2017) Cultivar competitiveness in pea-oat
408 intercrops under Mediterranean conditions. *Field Crops Res* 214:94–103.
409 <https://doi.org/10.1016/j.fcr.2017.08.024>
- 410 Bedoussac L, Journet E-P, Hauggaard-Nielsen H, et al (2015) Ecological principles underlying the
411 increase of productivity achieved by cereal-grain legume intercrops in organic farming. A review.
412 *Agron Sustain Dev* 35:911–935. <https://doi.org/10.1007/s13593-014-0277-7>
- 413 Bedoussac L, Justes E (2010a) Dynamic analysis of competition and complementarity for light and N
414 use to understand the yield and the protein content of a durum wheat–winter pea intercrop. *Plant Soil*
415 330:37–54. <https://doi.org/10.1007/s11104-010-0303-8>
- 416 Bedoussac L, Justes E (2010b) The efficiency of a durum wheat-winter pea intercrop to improve yield
417 and wheat grain protein concentration depends on N availability during early growth. *Plant Soil* 330:19–
418 35. <https://doi.org/10.1007/s11104-009-0082-2>
- 419 Bonke, V., Musshoff, O. Understanding German farmer’s intention to adopt mixed cropping using the
420 theory of planned behavior. *Agron. Sustain. Dev.* 40, 48 (2020). [https://doi.org/10.1007/s13593-020-](https://doi.org/10.1007/s13593-020-00653-0)
421 [00653-0](https://doi.org/10.1007/s13593-020-00653-0)
- 422 Brooker RW, George TS, Homulle Z, et al (2021) Facilitation and biodiversity–ecosystem function
423 relationships in crop production systems and their role in sustainable farming. *J Ecol* n/a:
424 <https://doi.org/10.1111/1365-2745.13592>
- 425 Cardinale BJ, Duffy JE, Gonzalez A, et al (2012) Biodiversity loss and its impact on humanity. *Nature*
426 486:59–67. <https://doi.org/10.1038/nature11148>
- 427 Cherièrè T, Lorin M, Corre-Hellou G (2020) Species choice and spatial arrangement in soybean-based
428 intercropping: Levers that drive yield and weed control. *Field Crops Res* 256:107923.
429 <https://doi.org/10.1016/j.fcr.2020.107923>
- 430 Corre-Hellou G, Fustec J, Crozat Y (2006) Interspecific competition for soil N and its interaction with
431 N-2 fixation, leaf expansion and crop growth in pea-barley intercrops. *Plant Soil* 282:195–208.
432 <https://doi.org/10.1007/s11104-005-5777-4>

433 Dong, N., Tang, M. M., Zhang, W. P., Bao, X. G., Wang, Y., Christie, P., & Li, L. (2018). Temporal
434 Differentiation of Crop Growth as One of the Drivers of Intercropping Yield Advantage. *Scientific*
435 *Reports*, 8(1), 1–11. <https://doi.org/10.1038/s41598-018-21414-w>

436 Duchene O, Vian J-F, Celette F (2017) Intercropping with legume for agroecological cropping systems:
437 Complementarity and facilitation processes and the importance of soil microorganisms. A review. *Agric*
438 *Ecosyst Environ* 240:148–161. <https://doi.org/10.1016/j.agee.2017.02.019>

439 Duru M, Therond O, Fares M (2015) Designing agroecological transitions; A review. *Agron Sustain*
440 *Dev* 35:1237–1257. <https://doi.org/10.1007/s13593-015-0318-x>

441 Garnier E, Navas M-L, Austin MP, et al (1997) A problem for biodiversity-productivity studies: how
442 to compare the productivity of multispecific plant mixtures to that of monocultures? *Acta Oecologica*
443 18:657–670. [https://doi.org/10.1016/S1146-609X\(97\)80049-5](https://doi.org/10.1016/S1146-609X(97)80049-5)

444 Gaudio, N., Violle, C., Gendre, X., Fort, F., Mahmoud, R., Pelzer, E., Médiène, S., Hauggaard-Nielsen,
445 H., Bedoussac, L., Bonnet, C., Corre-Hellou, G., Couëdel, A., Hinsinger, P., Steen Jensen, E., Journet,
446 E.-P., Justes, E., Kammoun, B., Litrico, I., Moutier, N., ... Casadebaig, P. (2021). Interspecific
447 interactions regulate plant reproductive allometry in cereal–legume intercropping systems. *Journal of*
448 *Applied Ecology*, 00, 1– 11. <https://doi.org/10.1111/1365-2664.13979>

449 Guinet M, Nicolardot B, Revellin C, et al (2018) Comparative effect of inorganic N on plant growth
450 and N₂ fixation of ten legume crops: towards a better understanding of the differential response among
451 species. *Plant Soil* 432:207–227. <https://doi.org/10.1007/s11104-018-3788-1>

452 Gurr GM, Lu Z, Zheng X, et al (2016) Multi-country evidence that crop diversification promotes
453 ecological intensification of agriculture. *Nat Plants* 2:1–4. <https://doi.org/10.1038/nplants.2016.14>

454 Hauggaard-Nielsen H, Gooding M, Ambus P, et al (2009) Pea-barley intercropping for efficient
455 symbiotic N₂-fixation, soil N acquisition and use of other nutrients in European organic cropping
456 systems. *Field Crops Res* 113:64–71. <https://doi.org/10.1016/j.fcr.2009.04.009>

457 Hauggaard-Nielsen H, Jørnsgaard B, Kinane J, Jensen ES (2008) Grain legume–cereal intercropping:
458 The practical application of diversity, competition and facilitation in arable and organic cropping
459 systems. *Renew Agric Food Syst* 23:3–12. <https://doi.org/10.1017/S1742170507002025>

460 Hiederer, R. (2013), 'Mapping Soil Properties for Europe: Spatial Representation of Soil Database
461 Attributes', Technical report, JRC, Luxembourg: Publications Office of the European Union,
462 EUR26082EN Scientific and Technical Research series, ISSN 1831-9424, Citeseer.
463 <https://doi.org/10.2788/94128>

464 Jensen ES, Carlsson G, Hauggaard-Nielsen H (2020) Intercropping of grain legumes and cereals
465 improves the use of soil N resources and reduces the requirement for synthetic fertilizer N: A global-
466 scale analysis. *Agron Sustain Dev* 40:5. <https://doi.org/10.1007/s13593-020-0607-x>

467 Kammoun, B., 2014. Analyse des interactions génotype x environnement x conduite culturale de
468 peuplement bi-spécifique de cultures associées de blé dur et de légumineuses à graines, à des fins de
469 choix variétal et d'optimisation de leurs itinéraires techniques (PhD Thesis). Toulouse, INPT, Toulouse,
470 France.

471 Knudsen MT, Hauggaard-Nielsen H, Jørgensen B, Jensen ES (2004) Comparison of interspecific
472 competition and N use in pea-barley, faba bean-barley and lupin-barley intercrops grown at two
473 temperate locations. *J Agric Sci* 142:617–627. <https://doi.org/10.1017/S0021859604004745>

474 Kruschke JK (2018) Rejecting or Accepting Parameter Values in Bayesian Estimation. *Adv Methods*
475 *Pract Psychol Sci* 1:270–280. <https://doi.org/10.1177/2515245918771304>

476 Kruschke JK, Meredith M (2020) BEST: Bayesian Estimation Supersedes the t-Test. R package version
477 052 <https://CRANR-project.org/package=BEST>

478 Launay M, Brisson N, Satger S, et al (2009) Exploring options for managing strategies for pea-barley
479 intercropping using a modeling approach. *Eur J Agron* 31:85–98.
480 <https://doi.org/10.1016/j.eja.2009.04.002>

481 Li C, Hoffland E, Kuyper TW, et al (2020a) Yield gain, complementarity and competitive dominance
482 in intercropping in China: A meta-analysis of drivers of yield gain using additive partitioning. *Eur J*
483 *Agron* 113:125987. <https://doi.org/10.1016/j.eja.2019.125987>

484 Li, C., Hoffland, E., Kuyper, T. W., Yu, Y., Zhang, C., Li, H., Zhang, F., & van der Werf, W. (2020b).
485 Syndromes of production in intercropping impact yield gains. *Nature Plants*, 6(6), 653–660.
486 <https://doi.org/10.1038/s41477-020-0680-9>

487 Loreau M, Hector A (2001) Partitioning selection and complementarity in biodiversity experiments.
488 *Nature* 412:72–76. <https://doi.org/10.1038/35083573>

489 Lynch JP (2019) Root phenotypes for improved nutrient capture: an underexploited opportunity for
490 global agriculture. *New Phytol* 223:548–564. <https://doi.org/10.1111/nph.15738>

491 Magrini M-B, Anton M, Cholez C, et al (2016) Why are grain-legumes rarely present in cropping
492 systems despite their environmental and nutritional benefits? Analyzing lock-in in the French agrifood
493 system. *Ecol Econ* 126:152–162. <https://doi.org/10.1016/j.ecolecon.2016.03.024>

494 Maier DS. (2012) Theories of Biodiversity Value. In: What's So Good About Biodiversity?. The
495 International Library of Environmental, Agricultural and Food Ethics, vol 19. Springer, Dordrecht.
496 https://doi.org/10.1007/978-94-007-3991-8_6

497 Mariotti M, Masoni A, Ercoli L, Arduini I (2009) Above- and below-ground competition between
498 barley, wheat, lupin and vetch in a cereal and legume intercropping system. *Grass Forage Sci* 64:401–
499 412. <https://doi.org/10.1111/j.1365-2494.2009.00705.x>

500 Martin-Guay M-O, Paquette A, Dupras J, Rivest D (2018) The new Green Revolution: Sustainable
501 intensification of agriculture by intercropping. *Sci Tot Environ* 615:767–772.
502 <https://doi.org/10.1016/j.scitotenv.2017.10.024>

503 McKane RB, Johnson LC, Shaver GR, et al (2002) Resource-based niches provide a basis for plant
504 species diversity and dominance in arctic tundra. *Nature* 415:68–71. <https://doi.org/10.1038/415068a>

505 Meynard, JM., Charrier, F., Fares, M. et al. Socio-technical lock-in hinders crop diversification in
506 France. *Agron. Sustain. Dev.* 38, 54 (2018). <https://doi.org/10.1007/s13593-018-0535-1>

507 Naudin C, Corre-Hellou G, Pineau S, et al (2010) The effect of various dynamics of N availability on
508 winter pea-wheat intercrops: Crop growth, N partitioning and symbiotic N₂ fixation. *Field Crops Res*
509 119:2–11. <https://doi.org/10.1016/j.fcr.2010.06.002>

510 Naudin C, van der Werf HMG, Jeuffroy M-H, Corre-Hellou G (2014) Life cycle assessment applied to
511 pea-wheat intercrops: A new method for handling the impacts of co-products. *J Clean Prod* 73:80–87.
512 <https://doi.org/10.1016/j.jclepro.2013.12.029>

513 Pelzer E, Bazot M, Guichard L, Jeuffroy M-H (2016) Crop Management Affects the Performance of a
514 Winter Pea-Wheat Intercrop. *Agron J* 108:1089–1100. <https://doi.org/10.2134/agronj2015.0440>

515 Pelzer E, Bazot M, Makowski D, et al (2012) Pea–wheat intercrops in low-input conditions combine
516 high economic performances and low environmental impacts. *Eur J Agron* 40:39–53.
517 <https://doi.org/10.1016/j.eja.2012.01.010>

518 Pelzer E, Hombert N, Jeuffroy MH, Makowski D (2014). Meta-analysis of the effect of nitrogen
519 fertilization on annual cereal–legume intercrop production. *Agronomy Journal*, 106(5), 1775–1786.
520 <https://doi.org/10.2134/agronj13.0590>

521 Podgórska-Lesiak M, Sobkowicz P (2013) Prevention of pea lodging by intercropping barley with peas
522 at different nitrogen fertilization levels. *Field Crops Res* 149:95–104.
523 <https://doi.org/10.1016/j.fcr.2013.04.023>

524 Postma JA, Lynch JP (2012) Complementarity in root architecture for nutrient uptake in ancient
525 maize/bean and maize/bean/squash polycultures. *Ann Bot* 110:521–534.
526 <https://doi.org/10.1093/aob/mcs082>

527 R Core Team (2020) R: A Language and Environment for Statistical Computing

528 Raseduzzaman Md, Jensen ES (2017) Does intercropping enhance yield stability in arable crop
529 production? A meta-analysis. *Eur J Agron* 91:25–33. <https://doi.org/10.1016/j.eja.2017.09.009>

530 Rodriguez C, Carlsson G, Englund J-E, et al (2020) Grain legume-cereal intercropping enhances the
531 use of soil-derived and biologically fixed nitrogen in temperate agroecosystems. A meta-analysis. *Eur*
532 *J Agron* 118:126077. <https://doi.org/10.1016/j.eja.2020.126077>

533 Sadras VO, Denison RF (2016) Neither crop genetics nor crop management can be optimised. *Field*
534 *Crops Res* 189:75–83. <https://doi.org/10.1016/j.fcr.2016.01.015>

535 Tang X, Placella SA, Dayde F, et al (2016) Phosphorus availability and microbial community in the
536 rhizosphere of intercropped cereal and legume along a P-fertilizer gradient. *Plant Soil* 407:119–134.
537 <https://doi.org/10.1007/s11104-016-2949-3>

538 Tilman D, Cassman KG, Matson PA, et al (2002) Agricultural sustainability and intensive production
539 practices. *Nature* 418:671–677. <https://doi.org/10.1038/nature01014>

540 Verret V, Pelzer E, Bedoussac L, Jeuffroy M-H (2020) Tracking on-farm innovative practices to support
541 crop mixture design: The case of annual mixtures including a legume crop. *Eur J Agron* 115:126018.
542 <https://doi.org/10.1016/j.eja.2020.126018>

543 Viguier L, Bedoussac L, Journet E-P, Justes E (2018) Yield gap analysis extended to marketable grain
544 reveals the profitability of organic lentil-spring wheat intercrops. *Agron Sustain Dev* 38:39.
545 <https://doi.org/10.1007/s13593-018-0515-5>

546 Wickham H, François R, Henry L, Müller K (2019) *dplyr: A Grammar of Data Manipulation*

547 Willey RW, Rao MR (1980) A Competitive Ratio for Quantifying Competition Between Intercrops†.
548 *Exp Agric* 16:117–125. <https://doi.org/10.1017/S0014479700010802>

549

550 Yu, Y., Stomph, T. J., Makowski, D., Zhang, L., & van der Werf, W. (2016). A meta-analysis of relative
551 crop yields in cereal/legume mixtures suggests options for management. *Field Crops Research*, 198,
552 269–279. <https://doi.org/10.1016/j.fcr.2016.08.001>

553

554 Yu, Y., Stomph, T. J., Makowski, D., & van der Werf, W. (2015). Temporal niche differentiation
555 increases the land equivalent ratio of annual intercrops: A meta-analysis. *Field Crops Research*, 184,
556 133–144. <https://doi.org/10.1016/j.fcr.2015.09.010>