



# Experimental evidence for extra proton exchange in ribulose 1,5-bisphosphate carboxylase/oxygenase catalysis

Camille Bathellier, Guillaume Tcherkez

## ► To cite this version:

Camille Bathellier, Guillaume Tcherkez. Experimental evidence for extra proton exchange in ribulose 1,5-bisphosphate carboxylase/oxygenase catalysis. *Communicative and Integrative Biology*, 2022, 15 (1), pp.68-74. 10.1080/19420889.2022.2039431 . hal-03585340

**HAL Id: hal-03585340**

**<https://hal.inrae.fr/hal-03585340>**

Submitted on 23 Feb 2022

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License



## Experimental evidence for extra proton exchange in ribulose 1,5-bisphosphate carboxylase/oxygenase catalysis

Camille Bathellier & Guillaume Tcherkez

To cite this article: Camille Bathellier & Guillaume Tcherkez (2022) Experimental evidence for extra proton exchange in ribulose 1,5-bisphosphate carboxylase/oxygenase catalysis, Communicative & Integrative Biology, 15:1, 68-74, DOI: [10.1080/19420889.2022.2039431](https://doi.org/10.1080/19420889.2022.2039431)

To link to this article: <https://doi.org/10.1080/19420889.2022.2039431>



© 2022 The Author(s). Published by Informa UK Limited, trading as Taylor & Francis Group.



Published online: 15 Feb 2022.



[Submit your article to this journal](#)



Article views: 33



[View related articles](#)



[View Crossmark data](#)

SHORT COMMUNICATION



## Experimental evidence for extra proton exchange in ribulose 1,5-bisphosphate carboxylase/oxygenase catalysis

Camille Bathellier<sup>a</sup> and Guillaume Tcherkez<sup>b,c</sup>

<sup>a</sup>ISARA, Department of Agroecology & Environment, Agrapole, Lyon, France; <sup>b</sup>Research School of Biology, Australian National University, Canberra, Australia; <sup>c</sup>Institut de Recherche en Horticulture et Semences, Université d'Angers, INRAE, Beaucouzé, France

### ABSTRACT

Despite considerable advances in the past 50 y, the mechanism of ribulose 1,5-bisphosphate carboxylase/oxygenase (Rubisco) catalysis is still not well understood. In particular, the movement and exchange of protons within the active site is not well documented: typically, kinetics of H exchange during the first steps of catalysis, i.e. abstraction of the H3 atom of ribulose 1,5-bisphosphate (RuBP) and enolization, are not clearly established. Here, we took advantage of reaction assays run in heavy water (<sup>2</sup>H<sub>2</sub>O) to monitor the appearance of deuterated RuBP and deuterated products (3-phosphoglycerate and 2-phosphoglycolate) with exact mass LC-MS. Enolization was reversible such that de-enolization generated not only monodeuterated RuBP (<sup>2</sup>H-[H-3]-ribulose 1,5-bisphosphate) but also dideuterated RuBP (<sup>2</sup>H<sub>2</sub>-[H-3,O-3]-ribulose 1,5-bisphosphate). Carboxylation yielded about one half deuterated 3-phosphoglycerate (<sup>2</sup>H-[H-2]-3-phosphoglycerate) and also a small proportion of dideuterated 3-phosphoglycerate (<sup>2</sup>H<sub>2</sub>-[H-2,O-2]-3-phosphoglycerate). Oxygenation generated a small amount of monodeuterated, but no dideuterated, products. (Di)deuterated isotopologue abundance depended negatively on gas concentration. We conclude that in addition to the first step of proton exchange at H3 occurring before gas addition (and thus influenced by the competition between de-enolization and gas addition), there is another proton exchange step between solvent water, active site residues, and the 2,3-enediol(ate) leading to deuterated OH groups in products.

### ARTICLE HISTORY

Received 23 November 2021  
Revised 2 February 2022  
Accepted 3 February 2022

### KEYWORDS

Rubisco; catalysis; isotope; kinetics; carboxylation; oxygenation

## Introduction

Ribulose 1,5-bisphosphate carboxylase/oxygenase (Rubisco) is the fundamental enzyme of photosynthesis and catalyses carboxylation (CO<sub>2</sub> addition) or oxygenation (O<sub>2</sub> addition) of D-ribulose-1,5-bisphosphate (RuBP) and subsequent carbon–carbon cleavage to form two molecules of 3-phospho-D-glycerate (PGA) (with CO<sub>2</sub>) or one molecule of PGA plus one molecule of 2-phospho-glycolate (with O<sub>2</sub>) (Figure 1(a)). The reaction proceeds through several elemental steps including enolization, yielding a 2,3-enediolate which is the substrate of gas addition [1,2]. Despite the importance of this reaction, which abstracts about 120·10<sup>9</sup> tons of carbon from the atmosphere each year, events in catalysis remain unclear, in particular how protons are transferred between residues and exchanged with the solvent. Enolization, which is believed to be partially rate-limiting [3,4], involves both proton abstraction at H3 (proton attached to C3) and protonation at O2 (negatively charged oxygen atom attached to C2) (Figure 1(a,b)). Proton abstraction involves


carbamylation of Lys 201 (numbering in spinach; carbamylated Lys 201 is denoted thereafter as Lys 201<sup>c</sup>) [5,6]. The origin and the fate of the proton given to O2 is unclear. Cleland and coworkers [2] have suggested that the H3 proton abstracted by Lys 201<sup>c</sup> could shuttle to O2 and then to Lys 175 and may eventually undergo proton exchange with the solvent. However, the role of Lys 175 in enolization remains unclear since it could be either a proton acceptor or a proton donor and play a role at later steps in PGA formation [2,5,7–9]. Also, other residues (Thr 175, Lys 177 and His 294) can be involved in protonation-deprotonation events. In fact, the protein structure at 1.5 Å and 1.85 Å resolution with Mg<sup>2+</sup> and 2-carboxy-D-arabinitol 1,5-bisphosphate (carboxylation transition state analogue) as a ligand suggests that O2 is close to Lys 175 and Thr 173, while O3 is close to His 294 and Lys 201<sup>c</sup> and the carboxylate is at a short distance to Lys 177 [10,11].

The role of water molecules in catalysis also seems to be rather complicated [reviewed in [12]]. The structure of

**CONTACT** Guillaume Tcherkez ✉ [guillaume.tcherkez@anu.edu.au](mailto:guillaume.tcherkez@anu.edu.au) Research School of Biology, Australian National University, Canberra, ACT 2601, Australia

© 2022 The Author(s). Published by Informa UK Limited, trading as Taylor & Francis Group.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.



**Figure 1. Formal scheme showing proton exchanges in Rubisco catalysis.** (a) Chemical H exchanges during catalytic processes. The starting substrate is protiated ribulose 1,5-bisphosphate (denoted as *h*-RuBP). Enolization reversibility can generate deuterated RuBP, with a  $^2\text{H}$  atom at C3 (*d*-RuBP) or both C3 and O3 (*d*<sub>2</sub>-RuBP) via one or two solvent exchange steps. Solvent heavy water molecules involved in different steps of catalysis are labeled with different colors to allow tracing the fate of deuterons ( $^2\text{H}^+$ ). For clarity, only one residue is shown in this figure, Lys 201 (numbering in spinach), which is responsible for proton abstraction in enolization in its carbamylated form. Note that the C3 proton abstracted during enolization is eventually lost during catalysis and reprotonation uses a solvent water molecule, thus forming a 3-phosphoglycerate (PGA) molecule (referred to as "upper") deuterated at C2. In this panel, hydroxyl groups are simply represented as O–H for simplicity while in the chemical mechanism, H atoms can be shared with basic groups of active site residues, forming a partially charged oxygen atom. Note that the H ( $^2\text{H}$ ) atom at O2 in PGA can be slowly exchanged with solvent water after having been released by the enzyme (discussed in main text). (b) Summary of H exchange history of protons in RuBP.

the protein crystallized with  $\text{Ca}^{2+}$  shows that a water molecule participates in the  $\text{Mg}^{2+}$  coordination sphere [13] and could participate in the reaction. However, water is believed to be mostly excluded from the chemistry of  $\text{CO}_2$  addition and cleavage of the six-carbon intermediate. In fact, the binding of RuBP appears to sequester Lys 201<sup>c</sup> from the bulk and no water molecule is seen near the carbamate, implying the latter residue experiences a substantially non-aqueous environment [6,14]. Also, structural analysis has shown that both gaseous substrates  $\text{CO}_2$  and  $\text{O}_2$  are embedded in a positively charged cavity located at the C-terminal end of the  $\beta$ -barrel of the catalytic domain and closed with a negatively charged lid formed by the N-terminal domain of an adjacent L subunit [15]. Labeling experiments with tritiated RuBP ( $[\text{}^3\text{H-3}]\text{-RuBP}$ ) and spinach Rubisco showed negligible labeling in PGA [16] and, therefore, the H3 proton is believed to be lost before the production of

upper PGA. By contrast, enolization involves protons from water, as shown by pioneering assays of isotopic exchange between the H3 proton and solvent water: with bacterial Rubisco (*Rhodospirillum rubrum*) assayed in tritiated water ( $^3\text{H}_2\text{O}$ ) with protiated RuBP, RuBP molecules not consumed by the reaction are eventually fully  $^3\text{H}$ -labeled [3]. Similarly, plant Rubiscos assayed in heavy water ( $^2\text{H}_2\text{O}$ ) yield deuterated RuBP [17,18], allowing calculation of the reverse commitment to catalysis (de-enolization to gas addition ratio) of about 7 (carboxylation) and 500 (oxygenation)  $\mu\text{M}$  [17]. Interestingly, in  $^2\text{H}_2\text{O}$ , there is an enhancement of the production of side product pyruvate, which comes from  $\beta$ -elimination (competing with PGA production) of the six carbon intermediate [19]. That is, with a deuterium atom, reprotonation (redeuteration) of C2 is relatively slow, thereby favoring wasteful cleavage of the carbon-phosphate bond. This shows that the proton used for reprotonation may have


exchanged with the solvent at some point before or during gas addition, but underlying specific chemical events are unknown. A better knowledge of the chemical path followed by protons during Rubisco catalysis is therefore needed. Here, we took advantage of assays with Rubisco from tobacco (*Nicotiana tabacum*), natural (protiated) RuBP and heavy water (98%  $^2\text{H}_2\text{O}$ ) as a solvent, and followed isotopologues of substrate RuBP and products using high-resolution (exact mass) LC-MS.

## Results and discussion

### Liberation of deuterated RuBP via de-enolization

When the carboxylation assay was run in  $^2\text{H}_2\text{O}$  as a solvent with protiated RuBP as a substrate, there was a progressive monodeuteration in RuBP (+0.503138 a.m.u. for the  $m/z$  feature of interest). This deuteration reflected the formation of  $^2\text{H}$ -3-RuBP ( $d$ -RuBP) via de-enolization (Figure 2(a)). Under our conditions, about 40% of RuBP appeared to be monodeuterated while total RuBP declined since it was consumed by the reaction (Figure 2(a)). When dissolved  $\text{CO}_2$  concentration was

increased to 24  $\mu\text{M}$  (800 ppm in equilibrating gas phase) and RuBP initial concentration increased to keep the same proportion of consumed RuBP, there was a strong decline in the appearance of  $d$ -RuBP (Figure 2(b), blue), demonstrating that de-enolization directly competes with gas addition. Importantly, we also found the dideuterated RuBP isotopologue ( $d_2$ -RuBP, +1.006277 a.m.u.). Its content, which was always very small, increased with time during the assay and was undetectable with no reaction. This indicates that a second proton exchange occurs and involves the enzyme. It is likely that in dideuterated RuBP, one deuterium atom is at H3 and the second deuterium atom is carried by an hydroxyl (OH) group (see H inventory in RuBP summarized in Figure 1(b)). The two O–H bonds that are known to be weakened during the reaction are at O2 and O3 of the 2,3-enediol(ate) intermediate. However, only the H atom at O3 is retained in RuBP (since RuBP has a C=O group at C2). Secondary H exchange at other sites (such as C1) are less likely considering their  $\text{pK}_a$  values [20,21]. Therefore, it is plausible that  $d_2$ -RuBP corresponds actually to  $^2\text{H}_2$ -[H-3, O-3]-ribulose 1,5-bisphosphate. The  $^2\text{H}$  atom carried by O3 comes from water either directly or via an active site




**Figure 2. RuBP isotopologues observed during Rubisco-catalyzed carboxylation**, monitored using high resolution (exact mass) LC-MS. Substrate ribulose 1,5-bisphosphate (RuBP) was natural (protiated RuBP) and the solvent was heavy water ( $^2\text{H}_2\text{O}$ ). Here, signals associated with the major ion are shown, i.e.,  $[\text{M}-2\text{H}]^{2-}$  with a monoisotopic  $m/z$  value of 153.984921 a.m.u. Units of the x-axis are in half a.m.u. because of the charge ( $-2$ ) of the ion of interest. (a) Kinetics of protiated  $^{12}\text{C}$  (monoisotopic, denoted as  $m$ ) and protiated  $^{13}\text{C}$  RuBP disappearance and formation of deuterated RuBP ( $^2\text{H}$ ) during the reaction (assay of 60 seconds with  $\text{CO}_2$  equilibrated at 100 ppm in the gas phase). The inset shows the appearance of  $^{12}\text{C}$  dideuterated and  $^{13}\text{C}$  monodeuterated RuBP. (b) Isotopologue composition at 60s when  $\text{CO}_2$  is equilibrated at 100, 400 or 800 ppm in the gas phase. In all cases, the initial amount of RuBP was adjusted to reach  $\approx 40\%$  of initial concentration at the end of the assay. Note the much higher content in deuterated RuBP at low (100 ppm) and moderate (400 ppm)  $\text{CO}_2$  concentration, compared to high  $\text{CO}_2$  (800 ppm). As in (a), the inset shows  $^{12}\text{C}$  dideuterated and  $^{13}\text{C}$  monodeuterated isotopologues.

residue if there is a proton relay between water, Lys residue(s) and H at O3.

### ***Production of deuterated products***

Isotopic species of products was also monitored with exact mass LC-MS (Figure 3). As expected, there was an increase in both PGA and 2PG during the oxygenase assay. Interestingly, monodeuterated PGA and 2PG also accumulated (at +1.006277 a.m.u.) during the reaction. Since there is no reprotonation to form 'lower' PGA during the oxygenase reaction, this shows that deuterated

species came from isotopic exchange within the active site. Such isotopic exchange probably occurred at C1, O4 and/or C5 (numbers refer to numbering in RuBP). It is also possible that deuterated PGA came from residual carboxylase activity (despite constant equilibration with CO<sub>2</sub>-free N<sub>2</sub>/O<sub>2</sub> to keep oxygenation conditions) caused by remaining traces of bicarbonate used beforehand to pre-activate the enzyme (i.e. to carbamylate Lys 201). However, this cannot explain 2PG deuteration (Figure 3 (b), inset), which must come from secondary isotopic exchange at C1. As expected, the carboxylase reaction generated monodeuterated PGA, coming from



**Figure 3. Isotopologues of products of Rubisco catalysis:** oxygenation (a,b) and carboxylation (c,d), monitored using high-resolution (exact mass) LC-MS analysis. Substrate ribulose 1,5-bisphosphate (RuBP) was natural (protiated RuBP) and the solvent was heavy water ( $^2\text{H}_2\text{O}$ ). Major ions of 2-phosphoglycolate and 3-phosphoglycerate are (monoisotopic):  $[\text{M}-\text{H}]^-$  at 154.9745 a.m.u. and  $[\text{M}-\text{H}]^-$  at 184.9851 a.m.u., respectively. As in Figure 2, the figure shows the monoisotopic species (m),  $^{13}\text{C}$  and  $^2\text{H}$  isotopologues (a, b,c) and double isotopologues (inset of b, and d). Results presented here are associated with a 60s-assay (a,b,c) or the effect of  $\text{CO}_2$  concentration (d) equilibrated with 100, 400 or 800 ppm in the gas phase. In (d), the asterisk (\*) stands for other features ( $m/z$ ) present in the assay and unrelated to 2-phosphoglycolate or 3-phosphoglycerate. In (a) and (b), no dideuterated 3-phosphoglycerate or 2-phosphoglycolate could be observed.




reprotonation with solvent water during catalysis (green  $^2\text{H}$  atom in Figure 1). Unlike the oxygenase assay, there were dideuterated molecules essentially detectable at low  $\text{CO}_2$  (Figure 3(d)). It is likely that the second deuterium atom was carried by an OH group rather than the C atom from C1 in RuBP, since the observed amount is proportionally much larger than that of the monodeuterated species of 2PG.

### Implications for catalysis

Amounts of isotopic species of RuBP, PGA and 2PG formed during assays are compiled in Figure 4. Results are expressed in % of total content of the metabolite of interest. We recognize that whenever OH deuteration is involved, values shown here are probably slightly underestimated due to possible isotopic exchange (i.e.,  $^2\text{H}$  loss) during sample processing for LC-MS analysis. Nevertheless, such spontaneous exchange is slow [22] and furthermore, there was a clear difference between samples with and without reaction, showing the role played by the enzyme. Interestingly, there was slightly less ( $\approx 48\%$ ) than 50% of monodeuterated PGA. This is explained by (i) the contribution of dideuterated PGA

(less than 0.5%, Figure 4(a), inset) and (ii) the fact the assay was not run in 100%, but 98%  $^2\text{H}$ -water. In both oxygenase and carboxylase reactions, monodeuterated RuBP formation by de-enolization decreased with gas concentration. Monodeuterated PGA was not detected at high dissolved  $\text{O}_2$ , suggesting that isotopic exchange at the OH group at C4 occurred prior to gas addition and/or that high  $\text{O}_2$  was required to prevent the enzyme from consuming traces of dissolved  $\text{CO}_2$  in the reaction medium. Monodeuterated 2PG was observed at all  $\text{O}_2$  concentrations, suggesting spontaneous isotopic exchange upon 2PG release from the active site. The content in dideuterated RuBP depended negatively on gas concentration (both  $\text{CO}_2$  and  $\text{O}_2$ ) showing that deuteration took place within the active site prior to gas addition.

The data presented here provides compelling evidence that Rubisco catalysis is accompanied by proton exchanges with the solvent at molecular sites other than C2/3 (abstraction at C3 during RuBP enolization and reprotonation at C2 in PGA). To our knowledge, the present results provide the first evidence that there is a proton exchange other than at H3, after the generation of the 2,3-enediol(ate) and before gas addition. The site concerned by this



**Figure 4. Abundance of deuterated isotopologues in substrate and products of Rubisco catalysis.** Substrate ribulose 1,5-bisphosphate (RuBP) was natural (protiated RuBP) and the solvent was heavy water ( $^2\text{H}_2\text{O}$ ). RuBP and 3-phosphoglycerate (PGA) after 60s of carboxylase assay (a), RuBP, PGA and 2-phosphoglycerate (2PG) after 60s of oxygenase assay (b). The inset in (a) is a magnification to facilitate reading the percentage of dideuterated species. The value shown here is the percentage (%) with respect to the total pool size of the metabolite of interest; for example, 40%  $d_1$ -RuBP means that monodeuterated RuBP represents 40% of total RuBP (total = non-deuterated + monodeuterated + dideuterated + other isotopic forms such as the  $^{13}\text{C}$  isotopologue). Data shown are mean  $\pm$  SE ( $n = 3$ ). Gases concentrations are shown with respect to the gas phase used to equilibrate the reacting medium. Letters stand for statistical classes ( $P < 0.05$ ). ND, not detected. NR, no reaction.

exchange is very likely a OH group prone to deprotonation-reprotonation (i.e., acid-base exchange with amino acid residues of the active site). In other words,  $d_2$ -RuBP formed by de-enolization carries a deuteron at H3 and probably at O3. In fact, from consideration of crystal structures and mutation effects, the H atom bound to O3 is likely to be exchanged with His 294 [10,23]. It has been proposed that in the last steps of enolization, it can be replaced by the H3 atom previously abstracted by Lys 201<sup>c</sup> in the first step of enolization (proton relay hypothesis; it is shown in Figure 1 whereby the gray proton at O3 is lost and replaced by the blue deuteron from Lys 201<sup>c</sup>) [2]. The proton relay hypothesis is also supported by the fact that the use of monodeuterated RuBP as a substrate ( $^2\text{H}$ -3-ribulose 1,5-bisphosphate) affects the  $^{12}\text{C}/^{13}\text{C}$  kinetic isotope effect during carboxylation [17]. It means that bonding in the reaction intermediate and/or the transition state involves somehow the abstracted H3 proton before it is washed out to the solvent. The most likely is that the H3 proton interacts with O2 and/or O3 in the 2,3-enediol(ate) because it is sufficiently close to the C atom attacked by  $\text{CO}_2$  in carboxylation and therefore, it can affect significantly the  $^{12}\text{C}/^{13}\text{C}$  kinetic isotope effect.

The involvement of several proton exchanges during enolization and before gas addition is thus of importance for catalysis since it has potential effects on gas addition itself. That is, there is an interdependence of reaction steps during catalysis. It further suggests that in terms of enzyme evolution, the adaptation of the active site to adjust kinetic properties when  $\text{CO}_2$  and  $\text{O}_2$  availability varies, should not only involve residues directly involved in  $\text{CO}_2$  attack but also H bonding during enolization. This conclusion is in line with the recognized catalytic trade-off during Rubisco catalysis suggested by the comparison of kinetic properties in different Rubisco forms [24].

## Methods

### Rubisco purification and assays

Tobacco (*Nicotiana tabacum* var. Wisconsin) Rubisco was purified after [25] by precipitation from tobacco leaf extracts using 12% PEG, and then crystallization in Tris (25 mM) at pH 7.2. Rubisco crystals were collected, washed and redissolved (in Tris 5 mM pH 7.6, 20% glycerol) for storage at  $-80^\circ\text{C}$ . Rubisco assays were

conducted as in [17]. Assays were carried out at  $25^\circ\text{C}$  in septum-capped 2-mL vials filled with 300  $\mu\text{L}$  buffer (HEPES 100 mM, 20 mM  $\text{MgCl}_2$ , pH 8). The reaction medium was first equilibrated for 1 h with a gas mixture  $\text{CO}_2/\text{O}_2$  in  $\text{N}_2$  produced with high precision mass-flow controllers (FC260, Tylan).  $\text{CO}_2$  and  $\text{O}_2$  mole fractions were continuously monitored with an IRGA (Li-6251, Li-Cor) and an oxygen sensor (MAX-250, Mextec). 10  $\mu\text{L}$  of RuBP solution was then injected to get a final concentration of about 3 mM (adjusted to have about 40% consumption after reaction). The reaction was started with 10  $\mu\text{L}$  of Rubisco extract that had been pre-activated for 20 min with 20 mM  $\text{NaHCO}_3$  and 15 mM  $\text{MgCl}_2$ . The reaction ran under constant gas bubbling and was quenched with 240  $\mu\text{L}$  EDTA at 100 mM followed by snap-freezing in liquid nitrogen and lyophilization (freeze-dried).

### LC-MS analyses

Isotopologue composition of RuBP and products was assessed using high-resolution LC-MS Orbitrap Q Exactive (Thermo Scientific). Lyophilized samples from assays were redissolved in water, ions were removed with washed Dowex 50 H+ and the eluate was instant-frozen, lyophilized, and redissolved in 100  $\mu\text{L}$  water. Samples were injected directly (i.e. by infusion). MS analysis was operated in negative polarity in the full MS scan mode (mass scan range 50–750  $m/z$ ) with the following source settings: source voltage 3,500 V, resolution 70,000, AGC target  $1 \cdot 10^6$ , mass scan range 60–600  $m/z$ , sheath gas 40, auxiliary gas 10, sweep gas 1.5, probe temperature  $300^\circ\text{C}$ , capillary temperature  $250^\circ\text{C}$  and S-lens RF level 50.

### Acknowledgments

The authors thank the *Région Pays de la Loire* and *Angers Loire Métropole* for the financial support via the grant Connect Talent Isoseed. They also thank the Joint Mass Spectrometry Facility for giving access to high-resolution LC-MS.

### Disclosure statement

No potential conflict of interest was reported by the author(s).

### Funding

This work was supported by the Conseil Régional des Pays de la Loire [Connect Talent Isoseed].



## References

- [1] Tommasi IC. The mechanism of Rubisco catalyzed carboxylation reaction: chemical aspects involving acid-base chemistry and functioning of the molecular machine. *Catalysts*. 2021;11:Article no. 813.
- [2] Cleland WW, Andrews TJ, Gutteridge S, et al. Mechanism of Rubisco: the carbamate as general base. *Chem Rev*. 1998;98:549–562.
- [3] Saver BG, Knowles JR. Ribulose 1,5-bisphosphate carboxylase: enzyme-catalyzed appearance of solvent tritium at carbon 3 of ribulose 1,5-bisphosphate reisolated after partial reaction. *Biochemistry*. 1982;21:5398–5403.
- [4] Van Dyk DE, Schloss JV. Deuterium isotope effects in the carboxylase reaction of ribulose 1,5-bisphosphate carboxylase/oxygenase. *Biochemistry*. 1986;25:5145–5156.
- [5] King WA, Gready JE, Andrews TJ. Quantum chemical analysis of the enolization of ribulose biphosphate: the first hurdle in the fixation of CO<sub>2</sub> by Rubisco. *Biochemistry*. 1998;37:15414–15422.
- [6] Taylor TC, Andersson I. The structure of the complex between rubisco and its natural substrate ribulose 1,5-bisphosphate. *J Mol Biol*. 1997;265:432–444.
- [7] Mauser H, King WA, Gready JE, et al. CO<sub>2</sub> fixation by rubisco: computational dissection of the key steps of carboxylation, hydration, and C-C bond cleavage. *J Am Chem Soc*. 2001;123:10821–10829.
- [8] Knight S, Andersson I, Brändén C-I. Crystallographic analysis of ribulose 1,5-bisphosphate carboxylase from spinach at 2.4 Å resolution: subunit interactions and active site. *J Mol Biol*. 1990;215:113–160.
- [9] Harpel MR, Hartman FC. Facilitation of the terminal proton transfer reaction of ribulose 1,5-bisphosphate carboxylase/oxygenase by active-site Lys 166. *Biochemistry*. 1996;35:13865–13870.
- [10] Matsumura H, Mizohata E, Ishida H, et al. Crystal structure of rice Rubisco and implications for activation induced by positive effectors NADPH and 6-phosphogluconate. *J Mol Biol*. 2012;422:75–86.
- [11] Valegård K, Hasse D, Andersson I, et al. Structure of Rubisco from *Arabidopsis thaliana* in complex with 2-carboxyarabinitol-1,5-bisphosphate. *Acta Crystallogr D Struct Biol*. 2018;74:1–9.
- [12] Tcherkez GG, Bathellier C, Stuart-Williams H, et al. D<sub>2</sub> O solvent isotope effects suggest uniform energy barriers in ribulose 1, 5-bisphosphate carboxylase/oxygenase catalysis. *Biochemistry*. 2013;52:869–877.
- [13] Taylor TC, Andersson I. Structural transitions during activation and ligand binding in hexadecameric Rubisco inferred from the crystal structure of the activated unliganded spinach enzyme. *Nat Struct Biol*. 1996;3:95–101.
- [14] Andersson I. Large structures at high resolution: the 1.6 Å crystal structure of spinach ribulose-1,5-bisphosphate carboxylase/oxygenase complexed with 2-carboxyarabinitol bisphosphate. *J Mol Biol*. 1996;259:160–174.
- [15] Stec B. Structural mechanism of Rubisco activation by carbamylation of the active site lysine. *Proc National Acad Sci USA*. 2012;109:18785–18790.
- [16] Fiedler F, Müllhofer G, Trebst A, et al. Mechanism of ribulose-diphosphate carboxydismutase reaction. *Eur J Biochem*. 1967;1:395–399.
- [17] Bathellier C, Yu L-J, Farquhar GD, et al. Ribulose 1,5-bisphosphate carboxylase/oxygenase activates O<sub>2</sub> by electron transfer. *Proc National Acad Sci USA*. 2020;117:24234–24242.
- [18] Sue JM, Knowles JR. Ribulose 1,5-bisphosphate carboxylase: primary deuterium kinetic isotope effect using [3-<sup>2</sup>H] ribulose 1,5-bisphosphate. *Biochemistry*. 1982;21:5410–5414.
- [19] Andrews TJ, Kane HJ. Pyruvate is a by-product of catalysis by ribulose biphosphate carboxylase/oxygenase. *J Biol Chem*. 1991;266:9447–9452.
- [20] Isbell HS, Frush HL, Wade CWR, et al. Transformations of sugars in alkaline solutions. *Carbohydr Res*. 1969;9:163–175.
- [21] Richard JP. Acid-base catalysis of the elimination and isomerization reactions of triose phosphates. *J Am Chem Soc*. 1984;106:4926–4936.
- [22] Wu J, Serianni AS, Vuorinen T. Furanose ring anomericization: kinetic and thermodynamic studies of the D-2-pentuloses by <sup>13</sup>C-NMR spectroscopy. *Carbohydr Res*. 1990;206:1–12.
- [23] Harpel MR, Larimer FW, Hartman FC. Multiple catalytic roles of His 287 of *Rhodospirillum rubrum* ribulose 1,5-bisphosphate carboxylase/oxygenase. *Protein Sci*. 1998;7:730–738.
- [24] Tcherkez G, Farquhar GD. Rubisco catalytic adaptation is mostly driven by photosynthetic conditions—not by phylogenetic constraints. *J Plant Physiol*. 2021;Article no. 153554. DOI:10.1016/j.jplph.2021.153554
- [25] Servaites JC. Crystalline ribulose bisphosphate carboxylase/oxygenase of high integrity and catalytic activity from *Nicotiana tabacum*. *Arch Biochem Biophys*. 1985;238:154–160.