



HAL
open science

Reducing a model of sugar metabolism in peach fruit to explore genetic diversity

Hussein Kanso, Bénédicte Quilot-Turion, Mohamed-Mahmoud Memah, Olivier Bernard, Jean-Luc Gouzé, Valentina Baldazzi

► **To cite this version:**

Hussein Kanso, Bénédicte Quilot-Turion, Mohamed-Mahmoud Memah, Olivier Bernard, Jean-Luc Gouzé, et al.. Reducing a model of sugar metabolism in peach fruit to explore genetic diversity. iCROP2020: Crop modelling for the Future, Feb 2020, Montpellier, France. pp.1-5. <hal-03587179>

HAL Id: hal-03587179

<https://hal.inrae.fr/hal-03587179v1>

Submitted on 24 Feb 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



HAL Authorization

Reducing a model of sugar metabolism in peach fruit to explore genetic diversity

Hussein Kanso^{1,2}, Bénédicte Quilot-Turion¹,
Mohamed-Mahmoud Memah², Olivier Bernard⁴, Jean-Luc
Gouzé⁴, and Valentina Baldazzi^{2,3,4}

¹INRA, UR1052, GAFL France

²INRA, UR1115, PSH France

³Université Côte d'Azur , INRA, CNRS, ISA, France

⁴Université Côte d'Azur , Inria, INRA, Sorbonne Université,
BIOCORE, France

September 11, 2019

1 Introduction

Nowadays, increasing efforts are made to select varieties that respond to a large panel of criteria, including abiotic and biotic stress tolerance, increased yield and quality of food products. Gene-to-phenotype models have been considered as the tools of the future since they can help to test the performance of new genotypes under different environment and management conditions. A kinetic model of sugar metabolism has been developed by Desnoues et al. (2018) to simulate the accumulation of different sugars during peach fruit development as a set of parametric ordinary differential equations. The model correctly accounts for annual variability and for the genotypic variations observed in ten peach genotypes issued from an inter-specific population. Two major drawbacks of this model are the number of parameters to estimate and its integration time that can be costly due to non-linearities and time-dependent input functions. Together, these issues hamper the use of the model for the whole genetic progeny of 106 genotypes, for which few data

are available (six data points or less per sugar). Several reduction and approximation approaches exist in literature, each one addressing a specific aspect of model complexity (Cariboni et al., 2007; Heinrich and Schuster, 1996). In this work, we present a model reduction scheme that combines different methods in several parallel steps. The purpose is i) to obtain a simplified model showing comparable predictions as the original model while reducing its calibration time and number of parameters. ii) to calibrate it for the genotypic progeny and develop an integrated genetic-kinetic model.

2 Methods

First, multivariate sensitivity analysis (Lamboni et al., 2009) was applied to identify those parameters having a significant influence on the outputs of the model, over the whole dynamics. Second, we operated three structural simplifications in terms of network and reactions rates to reduce the complexity of the model. Third, timescale-based approaches and quasi-steady-state approximation (López Zazueta et al., 2018) were applied to reduce the number of ODEs of the model. The quality of individual and combined reduction steps was systematically evaluated with respect to the original model according to three criteria of major importance for our application: the Akaike Information Criterion (AIC), the calibration time and the expected error over a population of virtual genotypes.

3 Results and conclusions

Results from the reduction steps were combined into a final reduced model (Fig. 1). This model has only 9 parameters to be estimated, linear flows, 9 ODEs and only one temporal enzymatic capacity, common to all genotypes. Comparison between the reduced and the original model showed an equivalent fit quality and confirmed a strong benefice for most genotypes. Results showed a satisfactory agreement between predictions and experimental data. Thanks to the reduced calibration time, the whole progeny of 106 genotypes ‘could be’ calibrated. The parameter values obtained were considered as quantitative genetic traits and used for a Quantitative Trait Loci analysis. Results from this analysis were then used to define a genetic model to be incorporated in the reduced kinetic model. This opens the possibility of predicting sugar metabolism of genotypes carrying diverse combinations of alleles at the markers controlling the parameters.




Figure 1: Graphical representation of the proposed strategy to calibrate 106 genotypes. Orange rectangle represent reduction methods. Yellow rectangle represent model evaluation steps by means of our 3 criteria: AIC, calibration time and expected error. The specificity of the original and reduced models are indicated in blue. Green dashed line represent the perspective part.

References

- Cariboni, J., Gatelli, D., Liska, R., and Saltelli, A. (2007). The role of sensitivity analysis in ecological modelling. *Ecological Modelling*, 203(1):167 – 182.
- Desnoues, E., Génard, M., Quilot-Turion, B., and Baldazzi, V. (2018). A kinetic model of sugar metabolism in peach fruit reveals a functional hypothesis of a markedly low fructose-to-glucose ratio phenotype. *The Plant Journal*, 94(4):685–698.
- Heinrich, R. and Schuster, S. (1996). *The regulation of cellular systems*. Chapman and Hall.
- Lamboni, M., Makowski, D., Lehuger, S., Gabrielle, B., and Monod, H. (2009). Multivariate global sensitivity analysis for dynamic crop models. *Field Crops Research*, 113(3):312–320.

López Zazueta, C., Bernard, O., and Gouzé, J.-L. (2018). Analytical reduction of nonlinear metabolic networks accounting for dynamics in enzymatic reactions. *Complexity*, 2018.