



**HAL**  
open science

**Is seed yield in seed orchards affected differently in presence of one insect species or a subset of species. Preliminary results from 2015-2016 in France and Sweden**

Olle Rosenberg, Weslien Jan, Alain Roques, Marie-Anne Auger-Rozenberg

► **To cite this version:**

Olle Rosenberg, Weslien Jan, Alain Roques, Marie-Anne Auger-Rozenberg. Is seed yield in seed orchards affected differently in presence of one insect species or a subset of species. Preliminary results from 2015-2016 in France and Sweden. IUFRO Seed Orchards conference - The Key to Sustainable Bio-Economy, SLU, Sep 2017, Bålsta, Sweden. hal-03592963

**HAL Id: hal-03592963**

**<https://hal.inrae.fr/hal-03592963>**

Submitted on 31 Mar 2022

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



*WP 2.09.01 Seed Orchards*

PROCEEDINGS

# IUFRO

## Seed Orchard Conference 2017

SEPTEMBER 4-6, 2017 BÅLSTA, SWEDEN



skogforsk



# IS SEED YIELD IN SEED ORCHARDS AFFECTED DIFFERENTLY IN THE PRESENCE OF ONE INSECT SPECIES OR A SUBSET OF SPECIES? PRELIMINARY RESULTS FROM FRANCE AND SWEDEN 2015-2016

**Olle Rosenberg**, Skogforsk, the Forestry Research Institute of Sweden, Uppsala, Sweden  
E-mail: [olle.rosenberg@skogforsk.se](mailto:olle.rosenberg@skogforsk.se)

Jan Weslien, Skogforsk, the Forestry Research Institute of Sweden, Uppsala, Sweden  
Alain Roques and Marie-Anne Auger-Rozenberg, INRA, Orleans, France

Seed orchards of coniferous tree species have been established in many countries to produce high quality seeds for forestation. However, the seed production in seed orchards is far below the demand, and one reason for the low and unpredictable seed production is damage caused by insects. Currently, only one insecticide is registered for use in seed orchards in both Sweden and France, and this is not active against all occurring pest species. The project in focus here aims to explore if it is a risk that pesticide targeted against certain pest species can affect other pest species through relieved intraspecific competition. This is an important issue in pest control in general when using more specific insecticides or pheromones (as opposed to broad-spectrum insecticides), when several species frequently inhabit the same resource. The risk is that species not or less affected by the insecticide can increase its feeding, resulting in a low effect of the treatment on seed yield and seed quality. To study an effect on yield and quality of seeds, strobili (female spruce flowers that will develop to cones) were protected by fine mesh bags from early season. We then inoculated the strobili with eggs of *Cydia strobilella* in two different densities and later the cones with larvae of *Dioryctria abietella* also in two densities. We also inoculated them with a combination of the two species in order to see if that had any effect on seeds. The preliminary results mainly focus on seed yield measured as easily extractable seeds, since seeds damaged by insects often get stuck to the cone by resin or due to damaged scales.

---

**Keywords:** *Picea abies*, spruce seed orchard, insecticide, pest management, cone and seed insects