



**HAL**  
open science

# Early detection of invasive seed chalcids (*Megastigmus* spp.) through extensive surveys in major European botanical gardens: the example of rose seed chalcids

Marie-Anne Auger-Rozenberg, Alain Roques

## ► To cite this version:

Marie-Anne Auger-Rozenberg, Alain Roques. Early detection of invasive seed chalcids (*Megastigmus* spp.) through extensive surveys in major European botanical gardens: the example of rose seed chalcids. IUFRO 125th anniversary congress, division 7 “Forest Health”, Sep 2017, Freiburg, Germany. hal-03592993

**HAL Id: hal-03592993**

**<https://hal.inrae.fr/hal-03592993>**

Submitted on 31 Mar 2022

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# BOOK OF ABSTRACTS

# IUFRO

Interconnecting Forests,  
Science and People

125<sup>th</sup> Anniversary  
Congress 2017



## 125<sup>TH</sup> ANNIVERSARY CONGRESS 2017

18 – 22 September 2017  
Freiburg, Germany



[www.iufro2017.com](http://www.iufro2017.com)

125th IUFRO Anniversary Congress - Book of Abstracts, 2017. Freiburg. 724 p.

Published by Forstliche Versuchs- und Forschungsanstalt (FVA) Baden-Württemberg  
ISBN 978-3-902762-88-7

Copyright FVA and IUFRO.

The publication is available for download at:

<https://www.iufro.org/events/anniversary-congress/#c24907>

All Division 7 (Forest Health) Meeting

120 - Insect pests and pathogens of tree reproductive structures in a changing world: assessing

---

K 8 (Konzerthaus Freiburg)

IUFRO17-3053 **Messing with megagametophyte gene expression allows a seed parasite (*Megastigmus spermotrophus*) to harness the storage reserves of its host, Douglas-fir.**

von Aderkas, P.\* (1); Donaleshen, K. (1); Ehlting, J. (1)

(1) *University of Victoria, Forest Biology University of Victoria, Victoria, Canada*

**Abstract:** Douglas-fir seed is susceptible to a parasitic chalcid wasp that lays its eggs into megagametophytes during a period corresponding to central cell and egg maturation, i.e. immediately before and during fertilization. Whether a conifer egg aborts or is fertilized, the megagametophyte that houses the eggs never aborts: instead, it is manipulated by the larva. Four types of ovules/seeds were studied: 1. pollinated and uninfested, 2. pollinated and infested, 3. unpollinated and uninfested, and 4. unpollinated and infested. A de novo reference transcriptome was assembled in Trinity. Expression values were estimated based on the alignment of the original reads back onto the reference transcriptome using RSEM. Transcripts were annotated based on sequence similarity to genes of *Pinus taeda*, *Arabidopsis thaliana*, *Nasonia vitripennis*, and the UniProt database. Differential gene expression in Douglas-fir suggests that *M. spermotrophus* is capable of manipulating its host. Infested treatments had more transcripts related to seed storage, cell division and growth, solute transport, programmed cell death and hormone signaling that were similarly expressed in pollinated than unpollinated seeds. Although larval feeding triggers some genes involved in wounding, defense genes against herbivory are not up-regulated, meaning the insect achieves an uncontested access to the seed's storage reserves.

Douglas-fir, seed parasite, gene expression

---

K 8 (Konzerthaus Freiburg)

IUFRO17-880 **Early detection of invasive seed chalcids (*Megastigmus* spp.) through extensive surveys in major European botanical gardens: the example of rose seed chalcids**

Roques, A. (1); Auger-Rozenberg, M.-A.\* (1)

(1) *INRA, Zoologie Forestiere, Orleans, France*

**Abstract:** Botanical gardens at Paris National Museum of Natural History and Kew were sampled during 2013-2015 for hips of wild roses, *Rosa* spp., in order to check the possible presence of seed consumers on the shrubs planted in these urban areas. Depending on the year, the survey concerned 19- 32 species in Paris and 56-70 in Kew. More than 50000 seeds were X-rayed of which 13-28% were infested by insect larvae in Paris but ca. 6% in Kew. Besides *Megastigmus aculeatus*, a native seed chalcid present in the surroundings of both sites, adult emergence revealed the additional infestation by 4 alien species of *Megastigmus* which were not recorded until now in these regions. Two species, confirmed by molecular analyses, originated from North America, one from Eastern Europe, the origin of the last one being yet unknown. The exchange/trade of seeds is the most likely pathway. The presence of a large set of rose species also allowed to test for specificity of chalcids in relation to botanical sections within the genus *Rosa*. Gardens and arboreta thus proved to constitute a tool for early detection of alien seed insects. The study is expected to be extended to other gardens and arboreta of Eurasia, and to tree and shrub species

Rosa, seed, seed chalcid, invasion

---

K 8 (Konzerthaus Freiburg)

IUFRO17-2355 **Cone and seed insects-mediated impacts of global change on forest health: observations, predictions, and management options.**

Candau, J.-N.\* (1); Boivin, T. (2)

(1) *Canadian Forest Service, Natural Resources Canada, Sault Ste Marie, Canada*; (2) *Institut National de la Recherche Agronomique, Avignon, France*

**Abstract:** The vast majority of the scientific literature on the interactions between global change and forest pests has focused on insects that are directly or indirectly affecting tree survival. In comparison, the interactions with insects affecting tree reproduction, particularly cone and seed insects, have received little attention although seed availability is likely to severely constrain the response of tree populations to climate change. The current state of knowledge suggests that cone and seed insects can have unexpected impacts on the ability of tree populations to respond to a changing environment. Indeed, recent results show that insects can limit the spatial migration of forest tree populations hindering their capacity to track ecological niches as the environment is changing. The alternative strategy for persistence of tree populations under climate change, i.e. the adaptation to new conditions in current locations, can also be affected by insect seed predation although the mechanisms of interactions are even less well understood. We argue that our capacity to predict how these insects will respond to and interact with climate change is hampered by the shortage of information on their long-term impacts on the functioning and sustainability of forest ecosystems. In this presentation we will review how cone and seed insects are expected to respond and interact with climate change, how some of the complex interactions between these insects and climate change are already observed and what are the information gaps that limit our capacity to predict their future impacts. Finally, we will discuss a number of considerations in the management of these insects in a changing climate.

cone and seed insects, climate change, adaptation

---