



HAL
open science

The Genetic Control of Nectary Development

Filip Slavković, Catherine Dogimont, Halima Morin, Adnane Boualem,
Abdelhafid Bendahmane

► **To cite this version:**

Filip Slavković, Catherine Dogimont, Halima Morin, Adnane Boualem, Abdelhafid Bendahmane.
The Genetic Control of Nectary Development. Trends in Plant Science, 2021, 26 (3), pp.260-271.
10.1016/j.tplants.2020.11.002 . hal-03633112

HAL Id: hal-03633112

<https://hal.inrae.fr/hal-03633112v1>

Submitted on 13 Feb 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial 4.0 International License

1 **The genetic control of nectary development**

2

3 **Filip Slavković¹, Catherine Dogimont², Halima Morin¹, Adnane Boualem¹ and Abdelhafid**
4 **Bendahmane¹**

5

6 ¹Université Paris-Saclay, INRAE, CNRS, Univ Evry, Institute of Plant Sciences Paris-Saclay,
7 91405, Orsay, France

8 ²INRAE, UR 1052, Unité de Génétique et d'Amélioration des Fruits et Légumes, BP 94, F-
9 84143 Montfavet, France

10 Correspondence: abdelhafid.bendahmane@inrae.fr (A. Bendahmane).

11

12 **Keywords:** Nectary, nectar, floral, development, angiosperms, pollinators

13 **Abstract**

14 Nectar is the most important reward offered by flowering plants to pollinators for pollination
15 services. Since pollinator decline has emerged as a major threat for agriculture, and the food
16 demand is growing globally, studying nectar gland is of utmost importance. Although the
17 genetic mechanisms that control the development of angiosperm flowers have been quite
18 well understood for many years, the development, the maturation of nectar gland and the
19 secretion of nectar in synchrony with the maturation of the sexual organs appears to be one
20 of the flower's best kept secrets. Here we review key findings controlling these processes.
21 We also raise key questions that need to be addressed to develop crop ecological functions
22 that take into consideration pollinators' needs.

23 **Nectar and Food Security**

24 Can we imagine the world without chocolate, coffee or vanilla ice cream, three goods
25 derived from crops that depend on pollinators! In fact, animal pollinators are vital for life on
26 earth. Pollinators have co-evolved with flowering plants for millions of years, ensuring their
27 reproduction and keeping biodiversity and ecosystems alive. Insect pollinators are also a key
28 to agriculture, contributing to the production of most fruits and vegetables necessary for
29 healthy human diets. Over the last decades, there is mounting evidence of pollinator decline
30 all over the world and consequences in many agricultural areas could be a major threat [1,2].

31 This is severe, knowing that yields of 87 out of 115 (76%) leading global food crops and 35%
32 of global production depend on animal-mediated pollination [2,3]. From the economical
33 perspective, the pollination services provided by insect pollinators have an estimated value
34 of \$29 billion in the US alone [4,5] and \$153 billion per year [3] worldwide, equivalent to
35 9.5% of the total world agricultural food production [6].

36

37 In agriculture, the main insect pollinators, by far, are bees. Unfortunately, honeybee colonies
38 have decreased by 25% to 50% in Europe [7,8] and elsewhere [9,10]. Wild bees are also
39 declining, with one tenth of the species extinct or in danger of extinction in Europe [11,12].
40 While pollinator's declines leading to large scale losses of agricultural productivity or local
41 and temporal pollination crisis remain in debate, all the specialists in the field recognize the
42 importance of pollination services, supporting continued research and monitoring of
43 pollinator biodiversity [1,11]. From a breeding point of view, one central issue is whether the
44 cultivated varieties have been selected to reward pollinator services. To our knowledge, the
45 answer is clearly no. Plant domestication and genetic selection have enhanced yield and
46 improved nutritional values of harvested food and feed [13–15]. But crop ecological
47 functions, such as plant-pollinator interaction, have been largely ignored. This is perplexing,
48 knowing that half of the habitable land is used for agriculture [16]. Plant breeding also led to
49 a reduction in genetic diversity with high risk of losing traits beneficial to pollinators.

50 Nectar and pollen are the main rewards to pollinators. Floral nectar is produced by
51 specialized glands, called nectaries, in a process that is under complex developmental
52 control. Since bees prefer flowers with larger rewards, usually in the form of pollen and
53 nectar [17], investigating nectar related traits including nectary development and nectar
54 secretion will be key to develop "pollinator friendly" cultivars that not only increase yield
55 and its stability but also contribute to the rewarding and the preservation of the bees [18].
56 Investigation of nectaries as they develop and mature holds great potential to identify novel
57 targets to improve crop-pollinators interactions. In this review we will not address the
58 question of the pollinators themselves as this has been reviewed elsewhere [19]. We review
59 the roles and interactions of key genes controlling nectary development and recent inspiring
60 findings regarding the gene networks regulating nectary maturation and nectar secretion, in
61 various plant species. We also discuss how domestication and crop selection could impact

62 crop pollinator interactions and how new findings could help to breed varieties with
63 ecological functions for the benefit of pollinators and food security.

64 **Nectaries in flowering plants**

65 Nectaries are secretory structures that produce nectar, a carbohydrate rich solution
66 composed mainly of sugars, [20,21] which connects the plants with their pollinators and
67 defenders. Specifically, floral nectar is produced to attract pollinators, whereas extrafloral
68 nectar acts to defend plants indirectly [22]. As nectaries are highly variable in their
69 morphologies, anatomies and locations, they are defined based on their shared function: the
70 secretion of nectar [23]. Although there have been reports of nectaries in ferns [24], and in
71 Gnetales [25], nectaries are most widespread in angiosperms, in which there is a conserved
72 floral organ patterning. Nevertheless, nectaries seem to play by their own rules, as they can
73 be found in various floral and extrafloral positions [26]. In basal angiosperms, nectaries are
74 usually associated with the perianth [27], whereas in eudicots they are associated with
75 carpels and stamens. For example, in *Brassicaceae*, the nectaries are found at the base of
76 the stamens [28], in the *Solanaceae*, at the base of the gynoecium, and in *Malvaceae*, they
77 are found on the abaxial side of the involucre bracts as well as the adaxial side of sepals
78 [22].

79

80 **Models for studying nectary development**

81 Due to the lack of an ideal single model for nectary biology, nectary development has been
82 studied on a variety of species. The majority of studies on transcriptional and hormonal
83 regulation of nectary biology were done using *Arabidopsis thaliana*, an outstanding model in
84 terms of genetic and genomic resources [29]. However, in this case, biochemical aspects of
85 nectary function were limited due to the rather small nectary size (~100 microns wide and
86 deep), and extremely low volumes of nectar [29]. On the other hand, owing to much larger
87 flowers and nectar volumes, excellent studies on nectary metabolomics were conducted
88 using tobacco [30] and *Cucurbita pepo* [31–33]. Moreover, some of the bee visitation
89 experiments with respect to different floral traits were done using *Vicia faba* [15], while
90 *Sinningia speciosa* served as a useful tool for studying co-evolution of the flower shape and
91 pollinator visitation [34].

92

93 When it comes to studying nectary biology, species bearing unisexual flowers such as the
94 cucurbits (Figure 1) have particular advantages. First, they offer the opportunity to study
95 synchronization of the nectar secretion with the maturation of the sexual organ. Second,
96 they are practical to dissociate gland development from the development of the sexual
97 organs. Third, they are attractive owing to the relatively large size and volumes of the
98 nectary gland which facilitates their manipulation and makes them suitable for biochemical
99 analyses. Forth, the wide spread of sex determination morphs in the Cucurbitaceae plant
100 family compel for insect-mediated fertilization [35,36]. Nevertheless, each of these different
101 models has its advantages, and combined together they allow a systems approach for a
102 comprehensive understanding of the co-evolution of nectary biology and plant pollinators.

103

104 **The nectary development and the ABC(E) genes in *Arabidopsis thaliana***

105 In arabidopsis, floral nectary development begins at the base of the stamens in the third
106 **whorl** around stage 9, approximately 3.5 days before anthesis [37], comprising a receptacle
107 tissue with six glands on the abaxial side of the stamens [26]. From the onset of nectary
108 development, there are two distinct nectary cell types: an outer epidermal layer and an
109 inner starch granule-containing parenchymal tissue [26].

110 Despite great variation in morphology and size, floral organ order is conserved across
111 angiosperms [38]. Four concentric floral whorls are specified by the synchronous overlapping
112 actions of various transcription factors, commonly referred to as the ABC(E) genes.
113 Specifically, the A genes specify sepals, the A and B genes together specify petals, the B and
114 C genes together specify stamens, and the C genes specify carpels [37,39,40]. Since
115 mutations in single A-, B- and C-class homeotic mutants still develop nectaries, Baum and
116 colleagues initially proposed a model which suggested that the arabidopsis nectary is an
117 ABC-independent structure associated with the third whorl [26]. Later on, it was proposed
118 that B-, C- and E- (SEPALATA) functions are redundantly required for nectary development
119 [41,42].

120 Here, we will review to what extent could the ABC(E) genes affect the sizes and positions of
121 nectaries. Firstly, both arabidopsis and petunia plants that lack C lineage genes, do not
122 develop nectaries [42]. In addition, nectary gland development is reduced in the B- (*pi-1*,
123 *ap3*) and the C- (*ag-1*) class mutants, based on the absence of nectary tissue in *pi ag* and *ap3*

124 *ag* flowers of arabidopsis [41] (Table 1). However, the lack of nectary tissue in these mutants
125 may be due to the completely different organization of the floral whorls, rather than due to
126 a direct effect of the B- function genes [42]. Nevertheless, the nectaries of the *ap3* and *pi*
127 mutants show changes in size and morphology and thus our understanding of the role of the
128 B- class genes in nectary development remains incomplete. Secondly, mutations in the E-
129 class genes (*SEP*), which are required for B and C gene activity [43–45], also result in a failure
130 of nectary development. In addition to the regulation by B- and C- MADS box genes, two
131 other genes, *LEAFY* (*LFY*) and *UFO*, which regulate both the homeotic genes and the
132 formation of the third whorl, also affect nectary development [26]. In both *lfy* and *ufo* single
133 mutants, nectaries were rarely found, whereas in *lfy ufo* double mutants, no nectaries
134 develop [46], (Table 1). Furthermore, the experiment with the superman (*sup-1*) mutant, in
135 which the third whorl is repeated multiple times, showed that nectaries are associated with
136 each of the third whorls [26].

137 In conclusion, using genetic analyses with floral homeotic mutants, it has been
138 demonstrated that arabidopsis nectary is a third whorl structure [26] whose development
139 requires a C-class gene: either *AG*, or in its absence, ectopically expressed *SHP* genes are also
140 sufficient. Nevertheless, as different eudicot species have nectaries in different locations, the
141 outline for arabidopsis will not necessarily be the same in other species that display
142 nectaries at different positions.

143 ***CRABS CLAW* (*CRC*) and the ABC genes in nectary development**

144 The last year marked two decades of the publication of a paper identifying *CRABS CLAW*, the
145 only example of a single gene required for nectary gland development in arabidopsis, as no
146 morphological or molecular signs of nectaries are observed in *crc* mutants [26]. In addition,
147 *CRC* is also implicated in FM determinacy and carpel development [47]. As a member of the
148 *YABBY* protein family, *CRC* is characterized by a C₂C₂ zinc finger domain located at N-
149 terminus and a helix-loop-helix motif (*YABBY* domain) at the C terminus which is similar to
150 the high mobility group (HMG) box motif [46]. Restricted to nectaries and carpels by the
151 action of the floral-meristem identity genes *AP1*, *LFY* and *UFO* [46], *CRC* expression
152 commences before the nectaries emerge and continues until after anthesis. Already from
153 stage 6 of flower development on, *CRC* expression occupies an almost continuous ring of

154 receptacle cells between the stamen and sepal primordia, including regions where nectaries
155 will develop, suggesting that *CRC* plays a role in the early specification of cells that will
156 become nectaries [46]. Using the ABC homeotic mutants, Baum and colleagues (2001)
157 demonstrated that *CRC* mRNA expression is negatively controlled by A and B functions in the
158 outer and the third whorl respectively, but can occur independently of C function outside
159 the third whorl. Although ectopic expression of *CRC* is not sufficient to induce ectopic
160 nectaries, *CRC* is necessary for nectary development in various genetic backgrounds,
161 indicating that it is one of the key genes directing nectary development in arabidopsis [26].

162

163 More recently, Gross et al. (2018) demonstrated that *CRC* forms homodimers and
164 heterodimers with *INO*, a member of the same protein family via the YABBY domain.
165 However, this interaction should not control the nectary development, but may control
166 other functions such as petal and sepal development and leaf structure as the two genes are
167 co-expressed in these tissues. Furthermore, their work showed that *CRC* has two distinct
168 functions: 1) it is involved in floral meristem termination via transcriptional repression, and
169 2) it acts as a transcriptional activator in nectary development and carpel fusion and growth
170 control [48].

171

172 ***CRC* and the ABC(E) genes: things get complicated**

173 The next question was how does *CRC* fit into the ABC(E) floral genes puzzle. Lee and
174 colleagues (2005) identified the *CRC* promoter region that is necessary and sufficient for
175 proper *CRC* expression [49]. They found it harbors two CArG [CC(A/T)6GG] boxes, known
176 binding sites for MADS box proteins. This section will review how MADS box proteins
177 *AGAMOUS*, *SHATTERPROOF1/2*, *PISTILLATA* and *SEPALLATA1/2/3* regulate *CRC* expression
178 and nectary development. Firstly, a study by Wuest and colleagues has shown that TFs AP3
179 and PI directly suppress the expression of *CRC* [50]. Secondly, mutations in the *SEP* genes,
180 which are redundantly required to specify petals, stamens and carpels [43–45,51,52], result
181 in a failure of nectary development despite having a third whorl, and are therefore required
182 for *CRC* activation in the third whorl. Thirdly, in the absence of B- and C-class gene activities,
183 *SHATTERPROOF1* (*SHP1*) and *SHP2*, which encode proteins similar to *AG*, might rescue
184 nectary development, if they are ectopically expressed, as in an A-class *ap2* mutant

185 background [42,49,53,54]. However, *ag shp1 shp2* triple mutants do not develop nectaries,
186 while nectaries still develop in *ag* and in *shp1 shp2* mutants (Table 1), suggesting that
187 SHP1/2 may not need to be in an *ap2* background to be functional [42].

188 All these experiments led to a conclusion that the C- and E- class gene activities are
189 redundantly required for *CRC* activation and nectary development. The lack of nectary
190 formation in BC double mutants, but their presence in B and C single mutants, would suggest
191 redundancy of these proteins in the complex with the SEP proteins [49]. Nevertheless, as the
192 architecture of these mutants is highly modified, it remains to be determined whether the
193 lack of nectaries is a direct effect. Finally, bearing in mind that the function and the
194 expression domain of the C- lineage genes and *CRC* is much broader than the nectary
195 development, the restriction of nectaries at the base of carpels in petunia, and at the base of
196 stamens in arabidopsis must depend on the presence of additional local genetic factors [42].
197 Moreover, it is important to note that the genetic analyses were carried out in eudicots, and
198 thus extrapolations from this data should be limited to these taxa. Furthermore, *CRC*'s DNA
199 binding motif and target genes in developing nectaries still need to be identified.

200 More recently, *CRC* was reported to bind promoter regions of *3-KETOACYL-COA SYNTHASE 7*
201 and *15 (KCS7 and KCS15)*, two genes that are involved in the synthesis of fatty acids [55],
202 which are then used as signaling molecules or in cuticular wax synthesis [56]. This report
203 sheds some new light on our understanding of *CRC* which seems to control other important
204 biological processes.

205

206 **Functional conservation of *CRC* in flowering plants**

207 What do we know on the identified regulators of nectary development in flowering plants?
208 Most of the rosids and asterid species have their nectaries associated either with stamens or
209 carpels [57]. Bearing in mind that both positions occupy the C-function domain and the fact
210 that *CRC* expression in nectaries has been shown to be conserved in a number of higher
211 eudicot species [58], Morel and colleagues suggested that floral nectary development in
212 rosids and asterids generally occurs via the C-lineage/*CRC* module [42]. Since all C-lineage
213 genes from petunia (*pMADS3* and *FBP6*) and arabidopsis (*AG*, *SHP1* and *SHP2*) are able to
214 activate *CRC* expression, this suggests that C-lineage gene dependent *CRC* activation already
215 existed before the split between rosids and asterids. This could further suggest a common

216 evolutionary origin for nectary development in the two major core eudicot lineages, at least
217 for species in which the nectaries are associated with the reproductive organs, and thus
218 residing within the classical C-function expression domain [42].

219 Interestingly, outside of the flower, *CRC* expression was also detected in extrafloral nectaries
220 of the rosid species *Capparis flexuosa*, as well as in nectaries that develop from the midvein
221 of leaves and on the involucre bracts in *Gossypium hirsutum* [58]. On the other hand, in
222 basal eudicot species, no evidence of *CRC* expression in nectaries was found in *Aquilegia*
223 *formosa* and *Epimedium sagittatum*, [58,59]. Nevertheless, Min and colleagues filled this gap
224 by demonstrating that in those two genera, nectary development is controlled by
225 the *STYLISH*-like genes instead, which encode a group of plant specific TFs that are required
226 for carpel fusion and the correct development of the style and stigma in arabidopsis [60,61].
227 In addition, their work showed that the expression of the *STY1* homologs is closely
228 associated with nectaries in the divergent members of both the *Ranunculaceae* and
229 *Berberidaceae*, both basal eudicots [60]. All these examples show that nectary development
230 may require *CRC* function, but its activation may not necessarily depend on C-lineage genes,
231 or that nectary development can even occur independently of *CRC* [42], which is in line with
232 the hypothesis that nectaries evolved multiple times independently [42,62].

233

234 While the involvement of *CRC* in carpel development was present in the ancestral
235 angiosperms [63], its involvement in nectary development, at least on present data, may be
236 restricted to the eudicots. [58,59]. Finally, the observation that nectaries are absent from
237 the flowers of ANITA (*Amborella*, *Nymphaeales*, *Illiciaceae*, *Trimeniaceae*,
238 *Austrobaileyaceae*) grade angiosperms [64] may suggest that nectaries evolved after the
239 separation of the ANITA clades from the remaining lineage [63].

240

241 ***BLADE ON PETIOLE (BOP)* gene and nectary development**

242 Similar to *CRC*, *BLADE ON PETIOLE1* and *BOP2* play an important role in nectary development
243 in arabidopsis. *BOP1* and *BOP2* are part of the NPR1 (NON-EXPRESSOR OF PR1) protein
244 family, which is characterized by a series of conserved cysteines and two protein-protein
245 interaction domains [51]. *BOP1/2* are expressed in undifferentiated cells at the base of
246 developing lateral organs and are needed to repress indeterminate growth and promote

247 differentiation in the proximal regions of lateral organs [51,65–67]. Localized to the cytosol
248 and nucleus [65,68], BOP1/2 can form homo- and heterodimers [69] and interact with the
249 TGA transcription factor, PERIANTHIA (PAN) [70] (Figure 2), mutations of which affect the
250 floral organ number in the first three whorls [71]. Unlike the *crc* mutant which completely
251 lacks nectaries, in *bop1/bop2* double mutant, nectaries are not entirely absent but rather
252 reduced in size and do not differentiate key nectary features such as parenchymal and
253 secretory tissue [72]. Similarly to *CRC*, *BOP* is expressed very early in nectary development
254 and may be controlling other downstream elements in conjunction with *CRC* [51]. Moreover,
255 phenotyping of *bop 1 bop 2 pan3* triple mutants revealed that BOP1/2 and PAN function in
256 the same genetic pathway and have a joint role in abaxial patterning of the floral meristem,
257 as no additive or synergistic increase in patterning defects was observed in these mutants
258 [70]. Taken together, it has been proposed that, once induced by an appropriate signal, BOP
259 proteins may interact in the nucleus with TGA transcription factors, such as PAN, to regulate
260 the transcription of floral patterning genes. The relationship between *BOP1/2* and other
261 floral homeotic genes such as *AG* is, however, lacking. Analysis of mutants, involving for
262 instance *bop1/bop2* and *ag*, could answer whether or not other floral homeotic genes feed
263 into the *BOP1/2*-dependent pathway of nectary development.

264 **Hormone signaling and nectary development**

265 Hormone action is often mediated by transcription factors such as the auxin response
266 factors (ARFs), some of which are microRNA (miRNA) regulated. For example, AUXIN
267 RESPONSE FACTOR 6 (ARF6) and ARF8, which are the cleavage targets of the
268 microRNA *miR167* [73], act redundantly to promote and coordinate maturation of nectary,
269 petal, stamen and gynoecium [74,75]. In *arf6/arf8* double mutants, nectaries are very small
270 and only detectable in a fraction of flowers, indicating that auxin signaling pathways are
271 required for proper nectary growth and function [76,77]. In addition, ARF6 and ARF8 have
272 been shown to activate jasmonate biosynthesis, which in turn activates MYB21 and MYB24
273 which are also expressed in nectaries. However, unlike ARF6 and ARF8, morphological and
274 gene expression analyses showed that MYB21 and MYB24 only affect nectary gene
275 expression, but not nectary formation [73]. Specifically, the *myb21/myb24* flowers have
276 reduced expression of arabidopsis terpene synthase genes *TPS11* and *TPS21*. In addition,
277 MYB21 was shown to promote the production of volatile sesquiterpenes, and together with

278 MYB24, to mediate secondary jasmonate responses in stamens, which may attract
279 pollinators and/or repel pathogens [73]. MYB21 also feeds back negatively on expression of
280 jasmonate biosynthesis pathway genes to decrease flower JA level, which also correlates
281 with termination of growth after the flowers have opened [73]. However, analysis of
282 jasmonate insensitive mutant *coi1-1* revealed no phenotypic alteration of nectaries,
283 excluding the possibility that JA regulates nectary development[76–78].

284

285 **Hormone signaling and nectar secretion**

286 The role of hormonal signaling in nectary development has not been studied in detail.
287 Nevertheless, auxin, gibberellin and JA have been reported to play important roles in
288 regulating nectar production [79–81]. For example, *PIN6*, that encodes an auxin efflux
289 transporter family protein, is a nectary-enriched gene whose expression is positively
290 correlated with total nectar production [79]. Moreover, plants with the knocked-out
291 *GA2OX6*, a gene encoding the enzyme that catalyzes inactivation of bioactive GAs, have
292 elevated levels of bioactive GAs, which, leads to decreased expression of genes involved in
293 nectar production, including *PIN6* [80]. In addition, there are nine other nectary-enriched
294 genes whose expression were reported to depend on GAs signaling [80] (Table 2).

295

296 According to the current model of nectar secretion (reviewed in [82]), (Figure 3), GAs
297 endogenous to nectaries negatively regulate nectar production [80], whereas GAs from
298 other floral tissues (developing stamens) seem to indirectly regulate nectary function
299 through induction of JA-mediated responses, which likely diffuse to nectaries to induce auxin
300 production in a positive feedback loop [73]. As a result, auxin may induce ARF6/8 expression
301 and lead to the expression of the MYB21/MYB24 genes which are required for nectary
302 maturation and function [73]. Moreover, jasmonate insensitive tobacco plants with silenced
303 *NtCOI1* gene have nectarless phenotype and were reported to act upstream of MYB305 [83],
304 a gene which plays a critical role in starch metabolism and nectar production [30].
305 Interestingly, ectopic expression of MYB305 in tobacco leaves was able to induce expression
306 of the *nec1* [84], the only reported gene that controls the development of extrafloral
307 nectaries in cotton [85]. In sum, JA, in interaction with other hormones, plays a central role
308 in the coordination of the maturation of the nectaries, stamen, gynoecium and petals, all to
309 attract pollinators, when the flower is competent for reproduction.

310

311 **Nectar synthesis and secretion**

312 A recent study on *Cucurbita pepo* nectary identified key genes in nectar synthesis and
313 secretion during starch synthesis (*STARCH BRANCHING ENZYME* - *CpSBE2*), starch
314 degradation (*BETA AMYLASE* - *CpBAM1*), sucrose synthesis (*SUCROSE PHOSPHATE SYNTHASE*
315 - *CpSPS*), sucrose export (*CpSWEET9*) and sucrose hydrolysis (*CELL WALL INVERTASE4* –
316 *CpCWIN4*) [31]. A common theme in nectar synthesis and secretion in different species is the
317 transformation of the starch breakdown products into sucrose by the action of the SPSs and
318 sucrose phosphate phosphatases (SPP), after which sucrose is exported from the nectary
319 cells in a concentration dependent manner via uniporter SWEET9 (Figure 3). However, the
320 final step of sucrose hydrolysis by CWIN4 seems to be species-specific and it might play
321 different roles. For example, in the hexose-rich nectar of arabidopsis, CWIN4 generates a
322 concentration gradient to drive sugar export, while in the sucrose-rich nectar of *C. pepo* [33]
323 its role is likely in dictating the final nectar quality [31].

324

325 In a study on ornamental tobacco, in which flower development is divided into 12 stages
326 [86], nectary starch degradation (20% by mass) was shown to rapidly produce a large
327 amount of glucose between stage S9 of flower development, characterized by enlarging of
328 the corolla tube, and S12 (anthesis) doubling the physiological cellular osmolarity (~300
329 Osm). This increase in cellular osmolarity will lead to a dramatic decline in the water
330 potential, triggering influx of water from the phloem via sieve elements. As a result, the
331 increased hydrostatic pressure within nectary causes nectar to exude through nectary pores
332 [87]. This report suggested that two processes, starch degradation and rapid sugar influx, are
333 determinants of sugar composition in floral nectar [87].

334

335 **Nectary transcriptome: There is more to nectaries than TFs, but not much more**

336 Global transcriptomics analyses of nectaries have been studied in several species [31,88–91].
337 The first report of a nectary transcriptome study was performed by Kram and colleagues
338 (2009), who identified 270 genes preferentially expressed in arabidopsis nectaries [88].
339 Interestingly, the short list of the nectary-enriched genes studied by Reeves and colleagues
340 showed that 18 genes were underrepresented in *arf6-2 arf8-3* mutants [73]. Among them
341 were *CRC*[46]; *YABBY5*, encoding a protein closely related to *CRC*; *CWIN4*, encoding a cell

342 wall invertase required for nectary sink strength and nectar production [92]; *SWEET9*,
343 encoding a nectary-specific glucose transporter [93,94]; and *JMT* encoding S-adenosyl-L-
344 methionine jasmonic acid carboxyl methyltransferase, which makes the volatile compound
345 methyl jasmonate [95]. Interestingly, each of these genes was underrepresented in both
346 *arf6-2 arf8-3* and *myb21-5 myb24-5* flowers, except for *CRC* which was underrepresented in
347 *arf6-2 arf8-3* flowers only, suggesting the requirement of auxin signaling for CRC-mediated
348 pathways.

349

350 A recent RNA-seq analysis demonstrated that CRC together with SUPERMAN (SUP), a gene
351 that encodes a C₂C₂-type zinc-finger protein involved in FM termination, coordinate
352 hormone-, stress-, and metabolic gene expression in stamen development [96]. This global
353 transcriptomic study identified and selected 263 differentially expressed genes in the *crc*
354 mutant which could help us better understand its roles in other biological processes [96].
355 Moreover, it seems that genes involved in auxin and gibberellin signaling might play
356 significant roles in nectary development and further research using candidate gene
357 approaches as well as 'OMICS' analyses are required to validate this hypothesis.

358 **Concluding Remarks and Future Perspectives**

359 Despite the central role of nectar glands in the interaction between flowering plants and
360 pollinators and the numerous investigations, we have only scratched the surface of
361 molecular mechanisms controlling their development. The knowledge is so limited today
362 that no breeding program can be conceived to favour the preservation of pollinators and to
363 improve fruit set. With the recent development of single cell omics technologies,
364 metabolome profiling, precise phenotyping and low cost of genome sequencing we can
365 foresee projects that tackle the interaction of plant and pollinators at the flower level, to
366 identify key genes controlling nectary development and nectar metabolism and secretion, as
367 well as at the population level, to bring new insights on the heritability and the variability of
368 the traits. Phenotyping of cultivated accessions, land races and related wild species for
369 pollinator foraging activities will also permit to investigate whether domestication have
370 filtered, in or out, certain nectar-related traits. Moreover, evo-devo analyses of the
371 identified genes shaping plant-pollinators interactions could help better understand the role

372 and the relationship of the controlled phenotypes in the context of the co-evolution of the
373 plant with the pollinators.

374 **Acknowledgements**

375 The authors would like to thank Judit Szecsi and Manash Chatterjee for careful proofreading
376 of the manuscript and the reviewers for constructive suggestions. Financial support was
377 provided by the ANR Nectar project (ANR-19-CE20), LabEx Saclay Plant Sciences - SPS (ANR-
378 10-LABX-40-SPS) and the Plant Biology and Breeding Department of INRAE.

379

- 380 1 Science for environment policy (2020) pollinators: importance for nature and human well-being,
381 drivers of decline and the need for monitoring. Future Brief 23. Brief produced for the European
382 Commission DG Environment. Bristol: Science communication unit, UWE Bristol.
- 383 2 Klein, A.-M. *et al.* (2007) Importance of pollinators in changing landscapes for world crops. *Proc. R.*
384 *Soc. B Biol. Sci.* 274, 303–313
- 385 3 Garibaldi, L. *et al.* (2009) Pollinator shortage and global crop yield: Looking at the whole spectrum
386 of pollinator dependency. *Commun Integr Biol* 2, 37–39
- 387 4 Roy, R. *et al.* (2017) Review: Nectar biology: From molecules to ecosystems. *Plant Sci.* 262, 148–
388 164
- 389 5 Calderone, N.W. (2012) Insect Pollinated Crops, Insect Pollinators and US Agriculture: Trend
390 Analysis of Aggregate Data for the Period 1992–2009. *PLoS ONE* 7, e37235
- 391 6 Gallai, N. *et al.* (2009) Economic valuation of the vulnerability of world agriculture confronted with
392 pollinator decline. *Ecol. Econ.* 68, 810–821
- 393 7 Potts, Simon Geoffrey, S.G. *et al.* (2010) Declines of managed honey bees and beekeepers in
394 Europe. *J Apic Res* 49, 15–22
- 395 8 Gray, A. *et al.* (2019) Loss rates of honey bee colonies during winter 2017/18 in 36 countries
396 participating in the COLOSS survey, including effects of forage sources. *J Apic Res* 58, 479–485
- 397 9 National Resource Council, Status of Pollinators in North America. . (2007) , National Academies
398 Press, Washington, DC,
- 399 10 van Engelsdorp *et al.* (2008) A survey of honey bee colony losses in the U.S., fall 2007 to spring
400 2008. *PLOS ONE* 3,
- 401 11 Biesmeijer, J. *et al.* (2006) Parallel Declines in Pollinators and Insect-Pollinated Plants in Britain
402 and the Netherlands. *Science* 313, 351–354
- 403 12 Nieto, A. *et al.* European Red List of bees. Luxembourg: Publication Office of the European Union.
404 . (2014)
- 405 13 Kobayashi, K. *et al.* (2010) Selective flower visitation behavior by pollinators in a radish F1 seed
406 production field. *Breed. Sci.* 60, 203–211
- 407 14 Richards, R.A. (2000) Selectable traits to increase crop photosynthesis and yield of grain crops. *J.*
408 *Exp. Bot.* 51, 447–458
- 409 15 Bailes, E.J. *et al.* (2018) An analysis of the energetic reward offered by field bean (*Vicia faba*)
410 flowers: Nectar, pollen, and operative force. *Ecol. Evol.* 8, 3161–3171
- 411 16 Ellis, E. *et al.* (2010) Anthropogenic transformation of the biomes, 1700 to 2000. *Glob. Ecol.*
412 *Biogeogr.* 19, 589–606
- 413 17 Brunet, J. *et al.* (2015) The Effects of Flower, Floral Display, and Reward Sizes on Bumblebee
414 Foraging Behavior When Pollen Is the Reward and Plants Are Dichogamous. *Int. J. Plant Sci.* 176,
415 811–819

- 416 18 Bailes, E.J. *et al.* (2015) How can an understanding of plant-pollinator interactions contribute to
417 global food security? *Curr. Opin. Plant Biol.* 26, 72–79
- 418 19 Prasifka, J.R. *et al.* (2018) Using Nectar-Related Traits to Enhance Crop-Pollinator Interactions.
419 *Front. Plant Sci.* 9, 812
- 420 20 Fahn, A. (1988) Secretory tissues in vascular plants. *New Phytol.* 108, 229–257
- 421 21 Schmid, R. (1988) Reproductive versus extra-reproductive nectaries—historical perspective and
422 terminological recommendations. *Bot. Rev.* 54, 179–227
- 423 22 Elias, T.S. (1983) Extrafloral nectaries: their structure and distribution. In *The Biology of Nectaries*
424 pp. 174–203, B. Bentley and T.S. Elias
- 425 23 Fahn, A. (1979) , Ultrastructure of nectaries in relation to nectar secretion. , *American Journal of*
426 *Botany*, 977–985
- 427 24 Darwin, F. (1877) On the nectar glands of the common brakeferns. *Bot. J. Linn. Soc.* 15, 407–409
- 428 25 PORSCHE, O. (1910) 57. O t t o P o r s c h: Ephedra campylopoda C. A. Mey., eine entomophile
429 Gymnosperme.
- 430 26 Baum, S.F. *et al.* (2001) The Arabidopsis nectary is an ABC-independent floral structure.
431 *Development* 128, 4657–4667
- 432 27 Endress, P.K. (2001) Origins of flower morphology. *J. Exp. Zool.* 291, 105–115
- 433 28 Arber, A. (1931) STUDIES IN FLORAL MORPHOLOGY. I. ON SOME STRUCTURAL FEATURES OF THE
434 CRUCIFEROUS FLOWER1. *New Phytol.* 30, 11–41
- 435 29 Kram, B.W. and Carter, C.J. (2009) Arabidopsis thaliana as a model for functional nectary
436 analysis. *Sex. Plant Reprod.* 22, 235–246
- 437 30 Liu, G. and Thornburg, R.W. (2012) Knockdown of MYB305 disrupts nectary starch metabolism
438 and floral nectar production: MYB305 regulates nectary starch metabolism. *Plant J.* 70, 377–388
- 439 31 Solhaug, E.M. *et al.* (2019) An integrated transcriptomics and metabolomics analysis of the
440 *Cucurbita pepo* nectary implicates key modules of primary metabolism involved in nectar
441 synthesis and secretion. *Plant Direct* 3, e00120
- 442 32 Chatt, E.C. *et al.* (2018) Sex-Dependent Variation of Pumpkin (*Cucurbita maxima* cv. Big Max)
443 Nectar and Nectaries as Determined by Proteomics and Metabolomics. *Front. Plant Sci.* 9, 860
- 444 33 Nepi, M. *et al.* (2001) Nectar Secretion, Reabsorption, and Sugar Composition in Male and
445 Female Flowers of *Cucurbita pepo*. *Int. J. Plant Sci.* 162, 353–358
- 446 34 Wang, C.-N. *et al.* (2015) Quantifying floral shape variation in 3D using microcomputed
447 tomography: a case study of a hybrid line between actinomorphic and zygomorphic flowers.
448 *Front. Plant Sci.* 6,
- 449 35 Boualem, A. *et al.* (2008) A Conserved Mutation in an Ethylene Biosynthesis Enzyme Leads to
450 Andromonoecy in Melons. *Science* 321, 836–838
- 451 36 Boualem, A. *et al.* (2015) A cucurbit androecy gene reveals how unisexual flowers develop and
452 dioecy emerges. *Science* 350, 688–691
- 453 37 Smyth, D.R. *et al.* (1990) Early flower development in Arabidopsis. *Plant Cell* 2, 755–767
- 454 38 Bowman, J.L. *et al.* (2012) The ABC model of flower development: then and now. *Development*
455 139, 4095–4098
- 456 39 Coen, E.S. and Meyerowitz, E.M. (1991) , The war of the whorls: genetic interactions controlling
457 flower development. , *Nature*, 31–37
- 458 40 Weigel, D. and Meyerowitz, E.M. (1994) The ABCs of floral homeotic genes. *Cell* 78, 203–209
- 459 41 Lee, J.-Y. *et al.* (2005) Activation of *CRABS CLAW* in the Nectaries and Carpels of Arabidopsis.
460 *Plant Cell* 17, 25–36
- 461 42 Morel, P. *et al.* (2018) The Floral C-Lineage Genes Trigger Nectary Development in Petunia and
462 Arabidopsis. *Plant Cell* 30, 2020–2037
- 463 43 Pelaz, S. *et al.* (2000) , B and C floral organ identity functions require SEPALLATA MADS-box
464 genes. , *Nature*, 200–203
- 465 44 Pelaz, S. *et al.* (2001) , Conversion of leaves into petals in Arabidopsis. , *Current Biology*, 182–184
- 466 45 Honma, B. and Goto, K. (2001) Complexes of MADS-box proteins are sufficient to convert leaves
467 into floral organs. *Nature* 409, 525–529

- 468 46 Bowman, J.L. and Smyth, D.R. (1999) CRABS CLAW, a gene that regulates carpel and nectary
469 development in Arabidopsis, encodes a novel protein with zinc finger and helix-loop-helix
470 domains. *Development* 126, 2387–2396
- 471 47 Alvarez, J. and Smyth, D.R. (1999) CRABS CLAW and SPATULA, two Arabidopsis genes that control
472 carpel development in parallel with AGAMOUS. *Development* 126, 2377
- 473 48 Gross, T. *et al.* (2018) CRABS CLAW Acts as a Bifunctional Transcription Factor in Flower
474 Development. *Front. Plant Sci.* 9, 835
- 475 49 Lee, J.-Y. (2005) Activation of CRABS CLAW in the Nectaries and Carpels of Arabidopsis. *PLANT
476 CELL ONLINE* 17, 25–36
- 477 50 Wuest, S.E. *et al.* (2012) Molecular basis for the specification of floral organs by APETALA3 and
478 PISTILLATA. *Proc. Natl. Acad. Sci.* 109, 13452
- 479 51 McKim, S.M. *et al.* (2008) The BLADE-ON-PETIOLE genes are essential for abscission zone
480 formation in Arabidopsis. *Development* 135, 1537–1546
- 481 52 Hugouvieux, V. *et al.* (2018) Tetramerization of MADS family transcription factors SEPALLATA3
482 and AGAMOUS is required for floral meristem determinacy in Arabidopsis. *Nucleic Acids Res.* 46,
483 4966–4977
- 484 53 Savidge, B. *et al.* (1995) , Temporal relationship between the transcription of two Arabidopsis
485 MADS box genes and the floral organ identity genes. , *The Plant Cell*, 721–733
- 486 54 Pinyopich, A. *et al.* (2003) Assessing the redundancy of MADS-box genes during carpel and ovule
487 development. *Nature* 424, 85
- 488 55 Han, X. *et al.* (2012) Co-expression Analysis Identifies CRC and AP1 the Regulator of Arabidopsis
489 Fatty Acid Biosynthesis. *J. Integr. Plant Biol.* 54, 486–499
- 490 56 Joubès, J. *et al.* (2008) The VLCFA elongase gene family in Arabidopsis thaliana: phylogenetic
491 analysis, 3D modelling and expression profiling. *Plant Mol. Biol.* 67, 547–566
- 492 57 Bernardello, G. (2007) A systematic survey of floral nectaries. In *Nectaries and nectar* Nicolson
493 S.W., Nepi M., Pacini E.pp. 19–128, Springer
- 494 58 Lee, J.-Y. (2005) Recruitment of CRABS CLAW to promote nectary development within the
495 eudicot clade. *Development* 132, 5021–5032
- 496 59 Sun, W. *et al.* (2013) Characterization of a Crabs Claw Gene in Basal Eudicot Species *Epimedium*
497 *sagittatum* (Berberidaceae). *Int. J. Mol. Sci.* 14, 1119–1131
- 498 60 Min, Y. *et al.* (2019) Homologs of the *STYLISH* gene family control nectary development in
499 *Aquilegia*. *New Phytol.* 221, 1090–1100
- 500 61 Kuusk, S. *et al.* (2002) STY1 and STY2 promote the formation of apical
501 tissues during Arabidopsis gynoecium development. *Development* 129, 4707
- 502 62 Brown, W.H. (1938) The bearing of nectaries on the phylogeny of flowering plants. *Proc. Am.
503 Philos. Soc.* 79, 549–595
- 504 63 Fourquin, C. *et al.* (2005) Evidence that CRABS CLAW and TOUSLED have conserved their roles in
505 carpel development since the ancestor of the extant angiosperms. *Proc. Natl. Acad. Sci. U. S. A.*
506 102, 4649
- 507 64 Endress, P.K. (2010) The evolution of floral biology in basal angiosperms. *Philos. Trans. R. Soc.
508 Lond. B. Biol. Sci.* 365, 411–421
- 509 65 Hepworth, S.R. *et al.* (2005) BLADE-ON-PETIOLE–Dependent Signaling Controls Leaf and Floral
510 Patterning in Arabidopsis. *Plant Cell* 17, 1434–1448
- 511 66 Norberg, M. (2005) The BLADE ON PETIOLE genes act redundantly to control the growth and
512 development of lateral organs. *Development* 132, 2203–2213
- 513 67 Ha, C.M. *et al.* (2007) *BLADE-ON-PETIOLE1* and 2 Control *Arabidopsis* Lateral Organ Fate through
514 Regulation of LOB Domain and Adaxial-Abaxial Polarity Genes. *Plant Cell* 19, 1809–1825
- 515 68 Jun, J.H. *et al.* (2010) *BLADE-ON-PETIOLE1* Coordinates Organ Determinacy and Axial Polarity in
516 *Arabidopsis* by Directly Activating *ASYMMETRIC LEAVES2*. *Plant Cell* 22, 62–76
- 517 69 Jun, J.H. *et al.* (2010) *BLADE-ON-PETIOLE1* Coordinates Organ Determinacy and Axial Polarity in
518 *Arabidopsis* by Directly Activating *ASYMMETRIC LEAVES2*. *Plant Cell* 22, 62–76

519 70 Hepworth, S.R. (2005) BLADE-ON-PETIOLE-Dependent Signaling Controls Leaf and Floral
520 Patterning in Arabidopsis. *PLANT CELL ONLINE* 17, 1434–1448

521 71 Chuang, C.-F. *et al.* (1999) The PERIANTHIA gene encodes a bZIP protein involved in the
522 determination of floral organ number in Arabidopsis thaliana. *Genes Dev.* 13, 334–344

523 72 McKim, S.M. *et al.* (2008) The BLADE-ON-PETIOLE genes are essential for abscission zone
524 formation in Arabidopsis. *Development* 135, 1537–1546

525 73 Reeves, P.H. *et al.* (2012) A Regulatory Network for Coordinated Flower Maturation. *PLoS Genet.*
526 8, e1002506

527 74 Nagpal, P. (2005) Auxin response factors ARF6 and ARF8 promote jasmonic acid production and
528 flower maturation. *Development* 132, 4107–4118

529 75 Wu, M.-F. *et al.* (2006) Arabidopsis microRNA167 controls patterns of ARF6 and ARF8 expression,
530 and regulates both female and male reproduction. *Development* 133, 4211–4218

531 76 Reeves, P.H. *et al.* (2012) A Regulatory Network for Coordinated Flower Maturation. *PLoS Genet.*
532 8, e1002506

533 77 Nagpal, P. (2005) Auxin response factors ARF6 and ARF8 promote jasmonic acid production and
534 flower maturation. *Development* 132, 4107–4118

535 78 Schmitt, A.J. *et al.* (2018) The octadecanoid pathway, but not COI1, is required for nectar
536 secretion in Arabidopsis thaliana. *Front. Plant Sci.* 9,

537 79 Bender, R.L. *et al.* (2013) PIN6 is required for nectary auxin response and short stamen
538 development. *Plant J.* 74, 893–904

539 80 Wiesen, L.B. *et al.* (2016) A role for GIBBERELLIN 2-OXIDASE6 and gibberellins in regulating
540 nectar production. *Mol. Plant* 9, 753–756

541 81 Radhika, V. *et al.* (2010) Towards elucidating the differential regulation of floral and extrafloral
542 nectar secretion. *Plant Signal. Behav.* 5, 924–926

543 82 Roy, R. *et al.* (2017) Review: Nectar biology: From molecules to ecosystems. *Plant Sci.* 262, 148–
544 164

545 83 Wang, W. *et al.* (2014) The F-box protein COI1 functions upstream of MYB305 to regulate
546 primary carbohydrate metabolism in tobacco (*Nicotiana tabacum* L. cv. TN90). *J. Exp. Bot.* 65,
547 2147–2160

548 84 Liu, G. *et al.* (2009) The MYB305 Transcription Factor Regulates Expression of Nectarin Genes in
549 the Ornamental Tobacco Floral Nectary. *Plant Cell* 21, 2672–2687

550 85 Hu, W. *et al.* (2020) Genetic and evolution analysis of extrafloral nectary in cotton. *Plant*
551 *Biotechnol. J.* DOI: 10.1111/pbi.13366

552 86 Koltunow, A.M. *et al.* (1990) Different Temporal and Spatial Gene Expression Patterns Occur
553 during Anther Development. *Plant Cell* 2, 1201

554 87 Ren, G. *et al.* (2007) Transient starch metabolism in ornamental tobacco floral nectaries
555 regulates nectar composition and release. *Plant Sci.* 173, 277–290

556 88 Kram, B.W. *et al.* (2009) Uncovering the Arabidopsis thaliana nectary transcriptome:
557 investigation of differential gene expression in floral nectariferous tissues. *BMC Plant Biol.* 9, 92

558 89 Hampton, M. *et al.* (2010) Identification of Differential Gene Expression in Brassica rapa
559 Nectaries through Expressed Sequence Tag Analysis. *PLoS ONE* 5, e8782

560 90 Thomas, J.B. *et al.* (2017) The pennycress (*Thlaspi arvense* L.) nectary: structural and
561 transcriptomic characterization. *BMC Plant Biol.* 17, 201

562 91 Liu, H. *et al.* (2019) Transcriptomic and microstructural analyses in *Liriodendron tulipifera* Linn.
563 reveal candidate genes involved in nectary development and nectar secretion. *BMC Plant Biol.*
564 19, 531

565 92 Ruhlmann, J.M. *et al.* (2010) CELL WALL INVERTASE 4 is required for nectar production in
566 Arabidopsis. *J. Exp. Bot.* 61, 395–404

567 93 Chen, L.-Q. *et al.* (2010) Sugar transporters for intercellular exchange and nutrition of pathogens.
568 *Nature* 468, 527–532

569 94 Lin, I.W. *et al.* (2014) Nectar secretion requires sucrose phosphate synthases and the sugar
570 transporter SWEET9. *Nature* 508, 546–549

571 95 Seo, H.S. *et al.* (2001) Jasmonic acid carboxyl methyltransferase: a key enzyme for jasmonate-
572 regulated plant responses. *Proc. Natl. Acad. Sci.* 98, 4788–4793
573 96 Lee, Z.H. *et al.* (2019) CRABS CLAW and SUPERMAN Coordinate Hormone-, Stress-, and
574 Metabolic-Related Gene Expression During Arabidopsis Stamen Development. *Front. Ecol. Evol.*
575 7, 437
576 97 Baylis, T. *et al.* (2013) *SHORT INTERNODES/STYLISH* genes, regulators of auxin biosynthesis, are
577 involved in leaf vein development in *Arabidopsis thaliana*. *New Phytol.* 197, 737–750
578 98 Bowman, J.L. *et al.* (1989) Genes directing flower development in Arabidopsis. *Plant Cell* 1, 37–52
579 99 Bowman, J.L. *et al.* (1991) Genetic interactions among floral homeotic genes of Arabidopsis.
580 *Development* 112, 1–20
581 100 Cao, D. *et al.* (2006) Gibberellin Mobilizes Distinct DELLA-Dependent Transcriptomes to
582 Regulate Seed Germination and Floral Development in Arabidopsis. *PLANT Physiol.* 142, 509–525
583
584

585 **Glossary**

586 **Nectar:** a sugar-rich, phloem derived solution which contains products of primary and
587 secondary metabolism.

588 **Nectary:** Secretory nectar-producing organ formed anywhere on the plant apart from the
589 root.

590 **Whorl:** a concentric ring of floral organs.

591 **Transcription factor (TF):** a molecule that binds to DNA-regulatory sequence to modulate
592 the rate of gene transcription.

593 **CRABS CLAW (CRC):** a putative TF which controls processes such as carpel development, floral
594 meristem termination, and floral nectary formation.

595 **YABBY (YAB):** the gene family named after the crabs claw-like appearance of the apically unfused
596 carpels of the *crc-1* mutant.

597

598 **Figure 1.** Transversely dissected flower of melon (*Cucumis melo*) prior to anthesis using
599 binocular (left) and scanning electron microscopy (SEM, right). Nectary develops as a single
600 dome-shaped structure at the base of the male flower (upper panel), while it completely
601 encircles the stigmatic tissue of the female flower (lower panel). Scale bar represents 2 mm.

602

603 **Figure 2.** Model of genetic control of nectary development. In arabidopsis, *CRC* is activated
604 by a combination of the C- (*AG*) and E- (*SEP*) class gene activities. In the absence of the *AG*
605 gene activity, *SHATTERPROOF1/2* can be sufficient to activate *CRC* (Proposed by Lee et al.
606 2005), while the B functions *AP3/PI* repress *CRC* [50]. *BOP1/2* genes control nectary size [51],
607 and they function in the same genetic pathway with *PAN*. In basal eudicots, the *STY* genes
608 play the key role in nectary formation [60] through the auxin biosynthesis pathway [97].

609 **Figure 3.** Model of nectar secretion at anthesis (S12). Hormones GA, auxin and JA regulate
610 nectar secretion. GAs endogenous to nectaries repress nectar production [80], while GA
611 outside of the nectary may induce JA production in stamen filaments which leads to auxin

612 production in a positive feedback loop [76]. In turn, IAA may trigger ARF expression and lead
613 to the expression of the MYB21/MYB305, which are required for nectary maturation and
614 transcription of the downstream genes in starch metabolism [reviewed in [4]. Upon starch
615 breakdown, the cellular osmotic pressure rises and triggers the influx of water from the
616 phloem. As a result, the increased hydrostatic pressure within nectary causes nectar to
617 exude through nectary pores [87].

618

619 **Table 1.** The effects of floral homeotic mutants on nectary phenotype in *Arabidopsis thaliana*

Gene (Mutant)	Function	Nectary phenotype	Refs.
<i>lfy-6</i>	Master regulator of flower meristem identity	Sometimes present in lateral domain, reduced in size, do not possess stomata	[26]
<i>ufo-2</i>	Regulator of floral meristem identity	Reduced in size, do not possess stomata. More glands present than in <i>lfy-6</i>	[26]
<i>lfy-6 ufo-2</i>		No nectaries	[26]
<i>pi-1</i>	B	Nectaries reduced in size; LN normal, MN not always present	[26]
<i>ag-1</i>	C	Disk shaped; nectaries develop between the 2 nd and the 3 rd , or outside the 3 rd whorl	[98]
<i>ap3</i>	B	Most flowers have both LN and MN but without stomata	[98]
<i>sep 1/2/3</i>	E	No nectaries	[41]
<i>shp1 shp2</i>	C	Normal	[42]
<i>pi-1 ag-1</i>	B C	No nectaries	[26], [98]
<i>ap3-3 ag-3</i>	B C	No nectaries	[98]
<i>ap2-2 pi-1</i>	A B	Nectaries develop interior of the lateral first whorl organs	[99]
<i>ag shp1 shp2</i>	C	No nectaries	[42]
<i>ap2 pi ag shp1 shp2</i>	A B C	No nectaries	[49]

620 LN – lateral nectaries; MN – median nectaries.

621

622

623 **Table 2.** Selected genes with altered expression in nectaries of *arf6-2 arf8-3* mutants relative
 624 to the wild type^a

Gene name	Function	Expression in <i>arf6 arf8</i> compared to Col-0 at stage 12	Dependence on additional hormonal signaling	Locus	Refs.
CRC	transcription factor	Down regulated	unknown	AT1G69180	[73]
YABBY5	transcription factor YABBY family protein	Down regulated	unknown	AT2G26580	[73]
CWIN4	beta-fructosidase, putative expressed protein	Down regulated	unknown	AT2G36190	[73]
SWEET9	nodulin MtN3 family protein	Down regulated	unknown	AT2G39060	[73]
TPS11	terpene synthase/cyclase family protein	Down regulated	DELLA-repressed	AT5G44630	[73], [100]
	cytochrome P450 family protein	Down regulated	DELLA-repressed	AT5G44620	[73], [100]
	copper-binding family protein	Down regulated	DELLA-repressed	AT5G24580	[73], [100]
JMT	S-adenosyl-L-methionine:jasmonic acid carboxyl methyltransferase	Down regulated	DELLA-repressed	AT1G19640	[73], [100]
	3-hydroxyisobutyryl-coenzyme A hydrolase, putative	unchanged	DELLA-repressed	AT2G30650	[73], [100]
	a putative zinc finger protein	Down regulated	GA-repressed	AT1G32540	[73], [100]
	unknown expressed protein	Down regulated	GA-induced	AT3G60780	[73], [100]
SAUR66	auxin-responsive protein	Down regulated	GA-induced		[73], [100]
	strictosidine synthase family protein	Down regulated	GA-induced	AT1G74020	[73]

625 ^aData are based on Affymetrix ATH1 gene chip array [76]

626

Figure 1




Figure 2




Figure 3

