



HAL
open science

Leaf and Branch Hydraulic Plasticity of Two Light-Demanding Broadleaved Tree Species Differing in Water-Use Strategy

Eele Öunapuu-Pikas, Jean-Stéphane Venisse, Arne Sellin, Philippe Label

► **To cite this version:**

Eele Öunapuu-Pikas, Jean-Stéphane Venisse, Arne Sellin, Philippe Label. Leaf and Branch Hydraulic Plasticity of Two Light-Demanding Broadleaved Tree Species Differing in Water-Use Strategy. *Forests*, 2022, 13 (4), pp.594. 10.3390/f13040594 . hal-03636792

HAL Id: hal-03636792

<https://hal.inrae.fr/hal-03636792>

Submitted on 11 Apr 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Article

Leaf and Branch Hydraulic Plasticity of Two Light-Demanding Broadleaved Tree Species Differing in Water-Use Strategy

Eele Õunapuu-Pikas ^{1,*}, Jean-Stéphane Venisse ^{2,3}, Philippe Label ^{2,3} and Arne Sellin ¹¹ Institute of Ecology and Earth Sciences, University of Tartu, 51005 Tartu, Estonia; arne.sellin@ut.ee² Physique Intégrative et Physiologie des Arbres Fruitières et Forestiers, Université Clermont Auvergne, 63000 Clermont-Ferrand, France; j-stephane.venisse@uca.fr (J.-S.V.); philippe.label@inrae.fr (P.L.)³ Integrative Physics and Physiology of Fruit and Forest Trees, INRAE, 63000 Clermont-Ferrand, France

* Correspondence: eele.ounapuu-pikas@ut.ee

Abstract: Global climate change creates new environmental scenarios and selective pressures; thus, a better understanding of the plasticity of plant functional traits is needed to predict how plant species will respond to shifts in climate. Among the important functional traits for plants are their hydraulic properties which ultimately determine their photosynthetic capacity, growth rate, and survival in a changing environment. In this study, the light sensitivity of leaf (K_L) and branch hydraulic conductance (K_B) to fast changes in irradiance, and hydraulic plasticity (PI_h) was studied in two broadleaved tree species differing in water-use strategy—silver birch (*Betula pendula*) and hybrid aspen (*Populus × wettsteinii*). The K_L increased by a factor of 3.5 and 1.5 from minimal values recorded in darkness to maximal values in high light conditions for birch and aspen, respectively, indicating a significantly higher PI_h for birch (0.72) than for aspen leaves (0.35). K_B increased 1.5-fold from dark to light conditions for both species. The high light sensitivity of K_L and K_B provides a regulatory mechanism to maintain a balance between transpirational demand and hydraulic supply. The plasticity of these traits increases the ability of plants to cope with a rapidly changing environment and to adapt to global climate change.

Keywords: acclimation; fluctuating irradiance; hydraulic conductance; ionic effect; light sensitivity; plasticity; plant–water relations; xylem sap

Citation: Õunapuu-Pikas, E.; Venisse, J.-S.; Label, P.; Sellin, A. Leaf and Branch Hydraulic Plasticity of Two Light-Demanding Broadleaved Tree Species Differing in Water-Use Strategy. *Forests* **2022**, *13*, 594. <https://doi.org/10.3390/f13040594>

Academic Editor: Marcello Vitale

Received: 11 March 2022

Accepted: 8 April 2022

Published: 10 April 2022

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2022 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

1. Introduction

Today, plant scientists solidly acknowledge the integral role of water transfer in soil–plant–atmosphere systems (i.e., plant hydraulics) in understanding plant and ecosystem functioning [1,2]. Water is an essential resource that all living organisms need for existence. On a global scale, water availability is an important limiting factor for plants, determining the productivity of crops and forests that humans rely on [3,4]. The structure of the water transport system places a physical limit on plant functioning, as the efficiency of water transport through their bodies determines their leaf water status, photosynthetic capacity, and growth rate [5–7]. Thus, plant hydraulic properties are fundamental determinants of plant fitness and their competitiveness and survival in plant communities. Moreover, hydraulic traits mediate the ways in which plants interact with their abiotic and biotic environments [1].

Leaf hydraulic conductance (K_L) is one of the most important physiological parameters in plant–water relations. It is the measure of water flow efficiency through the leaf (from stem to the sites of evaporation) and is defined as the water flow rate through the leaf per water potential difference across the leaf and generally expressed on a leaf area basis [8]. Leaves comprise more than $1/3$ of the total hydraulic resistance of a plant, thus representing a major bottleneck of the water transport pathway in plants [8–10]. Any

changes in leaf hydraulic efficiency will have a significant impact on the water balance of the whole plant.

Plants, as sessile organisms inhabiting temporally and spatially variable environments, present a sizeable capacity to acclimate to their growth environment to survive, grow, compete, and reproduce successfully [11]. Phenotypic plasticity is considered one of the major means by which plants can cope with heterogeneous and variable conditions [12]. By influencing the organisms' fitness and competitive ability, phenotypic plasticity plays a role in the survival of individuals and persistence of populations under changing climate conditions [13–15]. Phenotypic plasticity is defined as the capacity of a genotype to produce different phenotypes in response to environmental conditions [13,16]. A vast number of studies show that plants are plastic in numerous functionally important traits, including morphology, physiology, and anatomy [16,17], and plastic adjustments occur in response to changes in various environmental factors, such as water availability, light gradients, temperature, atmospheric CO₂ concentration, pollution, or combinations of global change drivers [18]. In terms of future perspectives, it is important to identify plant functional traits, the plasticity of which may play a determinant role in plant acclimation to global climate change [12].

Global climate change creates considerable geographical variability in climate, in which trends are different on a regional scale. For instance, although an increasing incidence of climate extremes is predicted for the whole northern hemisphere, future climate scenarios predict wetter conditions at high latitudes, including increasing precipitation in northern Europe, while southern Europe will experience substantial warming and drying [19]. Climate change scenarios for the year 2100 predict an increase in air temperature (by 2.3–4.5 °C) and precipitation (by 5–30%) in the Baltic region [20]. The rising amount and frequency of rainfall will increase soil water content, as well as air relative humidity (RH), on a local scale [21–23], especially within forest canopies [24]. Changes in precipitation and weather patterns will have significant consequences on the growth of trees and the functioning of forest ecosystems [25–27]. Results from the Free Air Humidity Manipulation (FAHM) experiment show that under increasing atmospheric humidity, trees experience less water loss and there is weaker environmental pressure to invest resources into developing effective water conducting tissues, resulting in reduced hydraulic conductance in leaves [28,29]. In the case of climate extremes (heat waves, severe droughts), which have been predicted to become more frequent during the 21st century across Europe, reduced hydraulic efficiency might represent a potential threat in hemi-boreal forest ecosystems [30]. Understanding which hydraulic traits are phenotypically plastic and to what extent is essential for understanding how tree species will respond to shifts in climate [31].

K_L is a highly plastic trait, showing great variability and dynamics both between and within species, on different timescales, and in response to various environmental factors [8]. Light is one of the most important environmental factors besides water availability and temperature that influences leaf hydraulic conductance. K_L has been shown to exhibit sensitivity to irradiance, e.g., [32–35], but the magnitude of the light response (i.e., plasticity) varies between species [36,37]. To our knowledge, only a few studies have specifically quantified the hydraulic plasticity of different species. Scoffoni et al. [38] studied the plasticity of K_L in differential growth light environments in endemic Hawaiian lobeliads and found species native to higher irradiance to show greater hydraulic plasticity. In one of our previous studies [39], we quantified the intracanalopy plasticity of K_L to incident light in common hazel (*Corylus avellana* L.) and found sun-exposed leaves to be considerably more plastic than leaves growing in deep shade. In the context of global climate change, a better understanding of the plasticity of plant functional traits is necessary to predict the adaptations of forest ecosystems to future conditions and to elaborate appropriate management practices. As plant hydraulics represents a central hub integrating plant and ecosystem function, greater hydraulic plasticity (i.e., a fast and responsive adjustment of hydraulic conductance depending on environmental factors) is likely to allow trees to adapt

more easily to climate change. Thus, it is important to investigate the variation of the plasticity of plant hydraulic traits responsible for adequate leaf water supply in more detail.

In this study, light sensitivity and plasticity of K_L was measured in silver birch (*Betula pendula* Roth) and hybrid aspen (*Populus × wettsteinii* Hämet-Ahti). Both species are considered typical light-demanding fast-growing species that are able to quickly colonise disturbed sites [40]. Silver birch is commercially the most important fast-growing early-successional tree species widely distributed in northern Europe [41]. Hybrid aspen is the fastest-growing deciduous tree species in the boreal zone and is commercially a valuable and promising species for short-rotation plantation forestry in northern Europe [42]. These two species were chosen for this study primarily because they grow at the study site. From an ecological perspective, these species were suitable for this study because they are similar in their geographical distribution and commercial importance, but at the same time they differ in their water-use strategy and thus might react to and cope with climate change differently. Silver birch represents an isohydric species with a conservative water-use strategy [43–45], whereas aspen inclines to anisohydric behaviour with a prodigal water-use strategy [40,45–47]. The objective of this study was to test whether and to what extent the plasticity of K_L to incident light differs between tree species with contrasting water-use strategies. Evidence is also presented on the light-mediated short-term modulation of branch hydraulic conductance. The results of this research will improve forest management in the future by broadening the understanding of the ability of trees to acclimate in changing environmental conditions and by providing additional information to maintain ecologically stable, diverse, and productive ecosystems. Hydraulically plastic tree species are more likely to adapt to a rapidly fluctuating environment. Hence, through forest selection and the identification of plastic genotypes, it is possible to shape the development of forest ecosystems that would be more stable against climate change.

2. Materials and Methods

2.1. Study Site and Sample Trees

This study was performed in the summer of 2019 at the Free Air Humidity Manipulation (FAHM) experimental site situated at Rõka village (58°14' N, 27°17' E, 40–48 m ASL) in eastern Estonia. The site belongs to the hemi-boreal forest zone and to the maritime to continental transitional climate zone. The mean annual precipitation in this area is 650 mm, with a mean air temperature of 17.0 °C in July and −6.7 °C in January. The growing period lasts 175–180 days, from mid-April to October. The total annual global short-wave radiation in the region averages 3518 MJ m⁻², and the annual radiation budget is 2552 MJ m⁻² [48]. The soil type at the site is fertile Endogleyic Planosol (WRB).

Based on climate change scenarios that predict increasing precipitation and concomitant rise in air relative humidity for northern Europe [19,49,50], the FAHM facility was established to investigate the effects of increasing air humidity on tree performance and the functioning of a deciduous forest ecosystem [51]. The experimental design and technical setup are described in detail by Kupper et al. [51], Tullus et al. [52], and Sellin et al. [53].

This study was performed on excised shoots of seven-year-old trees of silver birch (*Betula pendula* Roth clonal plants micropropagated from meristem cultures that originated from southern Finland, Vehmersalmi (62°45' N, 28°10' E); [54]) and six-year-old trees of hybrid aspen (*Populus × wettsteinii* Hämet-Ahti; root and stump regenerated offspring of clone C05-99-34 micropropagated from meristem culture) growing at the FAHM site. For hydraulic measurements, 252 shoots (14 shoots × 9 repetitions × 2 species) in total were sampled.

In the evening, prior to the measurement day, 14 birch (average length 28.0 cm) or aspen (20.5 cm) shoots were cut, recut under water to avoid airseeding of the shoot xylem, and inserted into test tubes with their basal ends submerged in water. Shoots were then

transported to the laboratory and put into plastic flasks filled with deionised, filtered (Direct-Q3 UV water purification system; Millipore SAS, Molsheim, France), and freshly degassed water (T-04-125 ultrasonic-vacuum degasser; Terriss Consolidated Industries, Asbury Park, NJ, USA). Shoots were then left to rehydrate overnight in a dark room until the hydraulic measurements were performed the next day.

2.2. Measuring Hydraulic Conductance

Hydraulic measurements were carried out by applying the water perfusion method using a high-pressure flow meter (HPFM; Dynamax, Houston, TX, USA) applied in quasi-steady state mode [32]. In the morning, the first two shoots were brought sequentially from the dark room and immediately measured for hydraulic conductance under dark conditions (light treatment dark-dark, DD). The shoots were recut under water and immediately attached to the HPFM apparatus. Water was perfused into the shoot until hydraulic conductance values became constant. Then, the leaves were removed with a razor blade, and the perfusion continued until again values became constant. Next, another two shoots were brought sequentially from the dark room and sampled for hydraulic conductance under dark conditions. Subsequently, the light was switched on, and the same shoots were sampled under light (PPFD 900–1000 $\mu\text{mol m}^{-2} \text{s}^{-1}$) conditions (light treatment dark-light, DL). Then, the light was switched off again, until the readings in the darkness became constant again (light treatment light-dark, LD). The leaf blades were then removed, and new constant values were again achieved. The remaining 10 shoots to be sampled that day were brought into the light at the same time (light treatment light-light, LL). Two shoots were exposed to light for 1, 2, 3, 4, or 5 h (1 hL, 2 hL, 3 hL, 4 hL, 5 hL). Each shoot was sampled first under light conditions; then, the light was turned off, and the shoot was sampled under dark conditions. Finally, the leaves were cut off and the leafless branch was sampled. The irradiance level chosen for our experiment was in accordance with Xiong et al. [37], who showed that the light intensity level required to saturate leaf hydraulic conductance (K_L ; $\text{kg m}^{-2} \text{s}^{-1} \text{MPa}^{-1}$) is around 1000 $\mu\text{mol m}^{-2} \text{s}^{-1}$ for most species. LED Fyto-Panels (Photon Systems Instruments, Drásov, Czech Republic) consisting of neutral white light sources supplemented with far-red LEDs (735 nm) were used for illumination prior to and in the course of the measurements. To minimise small-scale temperature gradients, the air in the vicinity of the shoots was stirred with a ventilator. During the measurements, leaf and branch temperature were recorded with an MT2 fast response temperature sensor (Delta-T Devices, Burwell, UK).

After hydraulic measurements, the shoot length was measured with a ruler. Additionally, shoot diameter was measured under bark from two perpendicular directions with a digital calliper for calculations of xylem cross-sectional area. The total leaf blade area of the shoot (A_L ; m^2) was determined with a LI-3100C optical area meter (LI-COR Biosciences, Lincoln, NE, USA). Leaves were then dried in an oven at 80 °C for 48 h, and the total leaf dry weight of the shoot (DW; g) was recorded. Leaf mass per area (LMA; g m^{-2}) was calculated for each shoot. The Huber value (HV) of each shoot was calculated by dividing the xylem cross-sectional area by A_L . The hydraulic conductance values were scaled by A_L and corrected for the dynamic viscosity of water at 23 °C. K_L was calculated as follows:

$$K_L = \frac{1}{K_S^{-1} - K_B^{-1}} \quad (1)$$

where K_S is the total shoot hydraulic conductance, and K_B ($\text{kg m}^{-2} \text{s}^{-1} \text{MPa}^{-1}$) is the hydraulic conductance of the leafless branch. K_L in the LL treatment represents the maximum leaf hydraulic conductance ($K_{L\text{max}}$), and K_L in the DD treatment represents the minimum leaf hydraulic conductance ($K_{L\text{min}}$).

The up-regulation time (t_{up} ; s) and down-regulation time (t_{down} ; s) of K_L were determined as the time elapsed between turning on or off the light until hydraulic conductance

values became constant, respectively. The up-regulation rate of K_L (dK_{Lup}/dt ; $\text{kg m}^{-2} \text{s}^{-2} \text{MPa}^{-1}$) was calculated as follows:

$$\frac{dK_{Lup}}{dt} = \frac{K_{Lmax} - K_{Lmin}}{t_{up}} \quad (2)$$

The down-regulation rate of K_L (dK_{Ldown}/dt ; $\text{kg m}^{-2} \text{s}^{-2} \text{MPa}^{-1}$) was calculated as follows:

$$\frac{dK_{Ldown}}{dt} = \frac{K_{Lmax} - K_{L,LD}}{t_{down}} \quad (3)$$

where $K_{L,LD}$ is leaf hydraulic conductance in the LD treatment. The initial up-regulation rate of K_L (b_{up} ; $\text{kg m}^{-2} \text{s}^{-2} \text{MPa}^{-1}$) was calculated as the slope of the linear part of K_L vs. time regression. Relative hydraulic resistance residing in leaves (R_L ; %) was calculated as follows:

$$R_L = \frac{K_L^{-1} \times 100}{K_S^{-1}} \quad (4)$$

2.3. Plasticity Index

The plasticity index [55] was calculated for both species and used to characterise hydraulic plasticity (PI_h):

$$PI_h = \frac{K_{max} - K_{min}}{K_{max}} \quad (5)$$

where K_{max} and K_{min} represent the species mean maximum and minimum hydraulic conductance, respectively, of leaves (K_L) or leafless branches (K_B).

2.4. Data Analysis

Statistical data analysis was performed using Statistica, Version 7.1 (StatSoft Inc., Tulsa, OK, USA). The normality of the data and the homogeneity of variance were checked using the Kolmogorov–Smirnov D -statistic and Levene’s test, respectively. Light exposition time (1–5 h) did not influence K_L ; thus, K_L data from light treatments 1 hL to 5 hL were pooled together and treated as a unitary light treatment (LL). To analyse the effects of categorical factors (species, light treatment) on leaf and branch hydraulic traits, an analysis of variance (ANOVA) was performed using the ‘general linear model’ module. Type IV sums of squares were used in the calculations. To fulfil the assumptions for ANOVA, logarithmic or complex transformations were applied to the data when necessary. Post hoc mean comparisons were conducted using Tukey’s HSD test. Bivariate relationships between the studied characteristics were assessed by Pearson’s correlations and simple linear or nonlinear least-squares regressions.

3. Results

Morphological traits of the sampled shoots are given in Table 1. A_L ($p < 0.001$), LMA ($p < 0.001$), and HV ($p = 0.024$) were significantly higher in aspen than in birch.

Table 1. Mean values (\pm SE) of shoot morphological traits in silver birch (*Betula pendula*) and hybrid aspen (*Populus × wettsteinii*). L_s —shoot length, A_L —total leaf blade area of the shoot, LMA—leaf mass per area, HV—Huber value.

Trait	<i>B. pendula</i>	<i>P. × wettsteinii</i>	Statistical Significance
L_s (cm)	27.9 \pm 0.34	20.5 \pm 0.38	$p < 0.05$
A_L (cm ²)	167 \pm 3.4	338 \pm 9.1	$p < 0.05$
LMA (g m ⁻²)	60.8 \pm 0.64	77.8 \pm 0.74	$p < 0.05$

HV ($\times 10^{-4}$; $\text{m}^2 \text{m}^{-2}$) 1.24 ± 0.023 1.34 ± 0.035 $p < 0.05$

Light treatment had a significant effect ($p \leq 0.001$) on both leaf and branch hydraulic conductance (Table 2). For both species, K_L was lowest in the initial dark conditions (treatment DD, $K_{L\text{min}}$), intermediate a few minutes after the light was turned on (treatment DL), highest in light lasting over an hour (treatment LL, $K_{L\text{max}}$), and it declined again in the second dark period (treatment LD) (Figure 1). The mean K_L for birch was $0.80 \pm 0.05 \times 10^{-4}$, $1.94 \pm 0.14 \times 10^{-4}$, $2.82 \pm 0.07 \times 10^{-4}$, and $2.26 \pm 0.04 \times 10^{-4} \text{ kg m}^{-2} \text{ s}^{-1} \text{ MPa}^{-1}$ for DD, DL, LL, and LD treatments, respectively. For aspen, K_L averaged $1.90 \pm 0.07 \times 10^{-4}$, $2.23 \pm 0.08 \times 10^{-4}$, $2.92 \pm 0.05 \times 10^{-4}$, and $2.84 \pm 0.06 \times 10^{-4} \text{ kg m}^{-2} \text{ s}^{-1} \text{ MPa}^{-1}$ in DD, DL, LL, and LD treatments, respectively.

Table 2. Results of ANOVA for the effects of species and light treatment on leaf and branch hydraulic properties.

Trait	Factor	Statistical Significance
Leaf hydraulic conductance, K_L ($\text{kg m}^{-2} \text{ s}^{-1} \text{ MPa}^{-1}$)	Species	$p < 0.001$
	Light treatment	$p < 0.001$
	Species \times Light tr.	$p < 0.001$
Branch hydraulic conductance, K_B ($\text{kg m}^{-2} \text{ s}^{-1} \text{ MPa}^{-1}$)	Species	$p < 0.001$
	Light treatment	$p = 0.001$
	Species \times Light tr.	$p = 0.174$
Up-regulation time of K_L , t_{up} (s)	Species	$p < 0.001$
Down-regulation time of K_L , t_{down} (s)	Species	$p < 0.001$
Up-regulation rate of K_L , ($dK_{L\text{up}}/dt$; $\text{kg m}^{-2} \text{ s}^{-2} \text{ MPa}^{-1}$)	Species	$p = 0.667$
Down-regulation rate of K_L , ($dK_{L\text{down}}/dt$; $\text{kg m}^{-2} \text{ s}^{-2} \text{ MPa}^{-1}$)	Species	$p < 0.001$
Initial up-regulation rate of K_L , (b_{up} ; $\text{kg m}^{-2} \text{ s}^{-2} \text{ MPa}^{-1}$)	Species	$p = 0.193$




Figure 1. Mean values (\pm SE) of leaf hydraulic conductance (K_L) depending on species and light treatment. DD—initial dark conditions; DL—initial light conditions (exposure to light for less than 1 h); LL—light conditions (exposure to PPFD 900–1000 $\mu\text{mol m}^{-2} \text{ s}^{-1}$ for 1–5 h); LD—final dark conditions

(light turned off after exposure to light for 1–5 h). Different letters express statistically significant ($p < 0.001$) differences between light treatments within a species; asterisks denote statistically significant ($p < 0.01$) differences between species within a light treatment.

K_L varied significantly ($p < 0.001$) between the species (Table 2); the mean K_{Lmin} in aspen was 2.4 times higher ($p < 0.001$) than that in birch (Figure 1). K_L in the DL treatment tended to be higher in aspen, but the difference was statistically not significant ($p = 0.08$). The means of K_{Lmax} did not differ ($p = 0.23$) between species. In the LD treatment, K_L for aspen was on average 1.26 times higher ($p < 0.01$; Figure 1) compared to birch leaves.

The interaction between species and light treatment was significant ($p < 0.001$); the species differed in the amplitude of K_L (i.e., the difference between K_{Lmax} and K_{Lmin}). K_L increased 3.5 ($p < 0.01$) and 1.5 times ($p < 0.01$) from dark to light conditions for birch and aspen, respectively (Figure 1). Thus, birch leaves exhibited a greater hydraulic plasticity than aspen; the PI_h for birch leaves (PI_{hL}) was 0.72 compared to 0.35 for aspen. The species also differed significantly in the decrease of leaf hydraulic conductance when transferring shoots from light to dark. Birch exhibited a 25% decrease ($p < 0.01$) and aspen a 3% decrease ($p = 0.263$) in K_L after the light was switched off (Figure 1). K_{Lmax} was significantly and positively correlated to HV for both birch ($r = 0.27$, $p = 0.017$) and aspen ($r = 0.26$, $p = 0.014$).

Branch hydraulic conductance also responded to the application of light and differed between the species (Table 2). K_B was lowest in the dark (DD and DL treatments) and reached a maximum after 3 h of light exposure (3 hL) in birch (Figure 2). In aspen, K_B was lowest under DD treatment and reached a maximum in light lasting for one hour (1 hL; Figure 2). For birch, the minimal and maximal K_B averaged $4.03 \pm 0.29 \times 10^{-4}$ and $6.06 \pm 0.54 \times 10^{-4} \text{ kg m}^{-2} \text{ s}^{-1} \text{ MPa}^{-1}$, respectively. For aspen, the minimal and maximal values of K_B were $7.54 \pm 0.60 \times 10^{-4}$ and $11.47 \pm 0.83 \times 10^{-4} \text{ kg m}^{-2} \text{ s}^{-1} \text{ MPa}^{-1}$, respectively. K_B increased 1.5 times ($p < 0.01$) from dark to light conditions for both species. Both minimal and maximal K_B were 1.9-fold ($p < 0.01$) higher for aspen compared to birch. The species did not differ in the hydraulic plasticity of branches; the PI_h for birch branches (PI_{hB}) was 0.33 and for aspen branches 0.34. K_B in light conditions was significantly and positively correlated with HV for both birch ($r = 0.54$, $p < 0.001$) and aspen ($r = 0.33$, $p = 0.001$). In light, K_B and K_L were positively correlated in birch ($r = 0.36$, $p = 0.001$) and aspen ($r = 0.40$, $p < 0.001$), but they were not correlated in the dark.




Figure 2. Mean values (\pm SE) of branch hydraulic conductance (K_B) depending on species and light treatment. DD—initial dark conditions; DL—initial light conditions (exposure to light for less than 1 h); 1 hL–5 hL—light conditions (exposure to PPFD 900–1000 $\mu\text{mol m}^{-2} \text{s}^{-1}$ for 1–5 h). Different letters express statistically significant ($p < 0.001$) differences between the light treatments within a species.

The times of up- and down-regulation of K_L (t_{up} and t_{down}) in response to turning the light on and off differed significantly between the species (Table 2). The mean t_{up} was 483 ± 32 s for birch and 150 ± 12 s for aspen ($p < 0.001$), and the mean t_{down} was 210 ± 5 s for birch and 251 ± 4 s for aspen ($p < 0.001$; Figure 3). t_{up} and t_{down} differed significantly from each other. In birch, t_{up} was 2.3 times longer ($p < 0.01$) than t_{down} , while the opposite trend was observed in aspen, in which t_{down} was 1.7 times longer ($p < 0.01$) than t_{up} (Figure 3). Across species, t_{up} was strongly and inversely related to LMA ($r^2 = 0.60$, $p < 0.001$; Figure 4).




Figure 3. Mean values (\pm SE) of the up- (t_{up}) and down-regulation (t_{down}) times of leaf hydraulic conductance of silver birch (*Betula pendula*) and hybrid aspen (*Populus × wettsteinii*). Different letters express statistically significant ($p < 0.01$) differences between the means.




Figure 4. Relationship between leaf mass per area (LMA) and time of K_L up-regulation (t_{up}) across species. All data points correspond to one shoot.

The up-regulation rate of K_L (dK_{Lup}/dt) did not differ between the species; the corresponding mean values were $2.94 \pm 0.44 \times 10^{-7}$ and $2.72 \pm 0.22 \times 10^{-7} \text{ kg m}^{-2} \text{ s}^{-2} \text{ MPa}^{-1}$ for birch and aspen, respectively ($p = 0.667$; Figure 5). Additionally, the initial up-regulation rate of K_L (b_{up}) did not differ significantly between the species ($p = 0.193$). However, the down-regulation rate of K_L (dK_{Ldown}/dt) differed substantially between species, and the corresponding means were $3.50 \pm 0.47 \times 10^{-7}$ and $0.83 \pm 0.08 \times 10^{-7} \text{ kg m}^{-2} \text{ s}^{-2} \text{ MPa}^{-1}$ for birch and aspen, respectively ($p < 0.01$), exhibiting 4.2 times faster down-regulation in birch leaves (Figure 5). For birch, there was no significant difference between the up- and down-regulation rates of K_L , whereas for aspen, dK_{Lup}/dt was 3.3-fold faster than dK_{Ldown}/dt ($p < 0.01$).




Figure 5. Mean values (\pm SE) of the rates of up- (dK_{Lup}/dt) and down-regulation of K_L (dK_{Ldown}/dt) of silver birch (*Betula pendula*) and hybrid aspen (*Populus × wettsteinii*). Different letters express statistically significant ($p < 0.01$) differences between the means.

The changes in K_L induced by exposure to light brought about a considerable redistribution of the hydraulic resistance within the sample shoots. The relative contribution of leaves to the whole-shoot hydraulic resistance (R_L) under dark conditions was similar for the two study species, averaging 83 and 80% for birch and aspen, respectively (Figure 6). After at least 1 h of illumination, R_L decreased to 65 and 77% for birch and aspen, respectively, indicating a significantly greater decline in birch.




Figure 6. Relative hydraulic resistance (mean \pm SE) residing in leaves (R_L) of silver birch (*Betula pendula*) and hybrid aspen (*Populus × wettsteinii*) under dark or light conditions (exposure to PPFD 900–1000 $\mu\text{mol m}^{-2} \text{s}^{-1}$ for at least 1 h). Different letters express statistically significant ($p < 0.01$) differences between the means.

4. Discussion

4.1. Light Effects on Leaf Hydraulic Properties

The results of this study confirm the high light sensitivity of leaf hydraulic conductance in silver birch and hybrid aspen; a rapid and substantial modulation of K_L took place in response to switching light on and off in both species (Figures 1 and 3). Nearly 20 years have passed since the earliest studies on the light sensitivity of K_L [32,56,57], and the enhancement of K_L by light has been generally acknowledged and amply confirmed in various species [35–37,39,58]. The light-induced rapid enhancement of K_L has been shown to originate from the outside-xylem compartment of the leaf [34,57] being specifically related to the up-regulation of aquaporins (AQPs) through enhanced gene expression and/or post-translational protein activation in response to irradiance. In *B. pendula*, a genome analysis has identified 33 putative genes encoding full-length AQP sequences [59]. The light-dependent increase of K_L shows a tight correlation with AQP gene transcript abundance [60–64]. The activation of AQPs by phosphorylation is also associated with enhanced leaf water transport capacity [65,66]. At the present time, it is not yet clear which mechanism is primarily responsible for light-induced rapid K_L enhancement, whether it is AQP expression or activation.

In the current study, K_L was lowest in the initial dark conditions, intermediate after the light was turned on, and highest in light lasting for an hour or more (Figure 1). Variation of the exposure time from 1 to 5 h did not influence leaf hydraulic capacity. In our

earlier study [33], also carried out in silver birch, K_L increased up to 9 h of illumination. This discrepancy is attributable to different experimental conditions; in that study, substantially lower irradiance was applied, which is why more time was needed for AQP up-regulation. We propose that the rapid changes in K_L that occurred directly after turning the light on/off probably took place through the post-translational activation of AQPs and/or their trafficking to the plasma membrane, whereas the maximal values of K_L were achieved through de novo AQP synthesis. In such a context, the PIP (plasma membrane intrinsic protein) and TIP (tonoplast intrinsic proteins) subfamilies would be appropriate candidates, as their functions in water and CO₂ homeostasis are very specialised [67]. Our present results concur with studies indicating that maximal K_L values and AQP transcript amounts occur at least 1 h after switching on the light [60–62]. Xiong et al. [37] indicated that for most species the sufficient light intensity required to saturate K_L is around 1000 $\mu\text{mol m}^{-2} \text{s}^{-1}$. Our results suggest that, for correct estimations of light-saturated K_L , it is also important to take into account the time of light exposure and that 1 h of illumination is enough to reach maximum K_L when really saturating light is applied. The light-induced rapid enhancement of K_L is a means of maintaining stable gas exchange in the event of rapid increases in VPD and evaporative water loss, while avoiding water potential drop and xylem embolism [35,60,68].

The magnitude of light sensitivity of K_L is highly species-specific, ranging from zero to a several-fold increase in K_L in response to incident light [32,34,37,60]. Xiong et al. [37] reported the lack of light sensitivity of K_L of ferns, gymnosperms, and one angiosperm, whereas the other angiosperms under study responded to light. In the current study, the species differed substantially in the magnitude of their response; K_L increased 3.5 and 1.5 times from dark to saturating light conditions for birch and aspen, respectively, reflecting higher hydraulic plasticity in birch compared to aspen. The differences in sensitivity of K_L to light between the species sampled in the current study probably ensued from the extravascular compartment of the leaf, specifically from species specificity in AQP gene expression and/or activation.

Additionally, the differing distribution of resistance between the vascular and extravascular compartments within leaves of different species and changes in preferential water transport pathways with varying light conditions probably contribute to the differences in K_L light sensitivity between the species. Cochard et al. [60] argued that, in the dark, when AQP accumulation/activation is low, hydraulic resistance of the extravascular (R_{ext}) compared to the vascular (R_{vasc}) compartment in the leaf is high and water flows preferentially through the vascular and apoplasmic compartment. At the onset of light, AQPs start to accumulate, thus reducing R_{ext} , and the preferential water pathway changes to the cell-to-cell route. The current consensus is that within the leaves the relative contribution of R_{vasc} and R_{ext} to total leaf hydraulic resistance is relatively equal [8], but depending on species specificity and environmental factors, either one can prevail. Trifilo et al. [69] reported that R_{ext} for *Populus nigra* contributes about 80% to the total leaf hydraulic resistance under low light, while at a high irradiance R_{ext} and R_{vasc} are equal. Considering this information, we can assume that in the current study, the higher hydraulic plasticity resulted, in part, from the higher proportion of R_{ext} under dark conditions in birch leaves compared to aspen.

This study revealed a strong inverse relationship between LMA and t_{up} across the species (Figure 4). No doubt, this is attributable to species specificity, but leaf general morphology probably also plays a role in this relationship: a lower LMA is accompanied by lower leaf vein density (vein length per area, VLA) in birch leaves compared to aspen [29,70]. The lower the VLA, the greater the contribution of the extravascular pathway and the smaller the role of vascular pathway in water movement in the leaf. Therefore, for birch leaves with lower LMA (and VLA) and thus with a higher proportion of R_{ext} , the time for K_L up-regulation tended to be longer. To unravel the basis of K_L light sensitivity more deeply, future studies on K_L light responses should involve the estimations of R_{ext}

and R_{vase} contributions to total leaf hydraulic resistance with concomitant AQP expression analyses.

High hydraulic plasticity is important from many aspects of tree biology, contributing to whole-tree performance through effects on energy, carbon, and water balance of individual leaves in changing or spatially heterogeneous environments. A high PI_h of *B. pendula* is associated with its isohydric behaviour; in addition to the strict stomatal control over water losses [71,72], an efficient and highly dynamic water conducting system is also necessary to maintain the leaf water status within definite limits. Silver birches coordinate their hydraulic capacity with changes in stomatal conductance depending on environmental conditions to prevent leaf water potential from reaching critical values, supporting the isohydric water-use strategy. Moreover, this coordination can also be seen from the leaf anatomical structure. Leaf vascular (size of vascular bundles and vessels, vessel number and density, vein density, etc.) and stomatal traits (stomatal size, density, pore area index) exhibit coordinated adjustment, reflecting the balance between transpirational demand and hydraulic supply in silver birch [29]. On the one hand, exposure of birch foliage to high irradiance (changing patterns of clouds and sun flecks within forest canopies) affords, through a fast increase in K_L and reduction of leaf resistance to water movement (Figures 1 and 6), an adequate hydraulic supply to keep stomata open, and supports evaporative cooling. On the other hand, the capability for efficient down-regulation of K_L helps to prevent excessive water loss in birches in the dark (i.e., arrival of the night) or in the case of shade (Figures 1 and 3).

In hybrid aspen, low PI_h and $dK_{L\text{down}}/dt$ and high K_L in the dark (Figures 3 and 5) afford lavish water use at night, as evidenced by high nocturnal stomatal conductance and low predawn leaf water potential [45,73]. However, high K_L cannot compensate for high water loss due to weak stomatal control day and night, which causes substantial native embolism (percentage loss of hydraulic conductivity 16–41%) in aspen coppice during drought periods [74]. At the leaf level, no solid evidence of coordination was found between vascular and stomatal traits and hydraulic functions, which were most likely associated with the anisohydric behaviour of this species [70].

The up-regulation time of K_L was significantly longer for birch compared to aspen; however, with a much bigger amplitude of K_L between dark and light values, the resulting up-regulation rate of K_L did not differ between the species (Figure 5). We measured similar values of $dK_{L\text{up}}/dt$ also in *Corylus avellana*, a shade-tolerant species [39]. The similar up-regulation rates of K_L among species may imply that the mechanism behind the fast light-induced enhancement of K_L is universal for vascular plants. However, birch exhibited a down-regulation rate over four times faster than aspen, and $dK_{L\text{up}}/dt$ and $dK_{L\text{down}}/dt$ did not differ for birch, whereas for aspen, $dK_{L\text{up}}/dt$ was 3.3 times faster than $dK_{L\text{down}}/dt$ (Figure 5). At present, an unequivocal explanation cannot be provided for why the down-regulation rate for aspen was so much slower than for birch. It may be related to aquaporin metabolism, principally with phosphorylation and dephosphorylation of AQP monomers that form the water channels in cell membranes [75,76]. The exact process of AQP activation and deactivation is subject to research even today [77].

4.2. Light Effects on Branch Hydraulic Properties

The results of this study confirm the light sensitivity and light-induced rapid (time scale in hours) enhancement of branch hydraulic conductance. K_B increased by approximately 50% from dark to light conditions for both species (Figure 2). Water movement in xylem was previously considered a passive process [78,79], determined only by the driving forces and anatomical features of the conducting system (i.e., density, length, diameter of the xylem conduits, structure of interconduit pits) and modified on a developmental time scale. The changes in xylem hydraulic efficiency on a short time scale were attributable to embolism. Presently, there is consensus that xylem hydraulic efficiency can be modulated by the plant on a short time scale also through the modification of xylem sap ionic concentration [80–82]. This ionic effect has been considered to be responsible for the

rapid enhancement of K_B in response to increases in incident light [43,83]. The ion-mediated modulation of xylem hydraulic efficiency is related to the interaction of cations (mainly K^+) with the pectin hydrogels of pit membranes that separate adjacent xylem vessels, and which alters the hydraulic resistance of the bordered pits through changes in the thickness of the pit membrane [84]. The ecophysiological implication of the phenomenon is probably to provide a regulatory mechanism of branch hydraulics to facilitate water flow towards fully illuminated foliage by reducing hydraulic resistance and water potential drop, maximising stomatal aperture and photosynthesis [43,83]. This is a crucial point for fast-growing light-demanding species, such as silver birch and hybrid aspen, contributing to their competitive advantage. A coordinated adjustment of different segments of the water transport pathway is also an essential point in this context. In both species, K_B and K_L were positively correlated in the light but not in the dark, i.e., at night when stomata are closed and atmospheric evaporative demand is low.

5. Conclusions

B. pendula and *P. × wettsteinii*, both fast-growing and light-demanding pioneer species, exhibit high light sensitivity in leaf and branch hydraulic conductance that provide a regulatory mechanism to maintain a balance between transpirational demand and hydraulic supply. However, the tree species differ in their hydraulic plasticity associated with their different water-use strategies; *B. pendula* with its isohydric behaviour exhibits a significantly higher hydraulic plasticity, allowing for a strict control over the water balance of the plant in changing or spatially heterogeneous environments, compared to *P. × wettsteinii* inclined to anisohydric behaviour. In light of climate change one can suppose that hybrid aspen as a species with weak stomatal control and low hydraulic plasticity will be potentially more susceptible to hydraulic impairment compared to silver birch, which will more likely suffer from carbon starvation in case of weather extremes. From a future perspective, it is necessary not merely to determine the plasticity in regard to light conditions but also to investigate possible shifts in the plasticity of plant functional traits (incl. leaf hydraulic conductance) in response to changes in other environmental factors, because most likely they play a determinant role in plant acclimation to climate change.

Author Contributions: Conceptualization, E.Õ.-P. and A.S.; Methodology, E.Õ.-P. and A.S.; Validation, A.S.; Formal Analysis, E.Õ.-P.; Investigation, E.Õ.-P. and A.S.; Resources, A.S.; Writing—Original Draft Preparation, E.Õ.-P.; Writing—Review and Editing, E.Õ.-P., J.-S.V., P.L., and A.S.; Supervision, A.S.; Project Administration, A.S.; Funding Acquisition, A.S. All authors have read and agreed to the published version of the manuscript.

Funding: This study was supported by a grant from the Estonian Research Council (personal research grant PRG1434).

Data Availability Statement: The data that support this study will be shared upon reasonable request to the corresponding author.

Conflicts of Interest: The authors declare no conflicts of interest.

Reference

1. Sack, L.; Ball, M.C.; Brodersen, C.; Davis, S.D.; Des Marais, D.L.; Donovan, L.A.; Givnish, T.J.; Hacke, U.G.; Huxman, T.; Jansen, S.; et al. Plant hydraulics as a central hub integrating plant and ecosystem function: Meeting report for ‘Emerging Frontiers in Plant Hydraulics’ (Washington, DC, May 2015). *Plant Cell Environ.* **2016**, *39*, 2085–2094.
2. Liu, Y.L.; Kumar, M.; Katul, G.G.; Feng, X.; Konings, A.G. Plant hydraulics accentuates the effect of atmospheric moisture stress on transpiration. *Nat. Clim. Chang.* **2020**, *10*, 691–695.
3. Lambers, H.; Chapin, F.S.; Pons, T.L. *Plant Physiological Ecology*, 2nd ed.; Springer: New York, NY, USA, 2008.
4. Taiz, L.; Zeiger, E. *Plant Physiology*, 5th ed.; Sinauer Associates Inc.: Sunderland, MA, USA, 2010.
5. Brodribb, T.J. Xylem hydraulic physiology: The functional backbone of terrestrial plant productivity. *Plant Sci.* **2009**, *177*, 245–251.
6. Scoffoni, C.; Chatelet, D.S.; Pasquet-kok, J.; Rawls, M.; Donoghue, M.J.; Edwards, E.J.; Sack, L. Hydraulic basis for the evolution of photosynthetic productivity. *Nat. Plants* **2016**, *2*, 16072. <https://doi.org/10.1038/NPLANTS.2016.72>.

7. Xiong, D.; Nadal, M. Linking water relations and hydraulics with photosynthesis. *Plant J.* **2020**, *101*, 800–815.
8. Sack, L.; Holbrook, N.M. Leaf hydraulics. *Annu. Rev. Plant Biol.* **2006**, *57*, 361–381.
9. Sack, L.; Cowan, P.D.; Jaikumar, N.; Holbrook, N.M. The ‘hydrology’ of leaves: Co-ordination of structure and function in temperate woody species. *Plant Cell Environ.* **2003**, *26*, 1343–1356.
10. Öunapuu, E.; Sellin, A. Daily dynamics of leaf and soil-to-branch hydraulic conductance in silver birch (*Betula pendula*) measured in situ. *Plant Physiol. Biochem.* **2013**, *68*, 104–110.
11. Newman, J.A.; Anand, M.; Henry, H.A.L.; Hunt, S.; Gedalof, Z. *Climate Change Biology*; CABI Publishing: Wallingford, UK, 2011.
12. Gratani, L. Plant Phenotypic Plasticity in Response to Environmental Factors. *Adv. Bot.* **2014**, 208747. <https://doi.org/10.1155/2014/208747>.
13. Nicotra, A.B.; Atkin, O.K.; Bonser, S.P.; Davidson, A.M.; Finnegan, E.J.; Mathesius, U.; Poot, P.; Purugganan, M.D.; Richards, C.L.; Valladares, F.; et al. Plant phenotypic plasticity in a changing climate. *Trends Plant Sci.* **2010**, *15*, 684–692.
14. Henn, J.J.; Buzzard, V.; Enquist, B.J.; Halbritter, A.H.; Klanderuds, K.; Maitner, B.S.; Michaletz, S.T.; Pötsch, C.; Seltzer, L.; Telford, R.J.; et al. Intraspecific trait variation and phenotypic plasticity mediate alpine plant species response to climate change. *Front. Plant Sci.* **2018**, *9*, 1548. <https://doi.org/10.3389/fpls.2018.01548>.
15. Mediavilla, S.; Martin, I.; Babiano, J.; Escudero, A. Foliar plasticity related to gradients of heat and drought stress across crown orientations in three Mediterranean *Quercus* species. *PLoS ONE* **2019**, *14*, e0224462. <https://doi.org/10.1371/journal.pone.0224462>.
16. Valladares, F.; Gianoli, E.; Gomez, J.M. Ecological limits to plant phenotypic plasticity. *New Phytol.* **2007**, *176*, 749–763.
17. Sultan, S.E. Phenotypic plasticity for plant development, function and life history. *Trends Plant Sci.* **2000**, *5*, 537–542.
18. Matesanz, S.; Gianoli, E.; Valladares, F. Global change and the evolution of phenotypic plasticity in plants. *Ann. N. Y. Acad. Sci.* **2010**, *1206*, 35–55.
19. IPCC. Climate Change 2021: The Physical Science Basis. *Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change*; Masson-Delmotte, V., Zhai, P., Pirani, A., Connors, S.L., Péan, C., Berger, S., Caud, N., Chen, Y., Goldfarb, L., Gomis, M.I.; et al., Eds.; Cambridge University Press: Cambridge, MA, USA, 2021.
20. Kont, A.; Jaagus, J.; Aunap, R. Climate change scenarios and the effect of sea-level rise for Estonia. *Glob. Planet. Chang.* **2003**, *36*, 1–15.
21. Betts, A.K.; Desjardins, R.; Worth, D.; Beckage, B. Climate coupling between temperature, humidity, precipitation, and cloud cover over the Canadian Prairies. *J. Geophys. Res. -Atmos.* **2014**, *119*, 13305–13326.
22. Westra, S.; Fowler, H.J.; Evans, J.P.; Alexander, L.V.; Berg, P.; Johnson, F.; Kendon, E.J.; Lenderink, G.; Roberts, N.M. Future changes to the intensity and frequency of short-duration extreme rainfall. *Rev. Geophys.* **2014**, *52*, 522–555.
23. Busuioc, A.; Birsan, M.V.; Carbutaru, D.; Baciu, M.; Orzan, A. Changes in the large-scale thermodynamic instability and connection with rain shower frequency over Romania: Verification of the Clausius-Clapeyron scaling. *Int. J. Climatol.* **2016**, *36*, 2015–2034.
24. Fanourakis, D.; Aliniaefard, S.; Sellin, A.; Giday, H.; Körner, O.; Nejad, A.R.; Delis, C.; Bouranis, D.; Koubouris, G.; Kambourakis, E.; et al. Stomatal behavior following mid or long-term exposure to high relative air humidity: A review. *Plant Physiol. Biochem.* **2020**, *153*, 92–105.
25. Niu, S.L.; Luo, Y.Q.; Li, D.J.; Cao, S.H.; Xia, J.Y.; Li, J.W.; Smith, M.D. Plant growth and mortality under climatic extremes: An overview. *Environ. Exp. Bot.* **2014**, *98*, 13–19.
26. Zang, C.; Hartl-Meier, C.; Dittmar, C.; Rothe, A.; Menzel, A. Patterns of drought tolerance in major European temperate forest trees: Climatic drivers and levels of variability. *Glob. Chang. Biol.* **2014**, *20*, 3767–3779.
27. Oksanen, E.; Lihavainen, J.; Keinänen, M.; Keski-Saari, S.; Kontunen-Soppela, S.; Sellin, A.; Söber, A. Northern forest trees under increasing atmospheric humidity. In *Progress in Botany*; Canovas, F.M., Lüttge, U., Matyssek, R., Pretzsch, H., Eds.; Springer: Cham, Switzerland, 2018; Volume 80.
28. Sellin, A.; Tullus, A.; Niglas, A.; Öunapuu, E.; Karusion, A.; Löhmus, K. Humidity-driven changes in growth rate, photosynthetic capacity, hydraulic properties and other functional traits in silver birch (*Betula pendula*). *Ecol. Res.* **2013**, *28*, 523–535.
29. Sellin, A.; Taneda, H.; Alber, M. Leaf structural and hydraulic adjustment with respect to air humidity and canopy position in silver birch (*Betula pendula*). *J. Plant Res.* **2019**, *132*, 369–381.
30. Sellin, A.; Niglas, A.; Öunapuu-Pikas, E.; Kupper, P. Rapid and long-term effects of water deficit on gas exchange and hydraulic conductance of silver birch trees grown under varying atmospheric humidity. *BMC Plant Biol.* **2014**, *14*, 72. <https://doi.org/10.1186/1471-2229-14-72>.
31. Pritzkow, C.; Williamson, V.; Szota, C.; Trouve, R.; Arndt, S.K. Phenotypic plasticity and genetic adaptation of functional traits influences intra-specific variation in hydraulic efficiency and safety. *Tree Physiol.* **2020**, *40*, 215–229.
32. Tyree, M.T.; Nardini, A.; Salleo, S.; Sack, L.; El Omari, B. The dependence of leaf hydraulic conductance on irradiance during HPFM measurements: Any role for stomatal response? *J. Exp. Bot.* **2005**, *56*, 737–744.
33. Sellin, A.; Öunapuu, E.; Kupper, P. Effects of light intensity and duration on leaf hydraulic conductance and distribution of resistance in shoots of silver birch (*Betula pendula*). *Physiol. Plant.* **2008**, *134*, 412–420.
34. Voicu, M.C.; Zwiazek, J.J.; Tyree, M.T. Light response of hydraulic conductance in bur oak (*Quercus macrocarpa*) leaves. *Tree Physiol.* **2008**, *28*, 1007–1015.
35. Guyot, G.; Scoffoni, C.; Sack, L. Combined impacts of irradiance and dehydration on leaf hydraulic conductance: Insights into vulnerability and stomatal control. *Plant Cell Environ.* **2012**, *35*, 857–871.

36. Ohtsuka, A.; Sack, L.; Taneda, H. Bundle sheath lignification mediates the linkage of leaf hydraulics and venation. *Plant Cell Environ.* **2018**, *41*, 342–353.
37. Xiong, D.L.; Douthe, C.; Flexas, J. Differential coordination of stomatal conductance, mesophyll conductance, and leaf hydraulic conductance in response to changing light across species. *Plant Cell Environ.* **2018**, *41*, 436–450.
38. Scoffoni, C.; Kunkle, J.; Pasquet-Kok, J.; Vuong, C.; Patel, A.J.; Montgomery, R.A.; Givnish, T.J.; Sack, L. Light-induced plasticity in leaf hydraulics, venation, anatomy, and gas exchange in ecologically diverse Hawaiian lobeliads. *New Phytol.* **2015**, *207*, 43–58.
39. Öunapuu-Pikas, E.; Sellin, A. Plasticity and light sensitivity of leaf hydraulic conductance to fast changes in irradiance in common hazel (*Corylus avellana* L.). *Plant Sci.* **2020**, *290*, 110299. <https://doi.org/10.1016/j.plantsci.2019.110299>.
40. Possen, B.J.H.M.; Oksanen, E.; Rousi, M.; Ruhanen, H.; Ahonen, V.; Tervahauta, A.; Heinonen, J.; Heiskanen, J.; Kärenlampi, S.; Vapaavuori, E. Adaptability of birch (*Betula pendula* Roth) and aspen (*Populus tremula* L.) genotypes to different soil moisture conditions. *For. Ecol. Manag.* **2011**, *262*, 1387–1399.
41. Hynynen, J.; Niemisto, P.; Vihera-Aarnio, A.; Brunner, A.; Hein, S.; Velling, P. Silviculture of birch (*Betula pendula* Roth and *Betula pubescens* Ehrh.) in northern Europe. *Forestry* **2010**, *83*, 103–119.
42. Tullus, A.; Rytter, L.; Tullus, T.; Weih, M.; Tullus, H. Short-rotation forestry with hybrid aspen (*Populus tremula* L. × *P. tremuloides* Michx.) in Northern Europe. *Scand. J. For. Res.* **2012**, *27*, 10–29.
43. Sellin, A.; Öunapuu, E.; Karusion, A. Experimental evidence supporting the concept of light-mediated modulation of stem hydraulic conductance. *Tree Physiol.* **2010**, *30*, 1528–1535.
44. Zapater, M.; Breda, N.; Bonal, D.; Pardonnet, S.; Granier, A. Differential response to soil drought among co-occurring broad-leaved tree species growing in a 15-to 25-year-old mixed stand. *Ann. For. Sci.* **2013**, *70*, 31–39.
45. Kupper, P.; Ivanova, H.; Söber, A.; Rohula-Okunev, G.; Sellin, A. Night and daytime water relations in five fast-growing tree species: Effects of environmental and endogenous variables. *Ecophysiology* **2018**, *11*, e1927. <https://doi.org/10.1002/eco.1927>.
46. Aasamaa, K.; Söber, A.; Hartung, W.; Niinemets, Ü. Drought acclimation of two deciduous tree species of different layers in a temperate forest canopy. *Trees-Struct. Funct.* **2004**, *18*, 93–101.
47. Aasamaa, K.; Kõivik, K.; Kupper, P.; Söber, A. Growth environment determines light sensitivity of shoot hydraulic conductance. *Ecol. Res.* **2014**, *29*, 143–151.
48. Russak, V. Päikesekiirgus. *Tartu Kliima Ja Selle Muutumine Viimastel Kümnenditel*; Ross, J., Ed.; Eesti Teaduste Akadeemia: Tartu, Estonia, 1990.
49. Jaagus, J.; Mändla, K. Climate change scenarios for Estonia based on climate models from the IPCC Fourth Assessment Report. *Est. J. Earth Sci.* **2014**, *63*, 166–180.
50. Jacob, D.; Petersen, J.; Eggert, B.; Alias, A.; Christensen, O.B.; Bouwer, L.M.; Braun, A.; Colette, A.; Déqué, M.; Georgievski, G.; et al. EURO-CORDEX: New high-resolution climate change projections for European impact research. *Reg. Environ. Chang.* **2014**, *14*, 563–578.
51. Kupper, P.; Söber, J.; Sellin, A.; Löhmus, K.; Tullus, A.; Räm, O.; Lubenets, K.; Tulva, I.; Uri, V.; Zobel, M.; et al. An experimental facility for free air humidity manipulation (FAHM) can alter water flux through deciduous tree canopy. *Environ. Exp. Bot.* **2011**, *72*, 432–438.
52. Tullus, A.; Kupper, P.; Sellin, A.; Parts, L.; Söber, J.; Tullus, T.; Tullus, H. Climate change at Northern latitudes: Rising atmospheric humidity decreases transpiration, N-uptake and growth rate of hybrid aspen. *PLoS ONE* **2012**, *7*, e42648. <https://doi.org/10.1371/journal.pone.0042648>.
53. Sellin, A.; Alber, M.; Keinanen, M.; Kupper, P.; Lihavainen, J.; Löhmus, K.; Tullus, A. Growth of northern deciduous trees under increasing atmospheric humidity: Possible mechanisms behind the growth retardation. *Reg. Environ. Chang.* **2017**, *17*, 2135–2148.
54. Heimonen, K.; Valtonen, A.; Kontunen-Soppela, S.; Keski-Saari, S.; Rousi, M.; Oksanen, E.; Roininen, H. Insect herbivore damage on latitudinally translocated silver birch (*Betula pendula*)—Predicting the effects of climate change. *Clim. Chang.* **2015**, *131*, 245–257.
55. Valladares, F.; Martinez-Ferri, E.; Balaguer, L.; Perez-Corona, E.; Manrique, E. Low leaf-level response to light and nutrients in Mediterranean evergreen oaks: A conservative resource-use strategy? *New Phytol.* **2000**, *148*, 79–91.
56. Sack, L.; Melcher, P.J.; Zwieniecki, M.A.; Holbrook, N.M. The hydraulic conductance of the angiosperm leaf lamina: A comparison of three measurement methods. *J. Exp. Bot.* **2002**, *53*, 2177–2184.
57. Nardini, A.; Salleo, S.; Andri, S. Circadian regulation of leaf hydraulic conductance in sunflower (*Helianthus annuus* L. cv Margot). *Plant Cell Environ.* **2005**, *28*, 750–759.
58. Scoffoni, C.; Pou, A.; Aasamaa, K.; Sack, L. The rapid light response of leaf hydraulic conductance: New evidence from two experimental methods. *Plant Cell Environ.* **2008**, *31*, 1803–1812.
59. Venisse, J.-S.; Öunapuu-Pikas, E.; Dupont, M.; Gousset-Dupont, A.; Saadaoui, M.; Faize, M.; Chen, S.; Chen, S.; Petel, G.; Fumanal, B.; et al. Genome-wide identification, structure characterization, and expression pattern profiling of the aquaporin gene family in *Betula pendula*. *Int. J. Mol. Sci.* **2021**, *22*, 7269. <https://doi.org/10.3390/ijms22147269>.
60. Cochard, H.; Venisse, J.-S.; Barigah, T.S.; Brunel, N.; Herbette, S.; Guilliot, A.; Tyree, M.T.; Sakr, S. Putative role of aquaporins in variable hydraulic conductance of leaves in response to light. *Plant Physiol.* **2007**, *143*, 122–133.
61. Ben Baaziz, K.; Lopez, D.; Bouzid, S.; Cochard, H.; Venisse, J.S.; Sakr, S. Early gene expression in the walnut tree occurring during stimulation of leaf hydraulic conductance by irradiance. *Biol. Plant.* **2012**, *56*, 657–666.

62. Ben Baaziz, K.; Lopez, D.; Rabot, A.; Combes, D.; Gousset, A.; Bouzid, S.; Venisse, J.S. Light-mediated K-leaf induction and contribution of both the PIP1s and PIP2s aquaporins in five tree species: Walnut (*Juglans regia*) case study. *Tree Physiol.* **2012**, *32*, 423–434.
63. Lopez, D.; Venisse, J.S.; Fumanal, B.; Chaumont, F.; Guillot, E.; Daniels, M.J.; Gousset-Dupont, A. Aquaporins and leaf hydraulics: Poplar sheds new light. *Plant Cell Physiol.* **2013**, *54*, 1963–1975.
64. Miniussi, M.; Del Terra, L.; Savi, T.; Pallavicini, A.; Nardini, A. Aquaporins in *Coffea arabica* L.: Identification, expression, and impacts on plant water relations and hydraulics. *Plant Physiol. Biochem.* **2015**, *95*, 92–102.
65. Prado, K.; Boursiac, Y.; Tournaire-Roux, C.; Monneuse, J.M.; Postaire, O.; Da Ines, O.; Maurel, C. Regulation of *Arabidopsis* leaf hydraulics involves light-dependent phosphorylation of aquaporins in veins. *Plant Cell* **2013**, *25*, 1029–1039.
66. Prado, K.; Cotelle, V.; Li, G.W.; Bellati, J.; Tang, N.; Tournaire-Roux, C.; Martinière, A.; Santoni, V.; Maurel, C. Oscillating aquaporin phosphorylation and 14-3-3 proteins mediate the circadian regulation of leaf hydraulics. *Plant Cell* **2019**, *31*, 417–429.
67. Groszmann, M.; Osborn, H.L.; Evans, J.R. Carbon dioxide and water transport through plant aquaporins. *Plant Cell Environ.* **2017**, *40*, 938–961.
68. Scoffoni, C.; Albuquerque, C.; Cochard, H.; Buckley, T.N.; Fletcher, L.R.; Caringella, M.A.; Sack, L. The causes of leaf hydraulic vulnerability and its influence on gas exchange in *Arabidopsis thaliana*. *Plant Physiol.* **2018**, *178*, 1584–1601.
69. Trifilo, P.; Petruzzellis, F.; Abate, E.; Nardini, A. The extra-vascular water pathway regulates dynamic leaf hydraulic decline and recovery in *Populus nigra*. *Physiol. Plant.* **2021**, *172*, 29–40.
70. Sellin, A.; Alber, M.; Jasińska, A.K.; Rosenthal, K. Adjustment of leaf anatomical and hydraulic traits across vertical canopy profiles of young broadleaved forest stands. *Trees-Struct. Funct.* **2022**, *36*, 67–80.
71. Sellin, A.; Kupper, P. Variation in leaf conductance of silver birch: Effects of irradiance, vapour pressure deficit, leaf water status and position within a crown. *For. Ecol. Manag.* **2005**, *206*, 153–166.
72. Sellin, A.; Eensalu, E.; Niglas, A. Is distribution of hydraulic constraints within tree crowns reflected in photosynthetic water-use efficiency? An example of *Betula pendula*. *Ecol. Res.* **2010**, *25*, 173–183.
73. Kangur, O.; Kupper, P.; Sellin, A. Predawn disequilibrium between soil and plant water potentials in light of climate trends predicted for northern Europe. *Reg. Environ. Chang.* **2017**, *17*, 2159–2168.
74. Sellin, A.; Alber, M.; Kupper, P. Increasing air humidity influences hydraulic efficiency but not functional vulnerability of xylem in hybrid aspen. *J. Plant Physiol.* **2017**, *219*, 28–36.
75. Azad, A.K.; Sawa, Y.; Ishikawa, T.; Shibata, H. Phosphorylation of plasma membrane aquaporin regulates temperature-dependent opening of tulip petals. *Plant Cell Physiol.* **2004**, *45*, 608–617.
76. Hedfalk, K.; Tornroth-Horsefield, S.; Nyblom, M.; Johanson, U.; Kjellbom, P.; Neutze, R. Aquaporin gating. *Curr. Opin. Struct. Biol.* **2006**, *16*, 447–456.
77. Maurel, C.; Boursiac, Y.; Luu, D.T.; Santoni, V.; Shahzad, Z.; Verdoucq, L. Aquaporins in plants. *Physiol. Rev.* **2015**, *95*, 1321–1358.
78. Tyree, M.T.; Ewers, F.W. The hydraulic architecture of trees and other woody plants. *New Phytol.* **1991**, *119*, 345–360.
79. Tyree, M.T.; Zimmermann, M.H. *Xylem Structure and the Ascent of Sap*, 2nd ed.; Springer: Berlin/Heidelberg, Germany, 2002.
80. Zwieniecki, M.A.; Melcher, P.J.; Holbrook, N.M. Hydrogel control of xylem hydraulic resistance in plants. *Science* **2001**, *291*, 1059–1062.
81. Nardini, A.; Salleo, S.; Jansen, S. More than just a vulnerable pipeline: Xylem physiology in the light of ion-mediated regulation of plant water transport. *J. Exp. Bot.* **2011**, *62*, 4701–4718.
82. Zwieniecki, M.A.; Secchi, F. Getting variable xylem hydraulic resistance under control: Interplay of structure and function. *Tree Physiol.* **2012**, *32*, 1431–1433.
83. Nardini, A.; Grego, F.; Trifilo, P.; Salleo, S. Changes of xylem sap ionic content and stem hydraulics in response to irradiance in *Laurus nobilis*. *Tree Physiol.* **2010**, *30*, 628–635.
84. Lee, J.; Holbrook, N.M.; Zwieniecki, M.A. Ion induced changes in the structure of bordered pit membranes. *Front. Plant Sci.* **2012**, *3*, 55. <https://doi.org/10.3389/fpls.2012.00055>.