



HAL
open science

A preliminary study of fin rays microchemistry data in *Salmo trutta*: Is it possible to trace back to the original watershed from fin rays ?

Guénard Maxime, Tewann Beauchard, Jean-Christophe Aymes, Françoise Daverat, Francois Gueraud

► **To cite this version:**

Guénard Maxime, Tewann Beauchard, Jean-Christophe Aymes, Françoise Daverat, Francois Gueraud. A preliminary study of fin rays microchemistry data in *Salmo trutta*: Is it possible to trace back to the original watershed from fin rays ?. Nowpas 2022, Mar 2022, Saint Etienne de Baigorri, France. hal-03643410

HAL Id: hal-03643410

<https://hal.inrae.fr/hal-03643410>

Submitted on 15 Apr 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.


L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

➤ A preliminary study of fin rays microchemistry data in *Salmo trutta* : Is it possible to trace back to the original watershed from fin rays ?

Guénard.M ^{1,2}, Beauchard. T², Aymes. J.C ¹, Daverat. F ¹, Guéraud. F ¹

1. INRAE, ECOBIOP, St-Pée-sur-Nivelle 2. University of Pau and Pays de l'Adour (UPPA), Master 2 DYNEA

➤ Kerguelen islands : an open-air laboratory



INRAE

A preliminary study of fin rays microch
04/03/2022 / NoWPas 2022 / Guénard Maxime



Réserve Naturelle
TERRES AUSTRALES
FRANÇAISES


2006



2019

➤ A context of colonization of the Kerguelen islands by the common trout (*Salmo trutta*)

| Species | Number of individuals released into the river | Number of rivers seeded | Number of rivers currently colonised |
|-----------------|---|-------------------------|--------------------------------------|
| <i>S.trutta</i> | 83 000 | 12 | 42 |



(Labonne et al., 2013)

Introduction of salmonids since 1962


Dispersion of *S.trutta* by anadromous phenotype



(F.GUERAUD, technician at INRAE on site)



➤ Life cycle of trout in Kerguelen islands



(Labonne et al., 2013)

★ Minimum age for going to sea is 2 years (Sea trout)

➤ A large collection of anatomical parts

INRAE (COLISA) has a large collection of anatomical parts dating back to 1962, such as scales in majority but also fin rays.

| Species | number of captures | Pockets of scales | Adipose fin | Tip of the fin | Ray of pectorals | Stomach content | Otoliths | Whole fish |
|--------------------------|--------------------|-------------------|-------------|----------------|------------------|-----------------|----------|------------|
| Salmo trutta | 91853 | 37593 | 6074 | 1177 | 583 | 3508 | 1612 | 1107 |
| Salvelinus fontinalis | 48983 | 14348 | 905 | 88 | 3 | 2146 | 458 | 65 |
| Salmo salar | 8909 | 4452 | 170 | 0 | 0 | 48 | 63 | 0 |
| Oncorhynchus kisutch | 5094 | 3134 | 100 | 0 | 0 | 15 | 451 | 3 |
| Salvelinus alpinus | 363 | 160 | 84 | 48 | 0 | 49 | 50 | 0 |
| Oncorhynchus tshawytscha | 51 | 51 | 0 | 0 | 0 | 0 | 4 | 0 |
| Oncorhynchus mykiss | 25 | 25 | 0 | 0 | 0 | 0 | 0 | 0 |
| Inconnu | 938 | 556 | 22 | 1 | 1 | 4 | 60 | 122 |
| Total | 156216 | 60319 | 7355 | 1314 | 587 | 5770 | 2698 | 1297 |

A study was being carried out on the fin rays of trout and one of the objectives of this study was to find out if :

Is it possible to trace back to the original watershed from microchemistry data collected by the use of fin rays ?

Why sclerochronology to study the dispersion of individuals?

Incorporation of chemical elements

Each watershed has its own chemical signature. It is the result of the geological composition and environmental conditions


These chemical elements are transported and then assimilated in the calcified parts (Bones) during the growth of the individuals.

Sr
Ba
Mn
Mg
Zn
Cu
Fe

Sr

Ba
Mn
Mg
Zn
Cu
Fe

Sr
Ba
Mn
Mg
Zn
Cu
Fe



(Rogissart d'après Campana, 1999)




INRAE

A preliminary study of fin rays microchemistry data in *Salmo trutta*


04/03/2022 / NoWPas 2022 / Guénard Maxime

➤ Method


From fin rays to microchemistry data



Calcium phosphate structure (Beamish, 1981)



(Pierre FEUTRY modifié d'après Tabouret 2009)



(M. GUENARD)




INRAE

A preliminary study of fin rays microchemistry data in *Salmo trutta*

04/03/2022 / NoWPas 2022 / Guénard Maxime

➤ Study site and sampling

Study on 3 of the main rivers studied





| Populations | Sedentary trouts | Sea trouts | Total |
|-------------|------------------|------------|-------|
| CHÂTEAU | 11 | 11 | 22 |
| MANCHOTS | 6 | 15 | 21 |
| NORVEGIENNE | 6 | 16 | 22 |
| Total | 23 | 42 | 65 |

Age > 4 years

➤ Results

Sedentary trouts




Clear watershed discrimination for sedentary individuals with the PCA and the HCA .

The chemical compositions of watersheds are sufficiently different to allow discrimination between individuals according to their river.

So we could in principle reallocate individuals to their watersheds



➤ Results Sea trouts



Overlapping data makes it impossible to discriminate groups in any dimensions of the PCA

Chemical data do not allow to discriminate the watersheds with the HCA



> Results

Sea trouts

QDA and Jackknife reclassification

| | CHÂTEAU (reassignment) % | MANCHOTS (reassignment) % | NORVEGIENNE (reassignment) % |
|----------------------|--------------------------|---------------------------|------------------------------|
| CHÂTEAU (origin) | 0 | 73 | 27 |
| MANCHOTS (origin) | 0 | 67 | 33 |
| NORVEGIENNE (origin) | 0 | 56 | 44 |

The QDA coupled with the jackknife reclassification shows us a very low percentage of good reclassification in hindsight.

So, we will not be able to reassign individuals to their watersheds of origin.

➤ Discussion

A disappointing record



Sedentary trouts

Satisfying discrimination of watersheds of origin explained by environmental conditions (hydromorphology and geology of the site).



Sea trouts

The discrimination of the watersheds of origin is not satisfying.

HYPOTHESE : remobilization (structural modification of the bone matrix) of elements during the stay at sea.


➤ To conclude


Because this study was not designed to incorporate geologically disparate natal sources, it may be more effective on a larger scale.

Moreover, an isotopic analysis ($^{86/87}\text{Sr}$) of fin rays could allow us to discriminate more efficiently Or may be not !

➤ Future tests

This year, we want to study the food web of the Brown trout in Kerguelen islands and study the variation since the 60s from carbon $\delta^{13}\text{C}$ and nitrogen $\delta^{15}\text{N}$ isotopes from the scales.





➤ Thank you for your attention !



INRAE


A preliminary study of fin rays microchemistry data in *Salmo trutta*
04/03/2022 / NoWPas 2022 / Guénard Maxime

➤ continuation of the method


Sclerochronology

Scale : Age, age at sea, number of spawnings, number of returns to fresh water...but **not chemistry data** (Vignon et al., 2020).

Otoliths are generally use to provide individual data on fish structure and composition of ecology (Age, migratory status, habitats, etc.) and chemistry data. However, otoliths require the **sacrifice** of the fish, that is why we are interested in fin rays.



➤ Elements linked to Habitats or Physiology



(De Pontual et Geffen, 2002 d'après Campana, 1999)