



HAL
open science

Nitrogen isotopic discrimination as a biomarker of between-cow variation in the efficiency of nitrogen utilization for milk production: A meta-analysis

M. Correa-Luna, M. Johansen, Pierre Noziere, C. Chantelauze, S.M. Nasrollahi, P. Lund, M. Larsen, A.R. Bayat, L.A. Crompton, C.K. Reynolds, et al.

► To cite this version:

M. Correa-Luna, M. Johansen, Pierre Noziere, C. Chantelauze, S.M. Nasrollahi, et al.. Nitrogen isotopic discrimination as a biomarker of between-cow variation in the efficiency of nitrogen utilization for milk production: A meta-analysis. *Journal of Dairy Science*, 2022, 105 (6), pp.5004-5023. 10.3168/jds.2021-21498 . hal-03658611

HAL Id: hal-03658611

<https://hal.inrae.fr/hal-03658611v1>

Submitted on 12 May 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License



Nitrogen isotopic discrimination as a biomarker of between-cow variation in the efficiency of nitrogen utilization for milk production: A meta-analysis

M. Correa-Luna,¹ M. Johansen,² P. Noziere,¹ C. Chantelauze,^{1,3} S. M. Nasrollahi,¹ P. Lund,² M. Larsen,² A. R. Bayat,⁴ L. A. Crompton,⁵ C. K. Reynolds,⁵ E. Froidmont,⁶ N. Edouard,⁷ R. Dewhurst,⁸ L. Bahloul,⁹ C. Martin,¹ and G. Cantalapiedra-Hijar^{1*}

¹Université Clermont Auvergne, INRAE, UMR Herbivores, F-63000 Clermont-Ferrand, France

²Department of Animal Science, Aarhus University, AU Foulum, PO Box 50, DK-8830, Tjele, Denmark

³Université Clermont Auvergne, INRAE, UNH, Plateforme d'Exploration du Métabolisme, MetaboHUB, Clermont, F-63000 Clermont-Ferrand, France

⁴Milk Production Solutions, Production Systems, Natural Resources Institute Finland (Luke), FI 31600 Jokioinen, Finland

⁵Centre for Dairy Research, Department of Animal Sciences, School of Agriculture, Policy, and Development, University of Reading, Reading, RG6 6AH, United Kingdom

⁶Walloon Agricultural Research Center (CRA-W), B-5030 Gembloux, Belgium

⁷INRAE, Agrocampus-Ouest, PEGASE, 35590 Saint-Gilles, France

⁸SRUC, West Mains Road, Edinburgh, EH9 3JG, United Kingdom

⁹Adisseo France S.A.S., 92160 Antony, France

ABSTRACT

Estimating the efficiency of N utilization for milk production (MNE) of individual cows at a large scale is difficult, particularly because of the cost of measuring feed intake. Nitrogen isotopic discrimination ($\Delta^{15}\text{N}$) between the animal (milk, plasma, or tissues) and its diet has been proposed as a biomarker of the efficiency of N utilization in a range of production systems and ruminant species. The aim of this study was to assess the ability of $\Delta^{15}\text{N}$ to predict the between-animal variability in MNE in dairy cows using an extensive database. For this, 20 independent experiments conducted as either changeover ($n = 14$) or continuous ($n = 6$) trials were available and comprised an initial data set of 1,300 observations. Between-animal variability was defined as the variation observed among cows sharing the same contemporary group (CG; individuals from the same experimental site, sampling period, and dietary treatment). Milk N efficiency was calculated as the ratio between mean milk N (grams of N in milk per day) and mean N intake (grams of N intake per day) obtained from each sampling period, which lasted 9.0 ± 9.9 d (mean \pm SD). Samples of milk ($n = 604$) or plasma ($n = 696$) and feeds (74 dietary treatments) were analyzed for natural ^{15}N abundance ($\delta^{15}\text{N}$), and then the N isotopic discrimination between the animal and the dietary treatment was calculated ($\Delta^{15}\text{N} = \delta^{15}\text{N}_{\text{animal}}$

$-\delta^{15}\text{N}_{\text{diet}}$). Data were analyzed through mixed-effect regression models considering the experiment, sampling period, and dietary treatment as random effects. In addition, repeatability estimates were calculated for each experiment to test the hypothesis of improved predictions when MNE and $\Delta^{15}\text{N}$ measurements errors were lower. The considerable protein mobilization in early lactation artificially increased both MNE and $\Delta^{15}\text{N}$, leading to a positive rather than negative relationship, and this limited the implementation of this biomarker in early lactating cows. When the experimental errors of $\Delta^{15}\text{N}$ and MNE decreased in a particular experiment (i.e., higher repeatability values), we observed a greater ability of $\Delta^{15}\text{N}$ to predict MNE at the individual level. The predominant negative and significant correlation between $\Delta^{15}\text{N}$ and MNE in mid- and late lactation demonstrated that on average $\Delta^{15}\text{N}$ reflects MNE variations both across dietary treatments and between animals. The root mean squared prediction error as a percentage of average observed value was 6.8%, indicating that the model only allowed differentiation between 2 cows in terms of MNE within a CG if they differed by at least 0.112 g/g of MNE (95% confidence level), and this could represent a limitation in predicting MNE at the individual level. However, the one-way ANOVA performed to test the ability of $\Delta^{15}\text{N}$ to differentiate within-CG the top 25% from the lowest 25% individuals in terms of MNE was significant, indicating that it is possible to distinguish extreme animals in terms of MNE from their N isotopic signature, which could be useful to group animals for precision feeding.

Key words: meta-analysis, milk nitrogen efficiency, biomarker, individual variability, ^{15}N

Received October 28, 2021.

Accepted February 21, 2022.

*Corresponding author: gonzalo.cantalapiedra@inrae.fr

INTRODUCTION

Dairy products are important sources of food protein along with a range of other essential nutrients (Visioli and Strata, 2014), and their increased consumption is driven by the growth of the world human population and their average incomes (Scott, 2017). Total food production is a significant contributor to global greenhouse gas emissions, which are undeniably related to climate change (Clark et al., 2020; Ocko et al., 2021). There are 2 main sources of environmental pollution in livestock systems: greenhouse gas emissions per se (carbon dioxide, methane, and nitrous oxide; Uwizeye et al., 2020; Ocko et al., 2021) and the negative impact of excreta (mainly N and P) on the quality of surface and ground water (Castillo et al., 2000; Uwizeye et al., 2020). In this context, mitigation strategies for the livestock industry are highly needed (Uwizeye et al., 2020).

In the lactating cow, the efficiency of N utilization for milk production (MNE; g of milk N/g of N intake) is commonly used to describe the conversion of feed N inputs into dairy products (Cantalapiedra-Hijar et al., 2016) and also as an indicator of N losses to the environment (Jonker et al., 1998; Castillo et al., 2000; Nousiainen et al., 2004). The main constraint to collection of accurate estimations of MNE at the individual cow level is the determination of feed intake, which is costly and laborious (Hellwing et al., 2015). The identification and consolidation of techniques to predict MNE accurately from easy-to-collect samples will contribute to the design of feed rations according to nutritional status and to increasing the collection of records for breeding programs (Brito et al., 2021).

In the context of animal physiology, a biomarker can be defined as “a naturally occurring molecule, gene, or characteristic by which a particular pathological or physiological process, disease, etc., can be identified or referred to” (Oxford Dictionary; <https://www.lexico.com>). Ruminants have an effective internal N recycling system, where most of the excess dietary N is converted to urea in the liver through ureagenesis, designed to avoid toxic effects if ammonia enters the systemic circulation (Lapierre et al., 2005). In turn, urea is transported from the plasma to other body fluids such as saliva to be recycled, as well as to the kidneys to be excreted. Because of its low molecular weight and neutral charge, urea easily diffuses across cellular membranes where it is incorporated to milk as MUN (Jonker et al., 1998). On this basis, MUN has been proposed as a biomarker for MNE and N excretion in dairy cows. However, the evidence regarding the potential of this biomarker to reflect the between-animal variation in MNE (Spek et

al., 2013; Huhtanen et al., 2015) and its association with N partitioning at the individual animal level (Spek et al., 2013; Beatson et al., 2019) is inconclusive.

Alternatively, the natural ^{15}N abundance ($\delta^{15}\text{N}$; $^{15}\text{N}/^{14}\text{N}$ ratio relative to atmospheric N_2) in animal protein is a promising biomarker for predicting MNE because of its direct link with ruminal microbial N metabolism (Wattiaux and Reed, 1995) and with the catabolism of AA in the liver (Cantalapiedra-Hijar et al., 2015). In short, it has been demonstrated across a variety of conditions and species that ^{15}N natural abundance in animal proteins is higher than in the diet consumed (Deniro and Epstein, 1981) and that N isotopic discrimination ($\Delta^{15}\text{N} = \delta^{15}\text{N}_{\text{animal}} - \delta^{15}\text{N}_{\text{diet}}$) is negatively correlated with N use efficiency (NUE), estimated as g of milk N or retained body N per grams of N intake (Cantalapiedra-Hijar et al., 2018). This discrimination phenomenon has been confirmed to differ at the individual level, which could be advantageous in the attempt to rank ruminants reared under similar conditions for NUE (Cheng et al., 2013; Cantalapiedra-Hijar et al., 2018) or for feed efficiency (Wheadon et al., 2014; Guarnido-Lopez et al., 2021). However, not all studies found a significant negative relationship between MNE and $\Delta^{15}\text{N}$ in lactating dairy cows (Cheng et al., 2011; Chen et al., 2020). In a recent study by Chen et al. (2020), the N isotopic signatures were strongly influenced by protein mobilization occurring during early lactation, and this resulted in positive, rather than negative, associations with MNE. Another explanation for the disparity in the associations between MNE and $\Delta^{15}\text{N}$ could be related to a high experimental error associated with the measurements of N intake, milk N, or N isotopic signatures. This experimental error can be assessed statistically by analyzing the consistency of repeated measurements (Harper, 1994). Although guidelines and quality standards for measuring these traits exist, it has been hypothesized that higher repeatability values (i.e., lower experimental errors) of both NUE and $\Delta^{15}\text{N}$ measurements could lead to improved model MNE prediction.

In the present study, we explored the ability of $\Delta^{15}\text{N}$ to predict between-animal variability in terms of MNE in lactating dairy cows and potential factors affecting the prediction ability of $\Delta^{15}\text{N}$. In our previous meta-analysis (Cantalapiedra-Hijar et al., 2018), the association between NUE and $\Delta^{15}\text{N}$ was explored as the proof of concept from a range of ruminant species and production conditions, employing a smaller data set. The present study brings an update and refinement of the model, with a larger data set comprising only lactating dairy cows.

MATERIALS AND METHODS

Experimental Data

A database including individual animal measurements was created from experiments proposed by the partners of the SmartCow Project (grant agreement no. 730924), a collaborative EU project aiming at the integration of research infrastructures for the European cattle sector (<https://www.smartcow.eu>). Data originated from 20 dairy milk production experiments (**ID1** to **ID20**) conducted in Belgium (n = 1), England (n = 1), Finland (n = 2), Denmark (n = 6), and France (n = 10). These experiments were conducted as either changeover (e.g., Latin square; n = 14) or continuous (n = 6) experiments. The initial data set included multiple observations from 425 cows (i.e., different sampling period and dietary treatments) representing a total of 1,300 individual observations of N intake, milk N, MNE, and $\Delta^{15}\text{N}$. A summary of studies, along with their corresponding designs, is presented in Table 1.

Laboratory Analysis and Calculations

For both individual animal observations and dietary treatment (**DT**) means, values of MNE were calculated as the ratio between milk N (**MN**, g/d) and N intake (**NI**, g/d), considering all observations of the corresponding sampling period (**SP**) to account for

daily variability in the observations. Nitrogen intake was calculated by multiplying dietary N content (g of N/100 g of DM) by the daily DMI corresponding to each SP for each cow. In the same manner for those experiments not including MN, this was calculated from average milk yield and the corresponding milk CP percentage reported for the same SP, which ranged from 4 to 42 d and averaged 9 d [standard deviation (SD) = 9.9]. The large SD corresponds to the difference in the experimental setup between changeover and continuous experiments. It was assumed that milk CP contained on average 95% of protein N, and thus total N was estimated with the following equation: [(milk yield \times protein percentage)/6.38]/0.95 (DePeters and Ferguson, 1992). Milk composition, including fat, protein, and lactose, was provided from each independent experimental data set and determined by infrared spectroscopy.

Samples of plasma (696 samples from 13 experiments) or milk (604 samples from 7 experiments) provided by the SmartCow partners were processed and analyzed for N isotopic signatures at the INRAE laboratory (INRAE, Saint-Genès-Champanelle, France). Similar relationships with NUE were previously reported when analyzing $\Delta^{15}\text{N}$ in either plasma or milk samples (Cantalapiedra-Hijar et al., 2018). Thus, only 1 of the 2 matrices was analyzed in those occasions where samples of plasma and milk were available for a single observation. Because the within-sample repeatability of

Table 1. Description of experiments included in the present meta-analysis study

Experiment	Design	Sampling periods	Dietary treatments	Reference ¹
ID1	Changeover	4	2	Saro et al., 2019
ID2	Changeover	2	2	Herremans et al., 2020
ID3	Continuous	36	3	Reynolds et al., 2021
ID4	Continuous	3	4	Pourazad et al., 2021
ID5	Changeover	4	8	Johansen et al., 2017
ID6	Changeover	4	6	Damborg et al., 2019
ID7	Changeover	4	4	Unpublished data ²
ID8	Changeover	4	6	Giagnoni et al., 2021
ID9	Changeover	4	4	Unpublished data ³
ID10	Changeover	4	4	Martin et al., 2019
ID11	Continuous	4	3	Bayat et al., 2022
ID12	Continuous	13	1	Wallace et al., 2019
ID13	Changeover	2	2	Guyader et al., 2016
ID14	Changeover	2	2	Guyader et al., 2017
ID15	Changeover	4	8	Mendowski et al., 2019
ID16	Changeover	4	4	Mendowski et al., 2020
ID17	Changeover	4	4	Edouard et al., 2018
ID18	Changeover	4	2	Unpublished data ⁴
ID19	Continuous	1	3	Coppa et al., 2020
ID20	Continuous	4	2	Bahloul et al., 2021

¹Details of references are included in the Appendix.

²Brask-Pedersen et al. (Department of Animal Science, Aarhus University, Tjele, Denmark).

³Brask-Pedersen et al. (Department of Animal Science, Aarhus University, Tjele, Denmark).

⁴Edouard et al. (INRAE, Agrocampus-Ouest, PEGASE, Saint-Gilles, France).

isotopic analysis is always greater when using plasma versus milk samples, we decided to prioritize analyzing samples of plasma over milk. Once thawed, milk and plasma samples were vortex-mixed for homogenization, pipetted onto tin capsules, and dried for 24 h at room temperature before analysis. Samples were analyzed for the determination of N isotopic signatures ($\delta^{15}\text{N}$) using an isotope-ratio mass spectrometer (Isoprime Vision; Elementar) coupled to an elemental analyzer (EA Vario Cube; Elementar), with glutamic acid used as the in-house standard. In the same manner, all dried feed ingredients and TMR samples received were analyzed to obtain their $\delta^{15}\text{N}$ values for each dietary treatment and measurement period. For this, subsamples of feed ingredients and TMR were weighed into the tin capsules (between 2 and 4 mg, according to N content). In the case of diets comprising separated ingredients, the average $\delta^{15}\text{N}$ of each ingredient was weighted by the percentage of N the ingredient represents in the diet to obtain a single value of $\delta^{15}\text{N}$ for each diet and period. To ensure reliable $\delta^{15}\text{N}$ determinations, 2 replicates for milk and plasma samples and 3 to 4 replicates for dietary ingredients were analyzed to obtain an average value with $\text{SD} < 0.2\%$. Then, the isotopic discrimination between animal proteins and diet ($\Delta^{15}\text{N}$, ‰) was calculated for each animal as the $\delta^{15}\text{N}$ in animal proteins minus $\delta^{15}\text{N}$ of the corresponding diet.

Statistical Analysis

The primary objective of the present study was to assess the ability of $\Delta^{15}\text{N}$ to predict the between-animal variability in MNE of lactating dairy cows. Therefore, the notion of a contemporary group (CG) is defined here as a set of experimental animals sharing the same DT and SP within a particular experiment (i.e., animals fed the same diet, at the same time and place). According to this definition, an experiment with a 4×4 Latin square design would have 16 CG, unless the period effect was not observed significant, in which case there would be only 4 CG (further explained). Between-animal variability will then be approached in the present study, through different statistical approaches, as the variance within CG, also including the experimental error in addition to the true animal variance. Consequently, when discussing the between-animal variability or relationships between 2 variables at the individual level, we refer to the within-CG level. For experiments containing CG with 3 or fewer observations (10 out of 20), a preliminary adjustment by SP was conducted on MN, NI, $\Delta^{15}\text{N}$, and MNE, according to the methodology described by St-Pierre (2001). For this, data were adjusted by SP using a simple linear model with SP (within experiment) as fixed factor, and

then the obtained residuals were added to the mean value (i.e., intercept) for that experiment. In situations where the period effect was not significant ($P > 0.05$), all animals sharing the same dietary treatment within experiment were considered as a CG. This process allowed us to include those CG with a limited number of observations (e.g., in the case of experiments with an unreplicated Latin square design). Otherwise, it would not have been possible to calculate regressions for those conditions with a low number of observations. For continuous variables, the distribution of values was checked for normality and analyzed for outliers (biologically impossible or unlikely values) using the boxplot function in R software version 3.7.2 (R Development Core Team, 2009). Observations with a residual beyond ± 3 SD were rejected if biological reasons justified their elimination.

Sources of Variation for N Isotopic Discrimination and MNE

Estimates of variance components were evaluated using a random intercept model, through the 'nlme' package (version 3.1-153) using R software with experiment, DT within experiment, and SP within experiment as grouping random factors. In this analysis, the sources of variability for MN, NI, $\Delta^{15}\text{N}$, and MNE were separately analyzed using the following model:

$$Y_{ij} = \beta_0 + \beta_i + e_{ij}, \quad [1]$$

where Y_{ij} is the observed variable (MN, NI, $\Delta^{15}\text{N}$, or MNE) for observation j in group i , β_0 is the mean value for the population, β_i is the random variable representing the deviation for the population mean for the i th group, and e_{ij} is the random variable error for observation j in group i . The residual error of this model represented the within-CG variance and thus included both the between-animal variability and the experimental error.

Repeatability accounts for the contribution of individual animal variability to the total variance not explained by the known experimental factors. In other words, repeatability provides an estimate of the correlation between values from consecutive measurements conducted on the same cow once the known experimental factors (dietary treatment and experimental period within the same experiment) have been accounted for. The repeatability of MN, NI, $\Delta^{15}\text{N}$, and MNE was calculated for each experiment separately with the following equation:

$$\text{Repeatability} = \sigma_{\text{Cow}}^2 / (\sigma_{\text{Cow}}^2 + \sigma_{\text{Residual}}^2), \quad [2]$$

where σ_{Cow}^2 and $\sigma_{\text{Residual}}^2$ are the animal variance (between-animal variability) and experimental error (within-animal variability), respectively. Accordingly, we estimated σ_{Cow}^2 and $\sigma_{\text{Residual}}^2$ for each experiment and variable by including the fixed effects of SP and DT and the random effect of the cow. In each case, the confidence intervals of estimates were checked after fitting the model, to monitor for potential problems in model definition (i.e., abnormally wide intervals; Pinheiro and Bates, 2000).

Analysis of the Relationship Between MNE and N Isotopic Discrimination

Initially, the 'lmList' function of the 'nlme' package (Pinheiro and Bates, 2000) was employed to fit linear regressions relating MNE to $\Delta^{15}\text{N}$ within experiment and within diet and experiment separately. The statistical significance of the response of MNE to $\Delta^{15}\text{N}$ variations was also computed with Pearson correlation coefficients and declared significant at $P \leq 0.05$.

The relationship between MNE and $\Delta^{15}\text{N}$ at the individual animal level was explored following different statistical approaches. In the first approach, the between-animal variability in $\Delta^{15}\text{N}$ was assessed separately along with that of MNE once the random effects of the experiment, SP and DT (i.e., between-CG variability), were removed from the actual values (i.e., MNE and $\Delta^{15}\text{N}$) according to Equation [1], to assess the ability of $\Delta^{15}\text{N}$ to predict between-animal variation. If a relationship between $\Delta^{15}\text{N}$ and MNE was still significant once the between-CG variability was removed from actual values, their residuals, the ability of the biomarker to capture between-animal variation in MNE would be demonstrated (Cantalapiedra-Hijar et al., 2018). In addition, a one-way ANOVA on the $\Delta^{15}\text{N}$ residuals of the 25% highest and 25% lowest cows in terms of MNE within CG was conducted to test on half of the population whether $\Delta^{15}\text{N}$ allowed us to differentiate these 2 contrasting groups of animals.

The second approach involved fitting mixed-effects models (St-Pierre, 2001) using the 'nlme' package in R to test the ability of $\Delta^{15}\text{N}$ to predict MNE variations at 2 levels. For this purpose, 2 tiers of equations were developed: predictions of MNE variations across dietary conditions within experiment using mean dietary values (tier 1) and prediction of the within-CG variability of MNE using individual observations (tier 2). Whereas tier 1 models were tested only at the level of the superior grouping factor (i.e., experiment level), tier 2 models were tested across all grouping factors proposed (i.e., experiment, period within experiment, and CG random effects). The random effects of these

structures were tested on the intercept, the slope, or both. A general positive-definite matrix was employed as variance-covariance structure. These variance-covariance structures obtained from the candidate models were evaluated with the Akaike information criterion to identify the best random effect structure to predict MNE [lowest Akaike information criterion and root mean square prediction error (**RMSPE**)]. Random effect structures were always compared using the restricted maximum likelihood method. The general form of the mixed-effect model was as follows:

$$Y_{ij} = (\beta_0 + b_{0i}) + (\beta_1 + b_{1i}) X_{ij} + e_{ij}, \quad [3]$$

where Y_{ij} is the MNE observed, X_{ij} corresponds to the observed values of $\Delta^{15}\text{N}$, β_0 and β_1 are the fixed effects for the intercept and the slope, respectively; b_i are the random effects of experimental factors; and e_{ij} is the identically distributed within-group error, assumed to be independent of the random effects. The coefficient of determination (R^2) was determined for all candidate models via the 'r.squaredGLMM' function of the R package 'MuMIN' (version 1.43.17). Residuals were checked for homoscedasticity (i.e., the dependent variable exhibits similar variance across the range of values for an independent variable). The models derived from this section were evaluated against the same developmental database (observed vs. predicted). This evaluation focused on evaluating the performance of $\Delta^{15}\text{N}$ to capture the between-animal variation in MNE from the selected models based on their best random effect structure. For this evaluation, the concordance correlation coefficient (**CCC**; Lin, 1989) was used, which was calculated as

$$\text{CCC} = r \times C_b, \quad [4]$$

where r is the Pearson correlation coefficient and C_b is the bias correction factor. The CCC indicates how far the best fit line deviates from the concordance or unity line of the observed values predicted values plot. The CCC ranges from 0 to 1, with greater values indicating better model performance. Although the r value provides a measure of precision, the CCC is indicative of the model accuracy. In addition, the ratio of the RMSPE and SD of observed values (**RSR**) was computed to compare the prediction performance of models.

How the Repeatability of Evaluated Traits Affects Model Fit

The present study tested the hypothesis that better repeatability values of both dependent and independent

variables would enhance model prediction performance. The selected mixed-effects model of $\Delta^{15}\text{N}$ to predict MNE resulting from the mixed-effects meta-analysis in tier 2 was then evaluated for each experiment separately. Then, the coefficients of regression obtained during this model evaluation analysis were regressed on the repeatability values of MNE and of $\Delta^{15}\text{N}$ obtained separately for each experiment (according to equation [2]). These relationships were computed with Pearson correlation coefficients and declared significant at $P \leq 0.05$. If the repeatability of MNE and $\Delta^{15}\text{N}$ values significantly correlated with the model fitting, our hypothesis about the influence of measurement precision on the ability of $\Delta^{15}\text{N}$ to predict MNE was accepted.

RESULTS

Description of the Data Set

Descriptive statistics for animal performances and diet composition are shown in Table 2. Consistency existed in the number of observations across animal performance data; however, fewer records were available for some of the feed chemical composition variables. Only 13% of the data set (165 out of 1,300 observations) were from experiments conducted with cows in early lactation (<50 DIM on average), with the remaining

87% of observations corresponding to the mid- and late lactation stages (DIM ≥ 50). Most of the experiments were conducted using Holstein Friesian cows, and only ID11 and ID12 were conducted using Nordic Red cows. From a total of 490 cows, 71% were multiparous. A wide range of DT ($n = 74$) was included in the initial data set. Corn silage, grass silage, and grass hay were the main forage ingredients used, but they were not present in all diets from all experiments. Feed chemical composition varied widely as a result of the heterogeneity of the experimental diets used in each independent experiment. Crude protein and NDF concentrations, measured in all experimental diets, averaged 157 and 379 g/kg of DM, respectively, and ranged from 110 to 268 and from 202 to 607 g/kg of DM, respectively. Large variation was also observed in ADF content, which ranged from 131 to 351 g/kg of DM. Based on the available information on chemical composition of diets, net energy content for lactation averaged 1.56, with a range of 1.44 to 1.70 Mcal/kg of DM.

Data Editing for Model Development

In the exploratory analysis of the initial data set, it was observed that the relationship between MNE and $\Delta^{15}\text{N}$ in early lactation (DIM <50) was different compared with those observed in mid- and late lac-

Table 2. Descriptions of animal performances and diets from experimental studies

Item	n	Mean	SD	Minimum	Maximum
Production data					
Days in milk	1,300	139	83	4	1,074
DMI, kg/d	1,300	22.4	3.5	12.0	35.4
BW, kg	1,300	663	79	394	1,033
Milk yield, kg/d	1,300	32.8	8.05	3.5	60.1
Milk fat concentration, g/kg	1,300	39.8	5.9	21.1	61.9
Milk protein concentration, g/kg	1,300	33.5	3.6	25.2	50.6
Milk lactose concentration, g/kg	1,219	48.5	2.6	37.4	57.1
Fat:protein ratio	1,300	1.19	0.16	0.70	1.93
Fat yield, kg/d	1,300	1.30	0.35	0.13	2.79
Protein yield, kg/d	1,300	1.09	0.28	0.12	1.96
Lactose yield, kg/d	1,219	1.58	0.40	0.13	3.05
Feed composition data					
Diet composition ¹					
Concentrate, % of DM	74	37.28	15.27	8.79	82.64
Corn silage, % of DM	55	30.77	17.66	4.63	75.76
Grass silage, % of DM	45	42.90	17.17	16.50	70.00
Grass hay, % of DM	28	22.44	8.88	3.03	35.78
Urea, % of DM	6	1.40	0.19	0.78	1.78
Chemical composition ²					
CP, g/kg of DM	74	157	25	110	268
NDF, g/kg of DM	74	379	94	202	607
ADF, g/kg of DM	36	241	66	131	351
Starch, g/kg of DM	52	186	56	92	293
NE _L , Mcal/kg of DM	48	1.56	0.06	1.44	1.70
OM, g/kg of DM	62	861	94	653	955

¹According to diet formulation.

²According to data availability.

tation (Figure 1). For instance, a strong and positive correlation ($r = 0.88$; $P < 0.01$) between $\Delta^{15}\text{N}$ and NUE was observed in one of the experiments (ID20), conducted with high-producing dairy cows during the first 50 d of lactation. Therefore, in the present meta-analysis we decided to restrict the analysis of MNE to mid- and late lactation stages ($\text{DIM} \geq 50$) to improve modeling quality in terms of MNE prediction accuracy by using $\Delta^{15}\text{N}$. This resulted in the exclusion of experiment ID20, dedicated to examining the performance of 8 cows on 2 diets in the peripartum period ($n = 32$ observations), and the removal of some observations corresponding to cows in experiments transiting the declared early lactation period (Figure 2).

Table 3 describes statistics and repeatability values for MN, NI, MNE, and $\Delta^{15}\text{N}$ for the experiments included in this meta-analysis. A large variation in these traits was expected due to the contrasting experimental methods and designs. For instance, NI ranged from 271 (ID4) to 1,152 g of N/d (ID5), and average MN ranged from 19 (ID9) to 291 g of N/d (ID8). As a result, the data set covered a large range of MNE values (from 0.04 to 0.47 g/g) and showed a moderate variability [coefficient of variation (CV) = 13%] in relation to its mean value (0.30 g/g). The difference in ^{15}N natural

abundance between the cow and its diet ($\Delta^{15}\text{N}$) averaged 2.143‰ and ranged from 0.101‰ (ID1) to 4.457‰ (ID8) across diets and experiments.

The repeatability of all traits across experiments varied widely. For instance, repeatability values averaged from 36.1% (for NI in ID14) to 95.3% (for MN in ID14). The high overall repeatability values obtained for MNE (63.0%) was mainly due to the overall high mean repeatability of MN observed across experiments. Removing observations from the early lactation period (column 3 of Table 3) led to greater repeatability estimates for all the variables analyzed when compared with the initial data set (column 2 of Table 3).

Sources of Variation for Nitrogen Partitioning and N Isotopic Discrimination

This analysis showed that the effect of experiment (ID) was the main grouping factor explaining the total variance for all traits included (Table 4). For instance, more than one-third of the variability observed in values for MNE and $\Delta^{15}\text{N}$ was explained by between-experiment variability. Around half of the variability observed in MN was explained by the experiment effect, and the largest source of variation for NI was captured




Figure 1. Relationship between milk N efficiency (MNE, g/g) and N isotopic discrimination ($\Delta^{15}\text{N}$, ‰) in lactating cows using individual values ($n = 1,300$) for early (solid line and closed circles), mid- (dashed line and open circles), and late lactation (dotted line and open triangles). Overall relationships: $\text{MNE}_{\text{EARLY}} = 0.408 - 0.011 \times \Delta^{15}\text{N}$ ($n = 165$; $R^2 = 0.02$; residual standard error (RSE) = 0.08; $P = 0.08$); $\text{MNE}_{\text{MID}} = 0.388 - 0.035 \times \Delta^{15}\text{N}$ ($n = 610$; $R^2 = 0.25$; RSE = 0.04; $P < 0.001$); $\text{MNE}_{\text{LATE}} = 0.374 - 0.039 \times \Delta^{15}\text{N}$ ($n = 525$; $R^2 = 0.28$; RSE = 0.04; $P < 0.001$).




Figure 2. Diagram illustrating the experiment (ID) compilation, data screening, and model development with its evaluation. AIC = Akaike information criterion; RMSPE = root mean square prediction error; CCC = concordance correlation coefficient; RSR = ratio of square root of the mean square prediction error to SD of observed values.

Table 3. Mean, variability, and repeatability values for milk N, N intake, N isotopic discrimination ($\Delta^{15}\text{N}$), and milk N efficiency (MNE) from experimental studies (ID1–19) used in the mixed-effect model analysis

Item	Initial data set	Data set for $\Delta^{15}\text{N}$																			
		ID1	ID2	ID3	ID4	ID5	ID6	ID7	ID8	ID9	ID10	ID11	ID12	ID13	ID14	ID15	ID16	ID17	ID18	ID19	
n	1,300	1,135	52	12	71	167	140	128	138	87	17	32	22	100	16	10	31	32	12	23	45
Milk N, g/d																					
Average	173.4	168.5	175.3	103.5	163	116.1	174.8	203.3	188.5	180	128.3	152.8	213.6	188.4	144.8	125.8	140.9	157.4	153.1	178.4	182.8
SD	42.4	40.1	26.4	7.7	29	17.2	24.2	34.4	38.1	31	54	18.4	21	28.8	27.6	31.8	17.5	17.4	15.6	14.9	30.2
Minimum	19	19	94	93.8	112	70.7	120.7	133.1	46.6	116.6	19	117.8	176.3	115.6	105.3	73.6	105.9	127	122.2	139.5	112.8
Maximum	323.8	291.4	222.1	116.8	240	154.8	233.2	276.3	251.5	291.4	216.2	189.3	254.9	247.8	189.3	198	174.8	183.7	170.2	203.6	230
Repeatability, ¹ %	70	72.4	69.8	79.8	58.6	87.2	82.1	80.1	93.5	81.4	88.8	69.6	68.9	—	—	95.3	79.7	60.3	87.6	71.6	—
N intake, g/d																					
Average	565.8	570.3	489.3	412.5	558.6	423.1	646.6	615	627.1	666.5	533.7	541.5	630.8	615.8	486.6	481.2	490.5	653.5	463.5	499.5	559.9
SD	122.9	123.4	53	18.5	101	59.4	168.5	92.9	81.6	73.7	56.8	49.2	108.7	76.7	86.7	79.7	59.2	39.1	72.3	22.1	74
Minimum	271.1	271.1	397.8	390.6	369	271.1	397.1	412.9	445.4	493.2	442.8	449.5	496.9	464.1	347.3	399.1	410.6	538.6	372.9	457.1	362.1
Maximum	1,151.8	1,151.8	623.5	437	819	689.6	1,151.8	856.4	801.8	826.9	672.1	645.1	851.3	822.3	635	672.5	618.6	734.6	564.1	545.5	724.2
Repeatability, %	67.6	74.8	46.9	87.2	56.1	85.7	76.9	65.7	86.8	85.8	56.5	61.1	68.7	—	—	36.1	80.9	50	91.8	60.1	—
$\Delta^{15}\text{N}$, ‰																					
Sample type ²																					
Average	2.209	2.239	1.104	2.928	1.983	2.041	2.713	1.999	2.112	3.164	2.286	2.023	2.653	2.818	1.558	2.018	2.298	2.775	1.384	1.362	1.497
SD	0.747	0.667	0.502	0.189	0.511	0.31	0.406	0.38	0.55	0.441	0.663	0.397	0.369	0.226	0.382	0.357	0.215	0.276	0.565	0.115	0.691
Minimum	0.038	0.101	0.101	2.626	1.122	1.297	1.82	0.943	0.763	2.208	1.562	1.307	2.079	2.303	1.06	1.503	1.843	2.15	0.584	1.131	0.2
Maximum	4.553	4.457	2.55	3.257	2.941	2.984	3.556	3.055	3.82	4.457	4.048	2.936	3.464	3.364	2.103	2.475	2.654	3.335	2.193	1.544	2.269
Repeatability, %	37.2	56.1	56.8	65.3	59.6	60	40.1	36.5	79.2	55.9	56.1	62.6	69.8	—	—	93.7	75.8	63.7	59.1	41.8	—
MNE, g/g																					
Average	0.31	0.3	0.36	0.25	0.31	0.28	0.28	0.33	0.3	0.27	0.24	0.28	0.36	0.31	0.3	0.26	0.29	0.24	0.34	0.36	0.33
SD	0.06	0.05	0.05	0.01	0.04	0.03	0.05	0.04	0.05	0.04	0.09	0.03	0.03	0.03	0.03	0.04	0.04	0.02	0.05	0.02	0.02
Minimum	0.04	0.04	0.2	0.23	0.24	0.14	0.16	0.23	0.09	0.18	0.04	0.23	0.3	0.21	0.24	0.17	0.24	0.2	0.24	0.32	0.28
Maximum	0.69	0.47	0.47	0.28	0.39	0.36	0.39	0.44	0.38	0.38	0.32	0.35	0.43	0.36	0.35	0.31	0.35	0.27	0.41	0.39	0.39
Repeatability, %	51.7	67.5	69.8	45.4	37.4	84.3	65.7	65.9	89.5	73.1	89.9	52.8	54.4	—	—	45.6	87.9	58.8	37.5	49.7	—
Correlation of MNE with $\Delta^{15}\text{N}$	—	—	-0.55*	-0.40 ^{NS}	-0.81*	-0.48*	-0.47*	-0.41*	-0.61*	-0.29*	-0.87*	-0.45*	-0.50**	-0.33*	-0.24 ^{NS}	-0.43 ^{NS}	-0.76*	-0.07 ^{NS}	-0.84*	-0.48**	0.44*

¹Repeatability values were calculated as $\frac{\sigma_{\text{Cow}}^2}{(\sigma_{\text{Cow}}^2 + \sigma_{\text{Residual}}^2)}$; where σ_{Cow}^2 and $\sigma_{\text{Residual}}^2$ are cow within a single experiment and residual variances, respectively.

²N isotopic signatures ($\delta^{15}\text{N}$) of cows were determined from either milk or plasma samples.

Within-study linear regressions between MNE and $\Delta^{15}\text{N}$ at * $P \leq 0.001$ or ** $P \leq 0.05$; ^{NS} nonsignificant.

Table 4. Variance-component estimates of animal performances, N isotopic discrimination ($\Delta^{15}\text{N}$), and milk N efficiency (MNE) from experimental studies used in the mixed-effect model analysis

Item ¹	Mean value \pm SD	Estimate	95% CI	ICC ² (%)
Milk N, g/d (n = 1,135)	168.5 \pm 40.1			
ID		29.1	20.7–41.0	45.3
SP		0.9	0.1–5.8	1.3
DT		6.8	4.6–10.1	10.6
Residual		27.5	26.3–28.7	42.7
N intake, g/d (n = 1,135)	560.7 \pm 128.4			
ID		75.8	48.7–117.9	34.3
SP		5.8	0.3–98.9	2.6
DT		73.8	60.1–90.5	33.4
Residual		65.6	62.6–68.7	29.7
$\Delta^{15}\text{N}$, ‰ (n = 1,135)	2.239 \pm 0.667			
ID		0.558	0.386–0.806	42.6
SP		0.177	0.150–0.209	13.3
DT		0.286	0.225–0.364	21.8
Residual		0.289	0.276–0.303	22.1
MNE, g/g (n = 1,135)	0.30 \pm 0.05			
ID		0.067	0.039–0.083	41.0
SP		0.019	0.015–0.024	13.7
DT		0.024	0.019–0.032	17.5
Residual		0.039	0.037–0.041	27.8

¹ID = experiment; SP = sampling period within experiment; DT = dietary treatment within period and experiment.

²Intra-class correlation coefficient = total variance explained by the corresponding random variable. For instance, for the nested random variables of DT, it refers to the proportion of variance explained only by the dietary treatment from the total variance.

by the dietary treatment effect. Approximately 20% of variance was captured by the dietary treatment (diets within each sampling period and experiment) in MNE and $\Delta^{15}\text{N}$. In the same manner, the random effect of experimental period further captured around 13% of the variability in $\Delta^{15}\text{N}$ and MNE. Furthermore, the resulting residuals for the variables analyzed were mainly due to between-animal variation and to unidentified random sources of error (within-animal variation).

Relationship Between MNE and N Isotopic Discrimination

The response of MNE to $\Delta^{15}\text{N}$ variation (slope) was negative within experiment for observations from 18 out of 19 experiments (Figure 3b and Table 3), and was different from 0 ($P < 0.05$) for 14 out of 19 experiments. Likewise, although most slopes were negative within diet (67 out of 72 diets; Figure 3c), only slopes for 29 out of 72 diets were different ($P < 0.05$) from 0, likely because the number of observations within diet was rather small (mode = 4 observations per condition). A high variability in the response of MNE to $\Delta^{15}\text{N}$ variation among experiments and dietary treatments was thus evident, suggesting the need for different response (slope) coefficients across experimental conditions in our model.

Relationships Between MNE and N Isotopic Discrimination at the Individual Level

When individual data for MNE and $\Delta^{15}\text{N}$ were independently adjusted by the random effects of experiment, sampling period (within experiment), and dietary treatment (within period and experiment), their residuals were still negatively correlated with each other ($P < 0.001$) with a moderate fit ($R^2 = 0.29$; Figure 4). Moreover, the one-way ANOVA performed to test the ability of $\Delta^{15}\text{N}$ to differentiate the top 25% from the lowest 25% of individuals in terms of MNE within CG was statistically significant ($P < 0.001$; Figure 5), indicating that it is possible to distinguish extreme animals in terms of MNE from their N isotopic signature in a given CG.

Tier 1 and 2 Models. Table 5 presents the mixed-effect regression predictive models of MNE from $\Delta^{15}\text{N}$. These models are displayed according to the data employed for their development: dietary treatment means (tier 1) and individual observations (tier 2). Although the overall slope obtained with the different models of MNE prediction from $\Delta^{15}\text{N}$ were all negative and highly significant ($P < 0.001$), the slope of model 4 was more pronounced and had a slightly lower error than the others (models 2 and 3). Additionally, model 4 had a better modeling fit than the model obtained from the




Figure 3. Relationship between milk N efficiency (MNE) and N isotopic discrimination ($\Delta^{15}\text{N}$) in lactating cows using individual values ($n = 1,135$): (a) Simple linear regression analysis [overall relationship: $\text{MNE} = 0.385 - 0.038 \times \Delta^{15}\text{N}$ [$n = 1,135$; $R^2 = 0.26$; residual standard error (RSE) = 0.04; $P < 0.001$], where open triangles represent multiparous cows and closed circles represent primiparous cows; (b) simple linear regression for each independent study ($n = 19$; within-study regression); (c) simple linear regression analysis for each independent diet ($n = 72$; within-diet regression). In (b) and (c), solid lines represents negative slopes, and dashed lines represents positive slopes. Correlation coefficients (and statistical significances) between MNE and $\Delta^{15}\text{N}$ are presented in Table 3.




Figure 4. Simple linear regression between residuals of milk N efficiency (MNE) in lactating dairy cows and N isotopic discrimination ($\Delta^{15}\text{N}$). CG = contemporary group. Residuals were obtained when variables were independently adjusted for the random effects of the study, period (within study), and diet (within period and study). Equation: $\text{MNE} = -0.067 (\pm 0.003) \times \Delta^{15}\text{N}, \text{‰}$ [$n = 1,135$; $R^2 = 0.29$; residual standard error (RSE) = 0.028; $P < 0.001$].

dietary treatment mean observations (model 1). Based on the Akaike information criterion, the best mixed-effects model for $\Delta^{15}\text{N}$ included the random effects of all known experimental factors defining the CG level (i.e., experiment, sampling period, and dietary treatment) on both the intercept and the slope; that is the most complex model structure (model 4).

Model Evaluation. The overall fit statistics of the selected tier 2 model for $\Delta^{15}\text{N}$ are presented for each experiment in Table 6. In line with the fluctuating overall correlation between MNE and $\Delta^{15}\text{N}$ observed from one experiment to another (Table 3), it was observed that the modeling performance, this time at the CG level, varied widely between experiments. Correlation coefficients (r) between actual and predicted MNE ranged between 0.20 and 0.91, but increases in these correlations did not necessarily result in lower RMSPE in a given experiment. For instance, ID9 had the highest r but also the highest RMSPE. The inclusion of RSR, which includes the SD of the observed MNE at the CG level, allowed us to evaluate the fitness of the selected model on contrasting subsets (experiments).

Analysis of Repeatability for Explaining Variations in MNE Prediction Across Studies

A significant correlation was found between the coefficients of regression obtained during model evaluation analysis (observed vs. predicted correlation coefficient) and the repeatability values of MNE ($R^2 = 0.49$, $P = 0.06$; Figure 6a) or $\Delta^{15}\text{N}$ ($R^2 = 0.54$, $P = 0.03$; Figure 6b) obtained separately for each experiment. In other words, increases in repeatability of either MNE or $\Delta^{15}\text{N}$ enhanced the prediction fitness of the model.

DISCUSSION

The compilation of experiments conducted across 5 countries in Europe and resulting in a data set comprising 1,300 individual observations of MNE in lactating dairy cows allowed us to explore the ability of $\Delta^{15}\text{N}$ as a candidate biomarker to predict the between-animal variability of MNE across a wide range of experimental conditions. In line with previous research, we observed that, on average, $\Delta^{15}\text{N}$ was negatively and

significantly correlated with MNE at the individual level (Cantalapiedra-Hijar et al., 2018), but, in agreement with the recent study by Chen et al. (2020), this association could not be confirmed in early lactation, given the considerable body protein mobilization occurring at this stage. Finally, we identified that higher repeatability estimates for both dependent (MNE) and independent variables ($\Delta^{15}\text{N}$) resulted in models with better prediction fitness.

Associations Between MNE and N Isotopic Discrimination in Periparturient Dairy Cows

In the peripartum, dairy cows often undergo a period of negative energy balance because of the inability to increase energy intake at the same rate at which the energy requirements for milk production increase (de Vries and Veerkamp, 2000; Xu et al., 2018). Body reserves are used (mobilized) to compensate for the resulting energy deficit, and this could alter the estimations of MNE if this phenomenon is not properly accounted for (McNamara et al., 2016; Daniel et al., 2017). Unless body mobilization is adequately measured (Friggens and Newbold, 2007), it is difficult to ascertain how

much of the feed N intake is actually contributing to the total N supply for milk synthesis.

In the same way as MNE measurements, N isotopic signatures are affected by protein mobilization occurring during early lactation. Recently, a study by Chen et al. (2020) tested the ability of milk isotopic signatures (^{15}N and ^{13}C) to predict MNE, energy balance, and milk production of early lactation cows. Our results support their conclusions and suggest that the natural ^{15}N enrichment of animal proteins relative to the diet ($\Delta^{15}\text{N}$) could have some drawbacks and limitations when dairy cows are experiencing net protein mobilization. Indeed, it is well demonstrated from ecophysiological research and human longitudinal studies that protein mobilization and body weight loss may lead to greater ^{15}N enrichment of animal pools relative to diets received (Fuller et al., 2005; Barboza and Parker, 2006). This is because organisms are using their own already ^{15}N -enriched proteins in addition to N from the diet for maintenance or functional purposes. The study by Chen et al. (2020) observed a positive and linear correlation of 0.55 between $\Delta^{15}\text{N}$ and MNE in healthy cows from 4 to 11 wk postpartum. In the present study, a strong and positive correlation between $\Delta^{15}\text{N}$ and




Figure 5. Within-contemporary group (CG) values for N isotopic discrimination ($\Delta^{15}\text{N}$) in the top 25% highest and lowest efficient animals within CG, according to milk N efficiency (MNE). Error bars represent SEM.

Table 5. Mixed-effect regression models of milk N efficiency (MNE, g/g) on the N isotopic discrimination ($\Delta^{15}\text{N}$) using either dietary treatment means or individual observations in mid- and late lactating dairy cows

Item ¹	Model no.	Fit statistics variables ³							
		Intercept	Slope	AIC ²	RMSPE (g/g)	RMSPE (%)	R ²	CCC	RSR
Tier 1: Dietary treatment means									
$\Delta^{15}\text{N}$, ‰ (n = 72)	1	0.378* ± 0.017	-0.037* ± 0.007	-289	0.034	8.9	0.28	0.478	0.822
Experiment random effects									
Tier 2: Individual observations									
$\Delta^{15}\text{N}$, ‰ (n = 1,135)	2	0.403* ± 0.014	-0.049* ± 0.007	-4,313	0.036	8.8	0.30	0.380	0.882
Experiment random effects	3	0.407* ± 0.013	-0.050* ± 0.007	-4,316	0.035	8.7	0.32	0.381	0.883
Experiment and period random effects	4	0.417* ± 0.013	-0.056* ± 0.007	-4,498	0.028	6.8	0.36	0.400	0.851

¹At the treatment means level, the model was tested with random effects on the intercept, and at the individual observation level, all models were tested with random effects on the intercept, slope, or both.

²AIC = Akaike information criterion.

³RMSPE = square root of the mean square prediction error, expressed in g/g and RMSPE% as a percentage of mean observed MNE; R² = coefficient of determination, calculated for equations according to the experimental factor nesting level included in each case; CCC = concordance correlation coefficient; RSR = ratio of square root of the mean square prediction error to SD of observed values.

⁴Best random structure model based on AIC.

*P ≤ 0.001; **P ≤ 0.05; ^{NS}Nonsignificant.

Table 6. Fit statistics of the models obtained¹ to predict milk N efficiency (MNE) from N isotopic discrimination ($\Delta^{15}\text{N}$) at the individual level for each experiment (ID1–19) used in the mixed-effect model analysis

Fit statistics variable ²	Modeling data set	ID1	ID2	ID3	ID4	ID5	ID6	ID7	ID8	ID9	ID10	ID11	ID12	ID13	ID14	ID15	ID16	ID17	ID18	ID19
n	1,135	52	12	71	167	140	128	138	87	17	32	22	100	16	10	31	32	12	23	45
RMSPE, g/g	0.026	0.044	0.011	0.021	0.028	0.019	0.028	0.035	0.032	0.056	0.025	0.023	0.027	0.025	0.026	0.026	0.021	0.026	0.018	0.018
RMSPE, %	8.8	12.2	4.4	6.7	10.2	6.8	8.4	11.5	11.6	23.6	8.8	6.4	8.8	8.3	9.8	8.9	8.7	7.7	4.9	5.6
CCC	0.391	0.433	0.510	0.389	0.340	0.172	0.262	0.483	0.394	0.604	0.453	0.345	0.245	0.322	0.541	0.452	0.227	0.283	0.237	0.339
r	0.52	0.55	0.61	0.43	0.45	0.20	0.40	0.72	0.49	0.91	0.51	0.40	0.33	0.45	0.71	0.80	0.25	0.45	0.43	0.41
RSR	0.857	0.829	0.764	0.922	0.891	1.044	0.914	0.767	0.866	0.653	0.855	0.906	0.945	0.867	0.710	0.762	1.035	0.860	0.893	0.911

¹Selected mixed-effect regression models obtained based on best random structure model based on Akaike information criteria (Table 5).

²Fit statistics variables: n = observations; RMSPE = square root of the mean square prediction error expressed in g/g and RMSPE % as a percentage of mean observed MNE; CCC = concordance correlation coefficient; r = correlation coefficient; RSR = square root of the mean square prediction error to standard deviation of observed values ratio.




Figure 6. Relationship between mixed-effects model of milk N efficiency (MNE) from N isotopic discrimination ($\Delta^{15}\text{N}$; $\text{MNE} = 0.415 - 0.052 \times \Delta^{15}\text{N}$) model evaluation (correlation coefficient between observed vs. predicted MNE) at the within-study level (Table 5) and repeatability of either (a) MNE ($R^2 = 0.49$; $P = 0.06$) or (b) $\Delta^{15}\text{N}$ ($R^2 = 0.54$; $P = 0.03$). Experiments ID12, ID13, and ID19 are not included because they lack repeated measurements.

MNE was observed in one of the experiments (ID20) conducted with high-producing dairy cows in the early lactation stage (Correa-Luna et al., 2021). Moreover,

the coefficient of determination (R^2) between MNE and $\Delta^{15}\text{N}$ in the present study for observations across all experiments in the first 50 d of lactation was lower

($R^2 = 0.02$) and nonsignificant compared with the R^2 obtained from the mid- and late lactation stages (Figure 1). In early lactation, body protein mobilization contributes to alterations in the natural ^{15}N enrichment of milk (or plasma) over the diet, which in turn affects the response in $\Delta^{15}\text{N}$ due to MNE variation.

Mobilization of body reserves has been associated with dairy cow milk production and reproduction performance (Buckley et al., 2003), and with health status (de Vries and Veerkamp, 2000; Xu et al., 2018). The alteration of isotopic signatures due to body reserves mobilization might provide additional evidence toward indirect or proximal detection for health events. More studies, ideally based on larger databases generated from real-world farming conditions, are required to confirm whether $\Delta^{15}\text{N}$ is suitable for these purposes.

N Isotopic Discrimination as a Predictive Biomarker of MNE

In line with the results by Cantalapiedra-Hijar et al. (2018), our analysis confirmed that the most important variance component for MNE (NUE in their study) was between-experiment variation. The contribution of experiment to the variance of MNE in the present study was around a third lower compared with Cantalapiedra-Hijar et al. (2018), probably due to differences between diets and production systems employed in both meta-analyses (dairy cows vs. multiple ruminant systems). Similarly, experiment was the main grouping factor for total variance of $\Delta^{15}\text{N}$, and it was observed that reduced mean MNE was associated with higher mean $\Delta^{15}\text{N}$ (Cantalapiedra-Hijar et al., 2018). Although, on average, a negative association between MNE and $\Delta^{15}\text{N}$ was observed in the present study, the responses were not the same across experiments and diets. The use of mixed-effects models on individual observations allowed the effects of experiment, sampling period, and diet to be removed and, thus, allowed evaluation of the overall association between this biomarker and MNE at the individual level.

Residual standard errors of models obtained in this present study and those reported by Cantalapiedra-Hijar et al. (2018) are comparable and ranged from 0.020 to 0.030 g/g of NUE or MNE, respectively. The differences in slopes obtained between this study and those obtained by Cantalapiedra-Hijar et al. (2018) could be due to the contrasting sets of diets employed. In Cantalapiedra-Hijar and colleagues' study, diets corresponded to different production systems, including beef cattle, dairy goats, and nonlactating sheep, whereas in the present study, diets were only from dairy production systems. Also, the larger intercepts obtained for the models of the present study are probably related

to employing observations from only lactating cows, specifically in mid-late lactation. A meta-analysis to evaluate the ability of MUN to predict MNE at the individual level was conducted by Huhtanen et al. (2015). In their study, the model residual error reported as residuals was in the range of what was obtained in this study and represented a slightly larger RMSPE percentage (8.1% vs. 6.8%). In the same way, the error obtained by Jonker et al. (1998) when using MUN for predicting MNE at the individual level was higher than ours (14.7% vs. 6.8%). Huhtanen et al. (2015) showed that employing MUN was not robust enough as a predictive biomarker of N partitioning at the individual level, and that the systematic addition of animal factors such as milk yield, BW, stage of lactation, dietary CP, and DMI had to be considered to achieve better characterizations of between-animal variability in N partitioning. The lack of response of MUN to predict between-cow variations in MNE could be due to diurnal variations in MUN (Spek et al., 2013), and some of this variation could depend on time of feeding and on milking time with respect to milk sampling (Gustafsson and Palmquist, 1993; Broderick and Clayton, 1997). Another factor of variation in MUN could depend on the method of analysis. A recent study by Portnoy et al. (2021) identified the need to perform regular calibrations for the mid-infrared spectroscopy method, as considerable within- and between-laboratory variation can occur in the reference values for MUN; frequent calibration can therefore improve the precision of a laboratory's determination. Alternatively, mid-infrared spectra of milk have been proposed as a proxy to predict animal variation in MNE in early lactation dairy cows (Grelet et al., 2020), but this methodology could be also conditioned to calibrations to achieve precise determinations, especially when the determination of mid-infrared spectra is performed in different laboratories. Compared with the selected model at the individual level in this study, the RMSPE percentage of the predictive model by Grelet et al. (2020) was more than 2-fold larger (6.8 vs. 17%). The large error obtained by Grelet et al. (2020) was considered not suitable to discriminate between low- and high-NUE cows, and, in this case, this was attributed to the artificially high MNE observed in early lactation related to the severe mobilization of body reserves in this period of the lactation. In this last study, as in that of Huhtanen et al. (2015), the researchers had to include additional parameters such as parity and milk production to reduce the residual error in the predictions. In our case, the significant association between MNE and $\Delta^{15}\text{N}$ across different experiments and dietary treatments confirmed the suitability of this biomarker to significantly discriminate between cows randomly selected from the

same CG if they differ by at least 0.112 g/g of MNE ($\pm 1.96 \times \text{RMSEP}$ at 95% confidence level). At this stage, even though ^{15}N signature in plasma has been proven to be a moderate heritable trait in ruminants (Guarnido-Lopez et al., 2021), the minimum detectable difference of MNE found here (0.112 g/g) is considered too high for use as a tool to assist genetic selection on MNE. Further studies are warranted to confirm this point.

Model Evaluation and Trait Repeatability

Based on different criteria employed to evaluate $\Delta^{15}\text{N}$ as a biomarker of MNE within CG, we observed contrasting performance across experiments (Table 6). The different modeling data set sizes from one experiment to another could have influenced some of these results (Fuentes-Pila et al., 2003). The RSR is a useful tool to compare the performance of models when different data are used. Ideally this indicator should be less than 0.70 for satisfactory prediction models (Moriassi et al., 2007). Moreover, the different prediction fitness between experiments may also be a consequence of the diets employed in each experiment. For instance, Cantalapiedra-Hijar et al. (2016) identified that the association between $\Delta^{15}\text{N}$ and NUE could be compromised when diets are high in rumen-degradable N. If the parallel between NUE and feed efficiency is permitted, Guarnido-Lopez et al. (2021) observed that feed conversion efficiency was poorly correlated with $\Delta^{15}\text{N}$ when employing diets high in fiber relative to diets high in starch, and attributed this to the rumen protein balance. Greater rumen ammonia concentration will increase fractionation of N isotopes at the rumen level (Wattiaux and Reed, 1995). Although beyond the objectives of this present study, mean increases in grass silage at the experiment level resulted in poorer prediction fitness (lower r and higher RSR; data not shown) due to the increased RDP in diets with higher proportions of grass silage relative to more starchy diets (Cantalapiedra-Hijar et al., 2018).

The consistency of a trait or phenotype across time (i.e., repeatability) is of utmost importance for genetic studies (Friggens and Newbold, 2007). For instance, in genetic evaluations, repeatability models based on test-days are used for production traits, to differentiate genetic from phenotypic variance (Berry et al., 2014). In the present study, the mean repeatability estimate for $\Delta^{15}\text{N}$ across experiments was higher than that observed by Wheadon et al. (2014) in growing heifers over a 3-mo period (0.62 vs. 0.56). Across experiments, the mean repeatability estimate for MNE was higher in the present study compared with another study (Ariyaratne et al., 2021) in 2 grazing herds with contrasting farming management in New Zealand. Ariyaratne and col-

leagues observed that the repeatability for efficiency of crude protein utilization (CP in milk divided by CP intake) fluctuated from 0.60 to 0.13 according to the stage of lactation throughout the grazing season, but their mean repeatability was still lower than our mean repeatability estimate for MNE across studies (0.38 vs. 0.65). Although both studies had access to individual records of milk N (often generated from calibrated milk-meters), the observations of the present study were generated in housed conditions with individual records of N intake, and in the study by Ariyaratne et al. (2021), the repeatability was computed based on estimations of N intake on herd level calculated from pasture disappearance, which might have resulted in lower figures (Berry et al., 2014). Moreover, repeatability can also be referred to as the consistency of repeated measurements (Harper, 1994). In other words, a repeatability of 1 indicates that the measurement is perfectly consistent, with no experimental error. The present study confirmed the hypothesis that better repeatability values of both dependent (MNE) and independent variables ($\Delta^{15}\text{N}$) would enhance model prediction performance, as we observed a positive and strong correlation along with the fitness prediction of the selected model (Figure 6a and 6b). This strong association highlights the importance of measurement precision for the identification of proxies for phenotyping animals.

Although in the present study we managed to establish and confirm the negative association of MNE with $\Delta^{15}\text{N}$ over a range of experimental conditions, some potential limitations of the predictive ability for MNE of this biomarker must be highlighted. The fact that in some particular CG the negative association of MNE with $\Delta^{15}\text{N}$ was not observed could be attributable to the uncertainty of reaching a new isotopic equilibrium when animals shifted from one dietary treatment to another, especially for those experiments with a changeover design. Nonetheless, in this study strong correlations were observed in 2 changeover studies (ID9 and ID15), which means that even if the isotopic equilibrium had not reached 100%, the biomarker is still working to predict MNE at the individual level if cows differed by at least 0.112 g/g. It was suggested that the period of transitioning between diets should be no less than 27 d to reach a new isotopic equilibrium, to ensure appropriate analysis of $\Delta^{15}\text{N}$ data (Cantalapiedra-Hijar et al., 2015). Moreover, the determination of the isotopic signatures of diets could also be a limitation. Even though it is more feasible to pipette liquid subsamples (milk or plasma) to a higher level of accuracy and consistency onto the tin capsules, it is difficult to accurately collect minuscule portions of homogeneous dried feed ingredients. Although samples

were ground after drying, those feed ingredients representing a combination of large and small particles, such as silages of pasture or corn, posed a major challenge, considering that the tiny fraction subsampled could substantially change from one portion to another. To avoid this, several repetitions had to be undertaken to reduce the CV, aiming to achieve reliable $\delta^{15}\text{N}_{\text{diet}}$ determinations. Nonetheless, our results show that $\Delta^{15}\text{N}$ is still a powerful biomarker for discriminating within CG a group of extreme cows in terms of MNE (Figure 5). This approach was recently employed to distinguish Brahman steers in terms of feed efficiency from isotopic signatures measured from tail hair and fed low-quality senesced C4 grasses (Costa et al., 2021). In line with our results, the steers with lower $\delta^{15}\text{N}$ had higher feed efficiency, less N excreted in the urine, and higher NUE compared with steers having higher $\delta^{15}\text{N}$. Also, $\delta^{15}\text{N}$ of the 20% highest feed efficiency steers proved to be statistically different from the $\delta^{15}\text{N}$ of the 20% lowest feed efficiency steers, indicating that N isotopic signatures could be used as a tool to identify animals with contrasting NUE. In our case, N isotopic signatures of milk or plasma could not differentiate all cows in terms of MNE in a given CG, but this biomarker permitted us to significantly differentiate the highest from the lowest quartile of lactating cows fed the same diet at the same place and time, in terms of MNE, without the necessity of measuring feed intake. In the context of precision feeding, the implementation of nutritional grouping aims at providing different diets to different groups of animals to better fulfill their nutrient requirements. For instance, N isotopic signatures could be used as a tool to subgroup cows with the highest $\Delta^{15}\text{N}$ values (recognized as of having less MNE) and assigned diets with enzyme or inoculant additives to protect CP from rumen degradation, or fed restricted-CP diets, aiming to increase their MNE and reduce their N excreta. Hence, cluster subgroupings toward more precise feeding without compromising the farm management would improve the overall feed efficiency while reducing the environmental footprint, which should be translated into economic and social benefits (Cabrera and Kalantari, 2016).

CONCLUSIONS

In the present study, we confirmed the negative and significant correlation between $\Delta^{15}\text{N}$ and MNE in lactating dairy cows regardless of experimental site, sampling period, and dietary treatment in mid- and late lactation stages. However, the obtained prediction error of the developed model (0.028 g/g) reveals that $\Delta^{15}\text{N}$ only allows us to differentiate extreme cows in terms of MNE. Hence, $\Delta^{15}\text{N}$ can be implemented as a tool to

group animals (25% highest vs. 25% lowest MNE) for precision feeding. In early lactation, both MNE and $\Delta^{15}\text{N}$ values might be artificially increased because of the considerable protein mobilization of body reserves. This was confirmed by observing a positive (rather than negative) association of $\Delta^{15}\text{N}$ along with MNE in early lactation. Increases in repeatability of either MNE or $\Delta^{15}\text{N}$, or both, improved the prediction fitness of the model to differentiate cows in terms of MNE when fed the same diet at the same time. This emphasizes the need to identify best sampling protocols and to monitor the accuracy of measurements toward the identification and improvement of proxies to phenotype animals.

ACKNOWLEDGMENTS

This work received funding from SmartCow, a project funded by the European Union's Horizon 2020 research and innovation program under grant agreement no. 730924. The authors thank the many people involved in each experiment used in this meta-analysis study. The authors have not stated any conflicts of interest.

REFERENCES

- Ariyaratne, H. B. P. C., M. Correa-Luna, H. T. Blair, D. J. Garrick, and N. López-Villalobos. 2021. Genetic parameters for efficiency of crude protein utilisation and its relationship with production traits across lactations in grazing dairy cows. *N. Z. J. Agric. Res.* 64:62–82. <https://doi.org/10.1080/00288233.2020.1726414>.
- Barboza, P. S., and K. L. Parker. 2006. Body protein stores and isotopic indicators of N balance in female reindeer (*Rangifer tarandus*) during winter. *Physiol. Biochem. Zool.* 79:628–644. <https://doi.org/10.1086/502811>.
- Beatson, P. R., S. Meier, N. G. Cullen, and H. Eding. 2019. Genetic variation in milk urea nitrogen concentration of dairy cattle and its implications for reducing urinary nitrogen excretion. *Animal* 13:2164–2171. <https://doi.org/10.1017/S1751731119000235>.
- Berry, D. P., M. P. Coffey, J. E. Pryce, Y. de Haas, P. Lovendahl, N. Krattenmacher, J. J. Crowley, Z. Wang, D. Spurlock, K. Weigel, K. Macdonald, and R. F. Veerkamp. 2014. International genetic evaluations for feed intake in dairy cattle through the collation of data from multiple sources. *J. Dairy Sci.* 97:3894–3905. <https://doi.org/10.3168/jds.2013-7548>.
- Brito, L. F., N. Bedere, F. Douhard, H. R. Oliveira, M. Arnal, F. Penagaricano, A. P. Schinckel, C. F. Baes, and F. Miglior. 2021. Review: Genetic selection of high-yielding dairy cattle toward sustainable farming systems in a rapidly changing world. *Animal* 15:100292. <https://doi.org/10.1016/j.animal.2021.100292>.
- Broderick, G. A., and M. K. Clayton. 1997. A statistical evaluation of animal and nutritional factors influencing concentrations of milk urea nitrogen. *J. Dairy Sci.* 80:2964–2971.
- Buckley, F., K. O'Sullivan, J. F. Mee, R. D. Evans, and P. Dillon. 2003. Relationships among milk yield, body condition, cow weight, and reproduction in spring-calving Holstein-Friesians. *J. Dairy Sci.* 86:2308–2319. [https://doi.org/10.3168/jds.S0022-0302\(03\)73823-5](https://doi.org/10.3168/jds.S0022-0302(03)73823-5).
- Cabrera, V. E., and A. S. Kalantari. 2016. Economics of production efficiency: Nutritional grouping of the lactating cow. *J. Dairy Sci.* 99:825–841. <https://doi.org/10.3168/jds.2015-9846>.
- Cantalapiedra-Hijar, G., R. J. Dewhurst, L. Cheng, A. R. J. Cabrita, A. J. M. Fonseca, P. Noziere, D. Makowski, H. Fouillet, and I. Ortigues-Marty. 2018. Nitrogen isotopic fractionation as a biomarker

- for nitrogen use efficiency in ruminants: A meta-analysis. *Animal* 12:1827–1837. <https://doi.org/10.1017/S1751731117003391>.
- Cantalapiedra-Hijar, G., H. Fouillet, J. F. Huneau, A. Fanchone, M. Doreau, P. Noziere, and I. Ortigues-Marty. 2016. Relationship between efficiency of nitrogen utilization and isotopic nitrogen fractionation in dairy cows: Contribution of digestion v. metabolism? *Animal* 10:221–229. <https://doi.org/10.1017/S1751731115002025>.
- Cantalapiedra-Hijar, G., I. Ortigues-Marty, B. Sepchat, J. Agabriel, J. F. Huneau, and H. Fouillet. 2015. Diet-animal fractionation of nitrogen stable isotopes reflects the efficiency of nitrogen assimilation in ruminants. *Br. J. Nutr.* 113:1158–1169. <https://doi.org/10.1017/S0007114514004449>.
- Castillo, A. R., E. Kebreab, D. E. Beever, and J. France. 2000. A review of efficiency of nitrogen utilisation in lactating dairy cows and its relationship with environmental pollution. *J. Anim. Feed Sci.* 9:1–32.
- Chen, Y. T., J. P. McNamara, G. L. Ma, J. H. Harrison, and E. Block. 2020. Milk ^{13}C and ^{15}N discriminations as biomarkers of feed efficiency and energy status in early lactation cows. *Anim. Feed Sci. Technol.* 269:114638. <https://doi.org/10.1016/j.anifeedsci.2020.114638>.
- Cheng, L., E. J. Kim, R. J. Merry, and R. J. Dewhurst. 2011. Nitrogen partitioning and isotopic fractionation in dairy cows consuming diets based on a range of contrasting forages. *J. Dairy Sci.* 94:2031–2041. <https://doi.org/10.3168/jds.2010-3763>.
- Cheng, L., A. J. Sheahan, S. J. Gibbs, A. G. Rius, J. K. Kay, S. Meier, G. R. Edwards, R. J. Dewhurst, and J. R. Roche. 2013. Technical note: Nitrogen isotopic fractionation can be used to predict nitrogen-use efficiency in dairy cows fed temperate pasture. *J. Anim. Sci.* 91:5785–5788. <https://doi.org/10.2527/jas.2012-5378>.
- Clark, M. A., N. G. G. Domingo, K. Colgan, S. K. Thakrar, D. Tilman, J. Lynch, I. L. Azevedo, and J. D. Hill. 2020. Global food system emissions could preclude achieving the 1.5° and 2°C climate change targets. *Science* 370:705–708. <https://doi.org/10.1126/science.aba7357>.
- Correa-Luna, M., M. Larsen, C. Chantelauze, L. Bahloul, and G. Cantalapiedra-Hijar. 2021. Predictions of N use efficiency from natural ^{15}N abundance in periparturient dairy cows are impaired by the protein mobilization. *J. Dairy Sci.* 104(Suppl. 1):64. (Abstr.)
- Costa, D. F. A., M. Reis, R. M. Dixon, M. T. Fletcher, and L. F. P. Silva. 2021. Measuring nitrogen use efficiency in cattle from stable isotope ratios in hair. Page 01/2021AAAS in Proc. 33rd Conf. Austral. Assoc. Anim. Scientists, Perth, Australia. North Australia Beef Research Council Publishers.
- Daniel, J. B., N. C. Friggens, H. Van Laar, C. P. Ferris, and D. Sauvant. 2017. A method to estimate cow potential and subsequent responses to energy and protein supply according to stage of lactation. *J. Dairy Sci.* 100:3641–3657. <https://doi.org/10.3168/jds.2016-11938>.
- de Vries, M. J., and R. F. Veerkamp. 2000. Energy balance of dairy cattle in relation to milk production variables and fertility. *J. Dairy Sci.* 83:62–69. [https://doi.org/10.3168/jds.S0022-0302\(00\)74856-9](https://doi.org/10.3168/jds.S0022-0302(00)74856-9).
- Deniro, M. J., and S. Epstein. 1981. Influence of diet on the distribution of nitrogen isotopes in animals. *Geochim. Cosmochim. Acta* 45:341–351. [https://doi.org/10.1016/0016-7037\(81\)90244-1](https://doi.org/10.1016/0016-7037(81)90244-1).
- DePeters, E. J., and J. D. Ferguson. 1992. Nonprotein nitrogen and protein distribution in the milk of cows. *J. Dairy Sci.* 75:3192–3209. [https://doi.org/10.3168/jds.S0022-0302\(92\)78085-0](https://doi.org/10.3168/jds.S0022-0302(92)78085-0).
- Friggens, N. C., and J. R. Newbold. 2007. Towards a biological basis for predicting nutrient partitioning: the dairy cow as an example. *Animal* 1:87–97. <https://doi.org/10.1017/S1751731107657772>.
- Fuentes-Pila, J., M. Ibañez, J. M. De Miguel, and D. K. Beede. 2003. Predicting average feed intake of lactating Holstein cows fed totally mixed rations. *J. Dairy Sci.* 86:309–323. [https://doi.org/10.3168/jds.S0022-0302\(03\)73608-X](https://doi.org/10.3168/jds.S0022-0302(03)73608-X).
- Fuller, B. T., J. L. Fuller, N. E. Sage, D. A. Harris, T. C. O’Connell, and R. E. Hedges. 2005. Nitrogen balance and $\delta^{15}\text{N}$: Why you’re not what you eat during nutritional stress. *Rapid Commun. Mass Spectrom.* 19:2497–2506. <https://doi.org/10.1002/rcm.2090>.
- Grelet, C., E. Froidmont, L. Foldager, M. Salavati, M. Hostens, C. P. Ferris, K. L. Ingvarsen, M. A. Crowe, M. T. Sorensen, J. A. Fernandez Pierna, A. Vanlierde, N. GenglerGplusE Consortium, and F. Dehareng. 2020. Potential of milk mid-infrared spectra to predict nitrogen use efficiency of individual dairy cows in early lactation. *J. Dairy Sci.* 103:4435–4445. <https://doi.org/10.3168/jds.2019-17910>.
- Guarnido-Lopez, P., I. Ortigues-Marty, S. Taussat, C. Fossaert, G. Renand, and G. Cantalapiedra-Hijar. 2021. Plasma proteins $\delta^{15}\text{N}$ vs urea as candidate biomarkers of between-animal variations of feed efficiency in beef cattle contrasting diets: Phenotypic and genetic evaluation. *Animal* 15:100318. <https://doi.org/10.1016/j.animal.2021.100318>.
- Gustafsson, A. H., and D. L. Palmquist. 1993. Diurnal variation of rumen ammonia, serum urea, and milk urea in dairy cows at high and low yields. *J. Dairy Sci.* 76:475–484. [https://doi.org/10.3168/jds.S0022-0302\(93\)77368-3](https://doi.org/10.3168/jds.S0022-0302(93)77368-3).
- Harper, D. G. C. 1994. Some comments on the repeatability of measurements. *Ring. Migr.* 15:84–90. <https://doi.org/10.1080/03078698.1994.9674078>.
- Hellwing, A. L. F., P. Lund, M. R. Weisbjerg, F. W. Oudshoorn, L. Munksgaard, and T. Kristensen. 2015. Comparison of methods for estimating herbage intake in grazing dairy cows. *Livest. Sci.* 176:61–74. <https://doi.org/10.1016/j.livsci.2015.01.013>.
- Huhtanen, P., E. H. Cabezas-Garcia, S. J. Krizsan, and K. J. Shingfield. 2015. Evaluation of between-cow variation in milk urea and rumen ammonia nitrogen concentrations and the association with nitrogen utilization and diet digestibility in lactating cows. *J. Dairy Sci.* 98:3182–3196. <https://doi.org/10.3168/jds.2014-8215>.
- Jonker, J. S., R. A. Kohn, and R. A. Erdman. 1998. Using milk urea nitrogen to predict nitrogen excretion and utilization efficiency in lactating dairy cows. *J. Dairy Sci.* 81:2681–2692.
- Lapierre, H., R. Berthiaume, G. Raggio, M. C. Thivierge, L. Doepel, D. Pacheco, P. Dubreuil, and G. E. Lobley. 2005. The route of absorbed nitrogen into milk protein. *Animal Sci.* (Cambridge) 80:11–22. <https://doi.org/10.1079/ASC41330011>.
- Lin, L. 1989. A concordance correlation coefficient to evaluate reproducibility. *Biometrics* 45:255–268. <https://doi.org/10.2307/2532051>.
- McNamara, J. P., K. Huber, and A. Kenéz. 2016. A dynamic, mechanistic model of metabolism in adipose tissue of lactating dairy cattle. *J. Dairy Sci.* 99:5649–5661. <https://doi.org/10.3168/jds.2015-9585>.
- Moriassi, D. N., J. G. Arnold, M. W. Van Liew, R. L. Bingner, R. D. Harmel, and T. L. Veith. 2007. Model evaluation guidelines for systematic quantification of accuracy in watershed simulations. *Trans. ASABE* 50:885–900. <https://doi.org/10.13031/2013.23153>.
- Nousiainen, J., K. J. Shingfield, and P. Huhtanen. 2004. Evaluation of milk urea nitrogen as a diagnostic of protein feeding. *J. Dairy Sci.* 87:386–398.
- Ocko, I. B., T. Sun, D. Shindell, M. Oppenheimer, A. N. Hristov, S. W. Pacala, D. L. Mauzerall, Y. Xu, and S. P. Hamburg. 2021. Acting rapidly to deploy readily available methane mitigation measures by sector can immediately slow global warming. *Environ. Res. Lett.* 16:054042. <https://doi.org/10.1088/1748-9326/abf9c8>.
- Pinheiro, J., and D. Bates. 2000. *Mixed-Effects Models in S and S-PLUS*. Springer.
- Portnoy, M., C. Coon, and D. M. Barbano. 2021. Infrared milk analyzers: Milk urea nitrogen calibration. *J. Dairy Sci.* 104:7426–7437. <https://doi.org/10.3168/jds.2020-18772>.
- R Development Core Team. 2009. R: A language and environment for statistical computing. R Foundation for Statistical Computing. <http://www.R-project.org/>.
- Scott, P. 2017. Global panel on agriculture and food systems for nutrition: food systems and diets: Facing the challenges of the 21st century. *Food Secur.* 9:653–654. <https://doi.org/10.1007/s12571-017-0678-y>.
- Spek, J. W., J. Dijkstra, G. Van Duinkerken, and A. Bannink. 2013. A review of factors influencing milk urea concentration and its relationship with urinary urea excretion in lactating dairy cattle. *J. Agric. Sci.* 151:407–423. <https://doi.org/10.1017/S0021859612000561>.

- St-Pierre, N. R. 2001. Integrating quantitative findings from multiple studies using mixed model methodology. *J. Dairy Sci.* 84:741–755. [https://doi.org/10.3168/jds.S0022-0302\(01\)74530-4](https://doi.org/10.3168/jds.S0022-0302(01)74530-4).
- Uwizeye, A., I. J. M. de Boer, C. I. Opio, R. P. O. Schulte, A. Falcucci, G. Tempio, F. Teillard, F. Casu, M. Rulli, J. N. Galloway, A. Leip, J. W. Erisman, T. P. Robinson, H. Steinfeld, and P. J. Gerber. 2020. Nitrogen emissions along global livestock supply chains. *Nat. Food* 1:437–446. <https://doi.org/10.1038/s43016-020-0113-y>.
- Visioli, F., and A. Strata. 2014. Milk, dairy products, and their functional effects in humans: A narrative review of recent evidence. *Adv. Nutr.* 5:131–143. <https://doi.org/10.3945/an.113.005025>.
- Wattiaux, M. A., and J. D. Reed. 1995. Fractionation of nitrogen isotopes by mixed ruminal bacteria. *J. Anim. Sci.* 73:257–266. <https://doi.org/10.2527/1995.731257x>.
- Wheaton, N. M., M. McGee, G. R. Edwards, and R. J. Dewhurst. 2014. Plasma nitrogen isotopic fractionation and feed efficiency in growing beef heifers. *Br. J. Nutr.* 111:1705–1711. <https://doi.org/10.1017/S0007114513004078>.
- Xu, W., J. Vervoort, E. Saccenti, R. van Hoesj, B. Kemp, and A. van Knegsel. 2018. Milk metabolomics data reveal the energy balance of individual dairy cows in early lactation. *Sci. Rep.* 8:15828. <https://doi.org/10.1038/s41598-018-34190-4>.
- Guyader, J., M. Eugene, M. Doreau, D. P. Morgavi, C. Gerard, and C. Martin. 2017. Tea saponin reduced methanogenesis in vitro but increased methane yield in lactating dairy cows. *J. Dairy Sci.* 100:1845–1855. <https://doi.org/10.3168/jds.2016-11644>.
- Herremans, S., V. Decruyenaere, G. Cantalapiedra-Hijar, Y. Beckers, and E. Froidmont. 2020. Effects of hydrolyzable tannin-treated grass silage on milk yield and composition, nitrogen partitioning and nitrogen isotopic discrimination in lactating dairy cows. *Animal* 14:771–779. <https://doi.org/10.1017/S175173111900226X>.
- Johansen, M., A. L. F. Hellwing, P. Lund, and M. R. Weisbjerg. 2017. Metabolisable protein supply to lactating dairy cows increased with increasing dry matter concentration in grass-clover silage. *Anim. Feed Sci. Technol.* 227:95–106. <https://doi.org/10.1016/j.anifeeds.2017.02.018>.
- Martin, C., A. Ferlay, M. Benaouda, and M. Eugène. 2019. Effect of starch substitution by rapeseed oil on enteric methane emission and performance in dairy cows fed grass-silage based diets. Page 116 in Proc. 7th GGAA—The Greenhouse Gases and Animal Agriculture Conference, Iguassu, Brazil. Embrapa Southeast Livestock.
- Mendowski, S., P. Chapoutot, G. Chesneau, A. Ferlay, F. Enjalbert, G. Cantalapiedra-Hijar, A. Germain, and P. Noziere. 2019. Effects of replacing soybean meal with raw or extruded blends containing faba bean or lupin seeds on nitrogen metabolism and performance of dairy cows. *J. Dairy Sci.* 102:5130–5147. <https://doi.org/10.3168/jds.2018-15416>.
- Mendowski, S., P. Chapoutot, G. Chesneau, A. Ferlay, F. Enjalbert, G. Cantalapiedra-Hijar, A. Germain, and P. Noziere. 2020. Effects of pretreatment with reducing sugars or an enzymatic cocktail before extrusion of faba bean on nitrogen metabolism and performance of dairy cows. *J. Dairy Sci.* 103:396–409. <https://doi.org/10.3168/jds.2019-17286>.
- Pourazad, P., C. Martin, M. Bouchon, Y. Rochette, T. Aumiller, T. Aubert, and K. Wendler. 2021. Effects of phytogenic feed additives on performance and enteric methane emissions in dairy cattle. Page 616 in Proc. Book of Abstracts of the 72nd Annual Meeting of the European Federation of Animal Science, Davos, Switzerland. Wageningen Academic Publishers.
- Reynolds, C., L. Crompton, D. Roberts, A. Jones, A. del Prado, J. M. Moorby, J. Humphreis, T. Misselbrook, and A. Thomson. 2021. Effects of diet protein concentration on digestion and N balance of dairy cows over 3 lactations. Page 612 in Proc. Book of Abstracts of the 72nd Annual Meeting of the European Federation of Animal Science, Davos, Switzerland. Wageningen Academic Publishers.
- Saro, C., D. Morgavi, Y. Rochette, M. Bouchon, A. Le Morvan, S. Duval, M. Kindermann, and C. Martin. 2019. 3-Nitrooxypropanol blocked postprandial enteric methane emissions from lactating dairy cows. Page 564 in Proc. XIIIth International Symposium on Ruminant Physiology (ISRP), Leipzig, Germany. Cambridge University Press.
- Wallace, R. J., G. Sasson, P. C. Garnsworthy, I. Tapio, E. Gregson, P. E. Bani, P. Huhtanen, A. R. Bayat, F. Strozzi, F. Biscarini, T. J. Snelling, N. Saunders, S. L. Potterton, J. Craigon, A. Minuti, E. Trevisi, M. L. Callegari, F. Piccioli Cappelli, E. H. Cabezas-García, J. Vilkki, C. Pinares-Patino, K. O. Fliegerová, J. Mrázek, H. Sechovcová, J. Kopečný, A. Bonin, F. Boyer, P. Taberlet, F. Kokou, E. Halperin, J. L. Williams, K. J. Shingfield, and I. Mizrahi. 2019. A heritable subset of the core rumen microbiome dictates dairy cow productivity and emissions. *Sci. Adv.* 5:eaav8391. <https://doi.org/10.1126/sciadv.aav8391>.

APPENDIX

References of Experiments Used in the Meta-Analysis.

- Bahloul, L., L. E. Hernández Castellano, H. Lapiere, C. Schwab, and M. Larsen. 2021. Abomasal amino acid infusions in postpartum dairy cows. *J. Dairy Sci.* 104(Suppl. 1):149. (Abstr.)
- Bayat, A. R., A. Razzaghi, M. Sari, P. Kairenius, A. Troscher, E. Trevisi, and J. Vilkki. 2022. The effect of dietary rumen-protected *trans*-10, *cis*-12 conjugated linoleic acid or a milk fat-depressing diet on energy metabolism, inflammation, and oxidative stress of dairy cows in early lactation. *J. Dairy Sci.* 105:3032–3048. <https://doi.org/10.3168/jds.2021-20543>.
- Coppa, M., J. Jurquet, M. Eugene, T. Dechaux, Y. Rochette, J. M. Lamy, A. Ferlay, and C. Martin. 2020. Repeatability and ranking of long-term enteric methane emissions measurement on dairy cows across diets and time using GreenFeed system in farm-conditions. *Methods* 186:59–67. <https://doi.org/10.1016/j.ymeth.2020.11.004>.
- Damborg, V. K., S. K. Jensen, M. Johansen, M. Ambye-Jensen, and M. R. Weisbjerg. 2019. Ensiled pulp from biorefining increased milk production in dairy cows compared with grass-clover silage. *J. Dairy Sci.* 102:8883–8897. <https://doi.org/10.3168/jds.2018-16096>.
- Edouard, N., A. Suzanne, P. Lambertson, B. Rouillé, and P. Faverdin. 2018. Complete nitrogen balance in dairy cows: Does every drop of sweat count? Page 641 in Proc. 10th International Symposium on the Nutrition of Herbivores, Clermont-Ferrand, France. Cambridge University Press.
- Giagnoni, G., P. Lund, J. Sehested, and M. Johansen. 2021. Effect of exogenous dietary phytase and concentrate mixtures based on faba beans, rapeseed meal or soybean meal as main protein source on phytate and total phosphorus excretion in dairy cows. *Anim. Feed Sci. Technol.* 276:114913. <https://doi.org/10.1016/j.anifeeds.2021.114913>.
- Guyader, J., M. Doreau, D. P. Morgavi, C. Gerard, C. Loncke, and C. Martin. 2016. Long-term effect of linseed plus nitrate fed to dairy cows on enteric methane emission and nitrate and nitrite residuals in milk. *Animal* 10:1173–1181. <https://doi.org/10.1017/S1751731115002852>.