



HAL
open science

Can nitrite-free recipes of cured meat products prevent the formation of nitroso-compounds during digestion?

Noémie Petit, Elena Keuleyan, Sylvie Bourillon-Blinet, Laurent Aubry,
Aurélie Promeyrat, Gilles Nassy, Véronique Santé-Lhoutellier

► To cite this version:

Noémie Petit, Elena Keuleyan, Sylvie Bourillon-Blinet, Laurent Aubry, Aurélie Promeyrat, et al.. Can nitrite-free recipes of cured meat products prevent the formation of nitroso-compounds during digestion?. ICFD 2022, May 2022, Cork, Ireland. hal-03662224

HAL Id: hal-03662224

<https://hal.inrae.fr/hal-03662224v1>

Submitted on 9 May 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Can nitrite-free recipes of cured meat products prevent the formation of nitroso-compounds during digestion?

Biochimie et Protéines du Muscle
Unité QuaPA

Noémie Petit¹, Elena Keuleyan¹, Sylvie Bourillon-Blinet¹, Laurent Aubry¹, Aurélie Promeprat², Gilles Nassy², Véronique Santé-Lhoutellier¹

References:

(1) Chazelas et al. (2022). Int. J. Epidemiol. 1-14

¹ INRAE, QuaPA UR 370, BPM, F-63122 Saint-Genès-Champanelle, France

² IFIP – Institut du Porc, La motte au Vicomte, BP 35104, 35561 Le Rheu Cedex, France

INTRODUCTION

The presence of nitrite in cured meat products can lead to the formation of nitroso-compounds (NOCs), some of which have been linked to a higher risk of developing colon cancer¹. The objective was to study the formation mechanism of NOCs in nitrite-free cured meat products during *in vitro* dynamic digestion.

RESULTS

Nitrosylheme

Nitrosylheme was <15% in products without added nitrite (NC) and YE.




Figure 1. Nitrosylheme content was assessed after A) 40 min, B) 120 min, and C) 200 minutes of digestion. NC = negative control, PC = positive control, YE = yeast extract, VS = vegetable stock.

Nitrosamines

Nitrosamines were higher at the beginning of gastric digestion for PC and VS, but decreased during digestion for all samples to reach very low levels.




Figure 2. Nitrosamines level throughout the digestion.

Lipid oxidation

In the intestinal phase of digestion, lipid oxidation increased for all samples regardless of the formulation.




Figure 3. Lipid oxidation level throughout the digestion.

CONCLUSIONS

Nitrite-free recipes can reduce NOCs formation in cured meat products *in vitro*. However, lipid oxidation in nitrite-free products is important in the intestinal phase, and could lead to the development of colorectal cancer. Studies *in vivo* are required to validate the results presented herein, and their impact on colon mucosa.

Nitrite-free recipes:

