



**HAL**  
open science

## Pepsin activity as a function of pH and digestion time under static in vitro conditions

Lea Salelles, Juliane Floury, Steven Le Feunteun

### ► To cite this version:

Lea Salelles, Juliane Floury, Steven Le Feunteun. Pepsin activity as a function of pH and digestion time under static in vitro conditions. 7th International Conference on Food Digestion, May 2022, Cork, Ireland. , 2022. hal-03662664

**HAL Id: hal-03662664**

**<https://hal.inrae.fr/hal-03662664v1>**

Submitted on 9 May 2022

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial - NoDerivatives 4.0 International License



## Background


The activity of pepsin, the gastric protease, is generally considered to be negligible for  $\text{pH} \geq 4$ , based on the results obtained with a few purified globular proteins. The present study aimed at studying the activity of porcine pepsin on egg white proteins and casein micelle micro-aggregates over a broad range of pH (from 1 to 7) for short (3 min) and long (2 h) digestion times.

## Materials & Methods

Two series of static *in vitro* gastric digestions (INFOGEST protocol) were conducted with porcine pepsin at different pH with both caseins and egg white proteins. The first series was used to assess the initial reaction rate of hydrolysis (after 3 min) using the OPA method. The second series was used to monitor the kinetics of protein hydrolysis for 2 h of digestion with a high temporal resolution using the pH-STAT method and converting the results in degree of hydrolysis (DH) thanks to OPA analyses of the end samples.


## Key Results

**After 3 min:** The pH activity profile of pepsin strongly depends on the substrate and can be substantial for  $\text{pH} \geq 4$ .





**Fig. 1.** pH dependence of porcine pepsin activity at short time on different protein substrates (our results and literature data), arbitrary setting 100% at  $\text{pH} = 2$ . Lines are guides for the eyes.

**After 2h:** The extent of casein hydrolysis was constant in a remarkably broad range of pH: from 1 to 5 !



**Fig. 2.**  $\text{DH}_{2h}$  at different pH for egg white (EW) and caseins (CA) (mean  $\pm$  sd over 3 replicates). The dotted line is a guide for eyes. The solid line is a linear regression ( $\text{DH}_{2h} = -2.87 \times \text{pH} + 17.27$ ,  $R^2 = 0.99$ )

**During the course of digestion:** Pepsin hydrolysis profiles can be accurately modelled by a simple power law (with a scaling factor,  $\alpha$ , and a shape parameter,  $\beta$ , that seems largely pH independent and characteristic of the substrate)



**Fig. 3.** DH monitored by pH-STAT during 2 h of static gastric *in vitro* digestions (mean  $\pm$  sd over at least 3 replicates). All lines represent fittings of the power law model:

$$\text{DH}(\%) = \alpha \times \text{time}^\beta$$

**Table 1.**  $R^2$  and estimated parameters ( $\alpha$ ,  $\beta$ ) of the power law model on various foods.

Food	pH	$\alpha$	$\beta$	$R^2$	
Egg white proteins	2	1.72		> 0.99	
	3	1.15	0.41	> 0.99	
	4	0.80		> 0.99	
	5	0.42		0.96	
	6	0.18		0.98	
Caseins	2	2.23		> 0.99	
	3	2.78		> 0.99	
	4	2.88	0.23	0.92	
	5	2.55		0.74	
	6	1.18		0.98	
Native whey proteins <sup>1</sup>	3	0.91	0.30	> 0.99	
	Gelled whey proteins <sup>1</sup>	3	0.37	0.57	> 0.99
		3	0.38	0.45	> 0.99
	Pea based cake <sup>2</sup>	3	0.53	0.39	> 0.99
		3	0.09	0.64	> 0.99
	Pea protein gel <sup>2</sup>	3	0.37	0.48	> 0.99

<sup>1</sup> Mat et al., *Food Chemistry*, 2020, 311

<sup>2</sup> Unpublished data of ours

## Main Conclusions

- Pepsin activity under weakly acidic conditions ( $\text{pH} \geq 4$ ) should not always be neglected, in particular, for milk caseins.
- Pepsin instantaneous activity seems to evolve proportionally to the power of time during static *in vitro* gastric digestion.

- For complementary information:



Salelles L., Flory J. & Le Feunteun S. (2021) *Food Funct.*, 12, 12468