



HAL
open science

Maternal exposure to dietary heme iron elicits sex-related gut microbiota reshape associated with gut barrier and glucose metabolism defects in mice offspring

Anaïs Mazenc, Claire Maslo, Corinne Lencina, Ingrid Ahn, Valérie Alquier-Bacquié, Valérie Bézirard, Nathalie Naud, Cécile Héliès-Toussaint, Françoise Guéraud, Fabrice H.F. Pierre, et al.

► To cite this version:

Anaïs Mazenc, Claire Maslo, Corinne Lencina, Ingrid Ahn, Valérie Alquier-Bacquié, et al.. Maternal exposure to dietary heme iron elicits sex-related gut microbiota reshape associated with gut barrier and glucose metabolism defects in mice offspring. *Virtual Keystone Symposia: The microbiome: From mother to Child.*, Jan 2021, Silverthorne, United States. hal-03669495

HAL Id: hal-03669495

<https://hal.inrae.fr/hal-03669495>

Submitted on 16 May 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Anais Mazenc¹, Claire Maslo¹, Corinne Lencina¹, Ingrid Ahn¹, Valérie Alquier-Bacquié¹, Valérie Bézirard¹, Nathalie Naud¹, Cécile Héliès-Toussaint¹, Françoise Guéraud¹, Fabrice Pierre¹, Vassilia Théodorou¹, Maïwenn Olier¹

¹ Toxalim (Research Centre in Food Toxicology), Université de Toulouse, INRAE, ENVT, INP-Purpan, UPS, Toulouse, France.

Maternal exposure to dietary heme iron elicits sex-related gut microbiota reshape associated with gut barrier and glucose metabolism defects in mice offspring

BACKGROUND:

The heme iron provided by meat products has been positively associated with an increased risk of cancers and diabetes (1,2). Heme-induced colorectal cancer promotion is explained at least in part by its ability to induce luminal peroxidation of dietary polyunsaturated fatty acids, thus leading to the formation of cytotoxic and genotoxic luminal reactive aldehydes in the colon (3). Gut microbiota involvement in this process has been recently shown in adult male rats fed for 21 days with heme-enriched diet (4). Interestingly, *Escherichia coli* strongly covaried with luminal heme-induced reactive aldehydes produced at adulthood.

As *E.coli* is a well-known maternally transmitted bacteria belonging to the first colonizers after birth, our aim was to explore whether a disruption of maternal gut homeostasis induced by heme-enriched diet during the perinatal period may influence (i) neoformation of lipid peroxydation-derived reactive aldehydes (LPRA), (ii) gut ecosystem and (iii) glucose metabolism in offspring.

METHODS:

Female C3H/HeN mice were fed AIN76 diets supplemented with ferric citrate (control) or heme iron during gestation and lactation. Diets were balanced for protein, fat and iron (0.036%). From weaning to 9 weeks of age, offspring was fed the control diet. At weaning, microbiota composition of mother and offspring (16S rDNA gene sequencing, Illumina, Miseq) and LPRA formation in fecal waters and urine (TBARS and DH-NMA respectively) were studied. These analysis were repeated once offspring reached adulthood and continued with an oral glucose tolerance test and *ex-vivo* jejunal permeability assessment (Ussing chambers).

RESULTS:

At the weaning stage, the maternal heme intake affected the fecal bacterial community of both mothers and offsprings. In mothers fed the heme-enriched diet, greater richness associated with lower evenness was observed. These α diversity indexes were lower in all offsprings at this stage due to microbiota immaturity but were not affected by the maternal diet. β -diversity was significantly different between groups according to the generation, gender and maternal iron form intake, although mother to offspring transfert of some heme-altered taxa were noted. This imprinting in reponse to heme-enriched maternal diet persisted in offspring microbiota at adulthood and was

associated with more marked sex-related alterations. Indeed only males developed an impaired phenotype characterized by (i) significant higher levels of LPRA, while luminal concentrations of heme iron remained comparable to those of control offspring, (ii) jejunal paracellular permeability increase and (iii) altered blood glucose regulation following the oral glucose challenge associated with a lower insulin concentration in blood.

CONCLUSION:

This work highlights how maternal exposure to dietary heme iron can markedly trigger gut homeostasis disruption through lipid peroxidation in male offspring, and how the contribution of the maternal environment during the perinatal period influences offspring propensity to develop metabolism disorders. Whether maternal transmitted gut microbiota is sufficient to promote intestinal barrier defects and glucose intolerance in a sex-dependent manner remains to be determined.

- (1) Misra R et al., *Red Meat Consumption (Heme Iron Intake) and Risk for Diabetes and Comorbidities?* *Curr Diab Rep.* 2018;18(11). doi:10.1007/s11892-018-1071-8
- (2) Bastide NM et al., *A central role for heme iron in colon carcinogenesis associated with red meat intake.* *Cancer Res.* 2015;75(5):870-879. doi:10.1158/0008-5472.CAN-14-2554
- (3) Baradat M et al., *4-hydroxy-2(E)-nonenal metabolism differs in Apc^{+/+}Cells and in ApcMin⁺cells: It may explain colon cancer promotion by heme iron.* *Chem Res Toxicol.* 2011;24(11):1984-1993. doi:10.1021/tx2003036
- (4) Martin OCB et al., *Haem iron reshapes colonic luminal environment: Impact on mucosal homeostasis and microbiome through aldehyde formation.* *Microbiome.* 2019;7(1):6-8. doi:10.1186/s40168-019-0685-7