



**HAL**  
open science

## Adaptation of slow- and rapid-growing broilers to alternative diets

Quentin Berger, Elodie Guettier, Séverine Urvoix, Elisabeth Le Bihan-Duval,  
Sandrine Mignon-Grasteau

► **To cite this version:**

Quentin Berger, Elodie Guettier, Séverine Urvoix, Elisabeth Le Bihan-Duval, Sandrine Mignon-Grasteau. Adaptation of slow- and rapid-growing broilers to alternative diets. 26. World Poultry Congress, Aug 2022, Paris, France. hal-03670582

**HAL Id: hal-03670582**

**<https://hal.inrae.fr/hal-03670582>**

Submitted on 17 May 2022

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

**26<sup>th</sup> World's Poultry Congress**  
**Book of abstracts 2021**



**POSTPONEMENT**  
07<sup>th</sup> > 11<sup>th</sup>  
**AUGUST**  
**2022**

PARIS, FRANCE  
PALAIS DES CONGRÈS



MORE INFORMATIONS AT

**[WPCPARIS2022.COM](http://WPCPARIS2022.COM)**



## **Book of abstracts**

**Abstracts submitted in 2019 and accepted in 2020**

**26<sup>th</sup> World's Poultry Congress**

**Organized by**

**French Branch of the World's Poultry Science Association**

The 26<sup>th</sup> World's Poultry Congress - 07-11 August 2022

**Publisher**

French Branch of the World's Poultry Science Association

**Editors**

Dr Michèle TIXIER-BOICHARD, chair of 26<sup>th</sup> WPC

Dr Michel DUCLOS, Chairman of the Scientific Committee of 26<sup>th</sup> WPC

Professional Congress Organizer (PCO) – Colloquium Group (Paris, France)

ID : 321

#### ADAPTATION OF SLOW- AND RAPID-GROWING BROILERS TO ALTERNATIVE DIETS

B. Quentin\*(1), É.Guettier(1), S.Urvoix(1), E.Le bihan-duval(1), S.Mignon-grasteau(1)

(1)INRA, Nouzilly, France

\* *Corresponding author:* quentin.berger@inra.fr

To refine feed efficiency selection of broilers, we developed an electronic feed station allowing a continuous record of body weight and feed intake of animals reared in group on floor. Using this station, we compared slow-growing Label Rouge (LR) and fast-growing standard (S) chickens fed either with a corn-soybean diet (CD) or with an alternative diet (AD) including DDGS, fava beans, sunflower and rapeseed meals. Within a genotype, diets were isoproteic, isoenergetic and fulfilled the need of the animals during the three phase of their growth (S birds : 2850, 2900, 2950 kcal.kg<sup>-1</sup> and 21.5, 20, 18.5 % CP; LR birds: 2750, 2850, 2900 kcal.kg<sup>-1</sup> and 20, 18, 16.5 % CP).

For each genotype, the diet effect was estimated by an ANOVA on daily data of body weight, feed intake and feed efficiency, and on carcass composition and meat quality traits at slaughter (84 and 35 d for LR and S, respectively). The potential effect of the diet on performance homogeneity was checked through an ANOVA on the coefficient of variation including the effects of the period (starter, grower, finisher), the diet and their interaction. Correlations between slaughter traits and daily data were calculated by strain and diet.

LR chickens fed with AD were 4 to 9% heavier and had a 6 to 12% lower FCR during the starter period, but no effect was observed on feed intake nor meat characteristics. For S chickens, AD had no impact on the daily BW and feed efficiency but led to a slightly less acidic (+1.7% for pH), less yellow meat (9.7% for the b\* index) and leaner carcass (-14% for abdominal fat yield). Unexpectedly, groups fed with AD were more homogeneous for all traits than with CD. For example, CV was respectively 20 and 22% lower for FCR in LR and S chickens.

Correlations between daily and slaughter traits differed between genotypes and diets. For example, for the group fed with the alternative diet, cumulated FCR was correlated with breast yield in LR chickens (-0.23), but not in S chickens (-0.13). Moreover, these correlations showed that some traits recorded during the starter period could be used as precocious indicators of slaughter traits. For example, bodyweight during the starter phase was positively correlated with fat yield in LR chickens fed with AD.

These first results indicate that animals can adapt to alternative feedstuffs, even those with a rapid growth rate. A complementary study is in progress on pedigree birds from a heavy line of broilers fed with the alternative diet, in order to suggest new criteria of selection related to feed efficiency and adaptability.