



HAL
open science

Interactions between Salivary Proteins and Dietary Polyphenols: Potential Consequences on Gastrointestinal Digestive Events

Martine Morzel, Francis Canon, Sylvain Guyot

► To cite this version:

Martine Morzel, Francis Canon, Sylvain Guyot. Interactions between Salivary Proteins and Dietary Polyphenols: Potential Consequences on Gastrointestinal Digestive Events. *Journal of Agricultural and Food Chemistry*, 2022, 70 (21), pp.6317-6327. <10.1021/acs.jafc.2c01183>. <hal-03682613>

HAL Id: hal-03682613

<https://hal.inrae.fr/hal-03682613v1>

Submitted on 31 May 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire HAL, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons CC BY-NC-ND 4.0 - Attribution - Non-commercial use - No Derivative Works - International License

Interactions between Salivary Proteins and Dietary Polyphenols: Potential Consequences on Gastrointestinal Digestive Events

Martine Morzel,* Francis Canon, and Sylvain Guyot



Cite This: <https://doi.org/10.1021/acs.jafc.2c01183>



Read Online

ACCESS |

Metrics & More

Article Recommendations

ABSTRACT: The present review documents the current knowledge and hypotheses on how polyphenols–saliva interactions may modulate the bioaccessibility or bioavailability of nutrients and highlights research prospects in the field. After an updated description of the different classes of dietary polyphenols and their modifications by food processing or digestion, an overview of interactions between salivary proteins and polyphenols (with an emphasis on tannins) is provided. *In vitro* studies show that the solubility of salivary protein–tannin complexes in gastric conditions depends on the degree of tannin polymerization, while complexes are partly solubilized by bile salts. Salivary proteins–polyphenols interactions may affect digestive processes. For example, polyphenols can bind to and inhibit salivary amylase, with downstream consequences on starch digestion. Some salivary proteins (PRPs) prevent tannin-induced reduced protein digestibility, probably through binding tannins before they interact with digestive proteases. Salivary proteins may also act as scavenger molecules to limit the intestinal uptake of tannins.

KEYWORDS: *digestion, saliva, polyphenols, tannins, proline-rich proteins, mucins*

INTRODUCTION

The health benefits of dietary polyphenols are tremendously well-documented, for example regarding cardiovascular health,¹ metabolic diseases,² or cognitive decline.³ When it comes to digestion, the impact of polyphenols is described mainly in relation to inhibition of enzymes^{4,5} or modulation of gut microbiota.⁶ Another aspect of polyphenols that is extensively studied is their contribution to the sensory properties of food: bitterness and astringency. For the latter, a key mechanism at the origin of this sensation resides in the ability of polyphenols to aggregate proteins including those forming the mucosal pellicle. Such binding may initiate changes in the friction forces at the surface of the oral mucosa and/or intracellular signaling leading to the release of a neurotransmitter.⁷ This mechanism is modulated by the presence in saliva of proteins presenting a high affinity for tannins. In fact, when polyphenols enter the digestive tract, it is expected that part of them will be in the form of salivary proteins–polyphenols complexes.

In parallel, the impact of oral physiology (mastication and salivation) on digestion is gaining interest. For example, *in vitro* digestion was applied to boli produced by *in vivo* mastication of protein-fortified sponge cakes⁸ or to boli produced by *in vitro* simulated normal or deficient mastication of meat products⁹ or pasta.¹⁰ The impact of saliva on digestion has also been described, focusing on its enzymatic content,¹¹ antioxidant properties,¹² or the effect it may have on emulsions.¹³

The present article aims at bringing these different fields closer together. This narrative review does not intend to summarize the plethora of literature on the effect of polyphenols on digestion but is focused on the polyphenols–saliva interactions and how these may modulate the bioaccessibility

or bioavailability of polyphenols themselves or of other nutrients. For that purpose, the review provides an updated description of the different classes of polyphenols and how they are modified by either food processing or digestion. A second part will deal with salivary proteins, especially their fate during digestion, their interactions with polyphenols, and how the resulting complexes may be affected by digestion. This will provide the necessary background to then consider the impact of such interactions (binding, complexation, and/or precipitation) on digestive processes. Some hypotheses have been tested experimentally and published, and they are reported in this article. We also report articles suggesting an impact of polyphenols–salivary proteins interactions, even when this was not tested specifically. Finally, we will also formulate novel hypotheses based on the current knowledge.

DIFFERENT CLASSES OF DIETARY POLYPHENOLS

Polyphenols are very widespread secondary metabolites mainly found in higher plants and in some brown algae. They constitute a vast family of compounds with very diverse molecular structures. However, the term “polyphenol” is often used inappropriately in the literature, in particular by mistakenly considering as polyphenols compounds having in their structure only a single phenolic nucleus. Thus, a comprehensive definition for polyphenols was recommen-

Received: February 17, 2022

Revised: April 29, 2022

Accepted: May 7, 2022




Figure 1. Three classes of vegetable tannins

ded:¹⁴ The term “polyphenol” should be used to define plant secondary metabolites derived exclusively from the shikimate-derived phenylpropanoid and/or the polyketide pathway(s), featuring more than one phenolic ring and being devoid of any nitrogen-based functional group in their most basic structural expression. Thus, very widespread simple phenolics such as hydroxybenzoic or hydroxycinnamic acids must not be considered as “true polyphenols”.

With regard to this definition and based on biosynthetic pathways, several categories of polyphenols are distinguished. A special mention is made for flavonoids, of which there are more than 8000 different polyphenolic structures which are subdivided into a large series of subclasses (anthocyanins, flavanols, flavonols, flavones, etc.) according to the oxidation state and chemical functionalization scheme of the heterocycle.¹⁵ Stilbenes (for instance, resveratrol) and lignans are two other less widespread polyphenol classes for which numerous papers related to their biological properties have been published.^{16,17}

The other method of polyphenol categorization refers to a specific physicochemical feature of certain polyphenols—their tanning property—that refers to the ability of some polyphenols to associate to proteins. Obviously, this property is of particular importance in the context of the present review focusing on polyphenol interactions with salivary proteins. Historically, the term “vegetable tannin” refers to a plant extract used for the conversion of animal skin into leather. Still today, the generally accepted definition for a “vegetable tannin” is the one first proposed by Theodore White, a chemist in the leather industry,¹⁸ and later completed by Swain and Bates-Smith.¹⁹ Vegetable tannins are water-soluble phenolic compounds having a molecular weight between 500 and 3000 Da and having the ability to precipitate alkaloids, gelatin, and other proteins. Nevertheless, the upper value of molecular

weight (3000 Da) must be now reconsidered since several recent studies have demonstrated that much higher molecular weight tannins are water-soluble and in general have even stronger capacity to precipitate proteins.^{20,21}

Tannins are classified into three categories as depicted in Figure 1. First, condensed tannins (proanthocyanidins) are flavanol dimers, oligomers, and polymers. They are divided into subclasses (procyanidins, prodelfinidins, and profisetinidins) depending on the nature of their flavanol constitutive units ((epi)catechin, (epi)gallocatechin, and fisetinidol, respectively). In some cases, a part of the flavanol units are esterified by gallic acid, thus forming galloylated proanthocyanidins. They are found in many edible plant materials, including fruits (apples, grapes, plums, red fruits, ...), some legumes (beans and lentils), nuts (almonds, hazelnuts, pecans, ...), grains (barley, sorgho, ...), and processed food (fruit juices, cocoa, wine, and ciders).²² Second, hydrolyzable tannins (gallo- and ellagitannins) are mainly gallic or ellagic acid polyesters of a polyol moiety, which is generally glucose. Simply considering ellagitannins, near to 1000 different hydrolyzable tannin structures have been identified.²³ They are less widespread in comparison to condensed tannins; however, they are found in diverse sources, such as strawberries, raspberries, pomegranates, mangos, pecans, walnuts, and wines, in particular those aged in oak barrels.^{14,22} Third, phlorotannins²⁴ are oligomers and polymers composed mainly of phloroglucinol units linked together by biaryl or/and diaryl ether bonds and are essentially found in brown algae.

Food Processing: Oxidation and Structural Changes of Polyphenols. During food processing, a part of native polyphenols is involved in biochemical and chemical reactions, undergoing significant changes in their molecular structure and, thus, generating new molecules that can be described as “secondary polyphenols”.²⁵ Those newly formed products

contribute to the organoleptic and nutritional properties of foods, as do native polyphenols. For instance, they contribute to the color and the flavor of various plant-based beverages such as tea, coffee, juices, ciders, and wines.^{25,26}

Among reactions involving polyphenols, oxidation is the one occurring the most frequently during plant raw material transformation.²⁷ From a chemical point of view, oxidation of polyphenols can be summarized by the ability of the phenolic nucleus to lose an electron or a hydrogen atom leading to the formation of reactive species. As a consequence, oxidation of polyphenols is also clearly the main reaction route which leads to the formation of many “secondary polyphenols” which can be described as “oxidation products”.

The oxidation of catechol or pyrogallol groups present in the structures of phenolic compounds (tannins, flavonoids, hydroxycinnamic and hydroxybenzoic acids, etc.) leads to the formation of very reactive intermediate species such as quinones and semiquinones. Note that depending on the situation, these species can be either chemically generated in low concentrations but over long periods, as for example in the case of wine aging, or rapidly produced in large quantities under enzymatic catalysis by polyphenoloxidases and peroxidases during a precise step of the transformation process. For black tea, for example, this occurs during the bruising and rolling operations,²⁸ while for apple juice or cider, this is observed during the crushing and pressing of the fruits.²⁹ The kinetic aspects of these oxidation steps strongly depend on environmental conditions such as the availability of oxygen, the pH, and the presence of transition metals capable of catalyzing oxidation. Thus, in a weakly acidic environment (which is the case with most plant foods), the autoxidation of phenolic compounds is all the more favored as the pH is close to neutrality.

Once formed, quinones and semiquinones then evolve rapidly according to various reaction pathways: (i) electron transfer reactions generating new quinone or radical species, (ii) intramolecular additions and rearrangements, and (iii) intermolecular additions with other phenolic or nonphenolic compounds. Finally, depending on the plant matrix considered (the nature, concentrations, and proportions of the different categories of polyphenols and other constituents present) and environmental conditions imposed by the transformation processes (pH, availability of oxygen, etc.), this multiplicity of reaction pathways logically leads to the generation of numerous oxidation products exhibiting structures whose molecular architectures are sometimes very different from those of their native phenolic precursors. Although our objective is not here to make an exhaustive review of these different structures of oxidation products, we can, however, cite a few relevant examples. Thus, the theaflavins from black teas with their specific benzotropolone nucleus are generated by oxidative coupling between epicatechin and epigallocatechins.³⁰ Interestingly, B-type procyanidins can be converted into A-type either enzymatically³¹ or chemically³² as a result of an intramolecular oxidative coupling reaction. Intra- and intermolecular autoxidative coupling of condensed tannins in wine was characterized in model solutions showing that intermolecular coupling was highly favored when tannins are more concentrated.³³ Still in the field of wine chemistry, the coupling of anthocyanins with catechins and condensed tannins (proanthocyanidin oligomers) often proceeds through oxidative mechanisms leading to the formation of pyranoan-

thocyanin derivatives that contribute to the color change during red wine aging.³⁴

In the case of apple condensed tannins, it has been shown that the structural modifications generated by oxidation (essentially additional intramolecular linkages) significantly modify some of their physicochemical properties, such as their solubility, their rigidity, and their solvation properties in water–ethanol mixtures.³⁵ Although little knowledge is currently available on this issue, it is expected that these structural modifications also affect the ability of tannins to interact with proteins, including salivary proteins. Recently, in relation to what could be the perception of astringency in juices made from some cider apple varieties, a series of dehydrodimers with original structures resulting from oxidative coupling of caffeoylquinic acid (i.e., chlorogenic acid) were studied for their ability to complex saliva proteins in model solutions.³⁶ Contrarily to what is generally expected in the case of tannin–salivary protein interactions, those dehydrodimers presented a low interaction with proline-rich proteins (PRPs) but revealed a specific interaction with statherins, P–B peptide, and cystatins. Obviously, this specific example does not allow establishment of general rules regarding the impact of oxidation of polyphenols on their ability to interact with salivary proteins, and further studies regarding catechins and condensed and hydrolyzable tannins are needed to better understand the structure–activity relationships that characterize those interactions.

Modification during Digestion. The stability and metabolization of polyphenols in the digestive tract depend on the class to which they belong and also vary mainly according to the gastrointestinal compartment considered. Polyphenols are relatively stable in an *in vitro* simulated gastric environment (acidic pH in the presence of pepsin) while they are much more degraded under the neutral pH conditions of the intestine, in the presence of pancreatin and bile salts, which are favorable to autoxidation reactions.^{37–39} For instance, catechins were significantly degraded under simulated digestion, and some of them, in particular those containing a pyrogallol group, were partly converted into dimers by autoxidative coupling.^{38,40} Interestingly, those dimers were analogues of theasinensins, a class of polyphenols also found in mildly fermented tea.³⁸ Furthermore, regarding glycosylated flavonoids, the small intestine is favorable to some deglycosylation reactions. Indeed, human small intestine epithelial cells produce β -glucosidase enzymes able to hydrolyze quercetin glycosides into free quercetin, which is much more bioavailable.⁴¹

However, considering the numerous studies published in this field, it seems that one of the most important biotransformations of dietary polyphenols occurs as a result of the biological action of the gut microbiota.^{42,43} Thus, flavonols,⁴⁴ catechins,⁴⁵ and proanthocyanidins (i.e., condensed tannins)^{46,47} are largely converted into a series of small molecules corresponding to phenolic acids, with most of them being benzoic, phenylacetic, or phenyl propionic acids and their hydroxylated derivatives. In the case of catechins⁴⁸ and proanthocyanidins,⁴⁹ a series of hydroxylated phenyl valeric acids are also produced. Interestingly, those metabolites are produced even from nonextractable polyphenols.⁴² When they are highly polymerized, procyanidins are poorly metabolized.⁴⁶ Regarding ellagitannins, they are partly metabolized into urolitin,⁵⁰ a specific tricyclic metabolite that exhibits phytoestrogenic activities.⁵¹

■ SALIVARY PROTEINS

Whole saliva is defined as the biological fluid present in the mouth. The reader is directed to excellent reviews for further information on saliva functions and composition.^{11,52,53} One should note, however, that saliva flow and composition are highly variable between subjects and depend on many factors, such as age,^{54,55} diet,^{56,57} medication and/or pathologies,⁵⁸ or sensory stimulation during food intake.^{59,60} In any case, apart from water, saliva is composed mainly of proteins. In fact, it contains thousands of different proteins and peptides. These are either secreted by the salivary glands or originate from the crevicular fluid (an exudate of plasma reaching the oral cavity through the gingival crevices), epithelial cells, oral microbiota, or food remnants. The estimation of the proportions of the different proteins suggests that the most abundant proteins are those secreted by salivary glands (mucins, α -amylase, cystatins, histatins, statherin, and PRPs) together with immunoglobulins and albumin.^{61,62} PRPs are of special interest for this review because of their high affinity for tannins. They represent up to 70% of proteins secreted by the parotid glands and 20–30% of proteins in whole saliva.⁶³ Fragments of PRPs also represent a large part of the adult saliva peptidome⁶⁴ and the vast majority of oligopeptides identified in human infant saliva.⁵⁴ The PRPs family is characterized by a high heterogeneity: the products of six genes are found in saliva as a complex mixture of entire, truncated, phosphorylated, glycosylated, and dimerized proteoforms.⁶⁵

Fate of Salivary Proteins along the Digestive Tract.

Except during the sleeping period, where salivary flow is virtually null,⁶⁶ saliva is constantly secreted and thus swallowed. It is estimated that an adult swallows around 0.6–1 L of saliva per day.^{60,67} There are large variations in the reported protein concentration in whole saliva, which may depend on the assay used or the sampling conditions. Mean values are, for example, 0.6 mg·mL⁻¹⁶⁸ in unstimulated saliva of individuals ($n = 200$) included in an observational cohort or 1.06 mg·mL⁻¹ ($n = 19$),⁶⁹ 1.77 mg·mL⁻¹ ($n = 30$),⁷⁰ or 2.2 mg·mL⁻¹ ($n = 13$)⁷¹ in unstimulated saliva of healthy individuals. In any case, this implies that the total amount of protein originating from saliva would not exceed 2 g per day, well below the daily dietary protein intake.

Like any protein entering the digestive tract, it is expected that salivary proteins will be rapidly degraded by proteolytic enzymes. A study unexpectedly reported that secreted carbonic anhydrase 6 was present in mouse stools, with an apparent molecular weight on electrophoretic gels suggesting that it was intact or at least little degraded,⁷² but such a result would need confirmation. The particular case of proline-rich proteins (PRPs) also deserves some attention. Their unique composition in amino acids renders them particularly resistant to digestive hydrolysis. Indeed, the presence of proline residues after basic amino acids (Lys-Pro and Arg-Pro) precludes their cleavage by trypsin. For example, after a protein-free meal, oligopeptides in the mass range 500–5000 kDa were sought in mini-pig jejunal aspirates: 23 endogenous peptides were identified, among which 20 corresponded to PRP fragments of 8 to 22 amino acids (personal data).

Salivary Proteins Interacting Preferentially with Polyphenols. The presence of proteins which are effective precipitators of tannins has been reported in the saliva of some mammals. These proteins are called tannin-binding salivary proteins (TBSPs) and include mainly PRPs and histatins.

Their presence in saliva is thought to be related to diet, as they are abundant in species consuming high amounts of tannins, such as herbivores, lagomorphs, rodents, and omnivores, while they are absent in the saliva of mammals weakly exposed to tannin.^{73–75} Hypotheses explaining the presence of TBSPs in mammalian saliva postulate that these proteins participate in the maintenance of oral homeostasis, counteract the effect of tannins, and/or trigger the sensation of astringency, which signals a high tannin concentration in food. In rats, PRP secretion is induced by a tannin-rich diet.⁷⁶ This secretion precludes the antinutritional effect of tannins⁷⁶ and improves the consumption of solutions containing tannins,⁷⁷ suggesting that their presence reduces the intensity of unpleasant sensations (bitterness and/or astringency) induced by tannin. In agreement with these observations, PRPs protect the oral mucosal pellicle from aggregation by tannins.⁷⁸ Therefore, rather than inducing astringency, TBSPs can be considered as counterdefenses inherited from natural selection, which modulate astringency intensity by scavenging tannins.

PRPs have a higher capacity to protect proteins from aggregation by tannins than other salivary proteins.⁷⁶ This capacity has been attributed to their intrinsically disordered conformation⁷⁹ and to their tandem repeat sequences composed of proline clusters surrounded by flexible amino acids.⁸⁰ The rigid proline clusters provide stable binding sites to initiate the interaction with tannins,^{81,82} while the flexibility of surrounding amino acids allows structural rearrangement,^{80,83} stabilizing the interaction through the establishment of additional hydrogen bonds between the peptide chain and tannins.⁸⁴ This structural rearrangement may explain the higher affinity of full PRPs compared to proline-rich peptides.⁸⁵ The several tandem repeat sequences also provide several binding sites, allowing several stoichiometries of interaction.⁷⁹ Once soluble noncovalent complexes are formed, bound tannins, which are described as multidentate ligands, may bridge tannin-PRP complexes, leading to the formation of aggregates.⁸¹ A number of at least three tannins per protein is required to form aggregates. PRP–tannin aggregates grow with tannin concentration up to their precipitation.⁸¹ Beside PRPs, histatins have been reported to precipitate tannin more effectively than acidic PRPs (aPRPs) at pH 7.4, while acidic PRPs show a greater ability at pH 3.0. This difference may result from different mechanisms of interactions. Nevertheless, both histatins and aPRPs efficiently protected salivary amylase from tannin inhibition.⁸⁶ Interaction of tannin with histatins has been reported to involve π – π stacking between the ring of aromatic amino acids and the phenolic ring of tannins.⁸⁷ Other works have evidenced that aPRP, histatins, and statherins are more prone to be precipitated by tannins than basic PRP (bPRP) and glycosylated PRP (gPRP).⁸⁸ For bPRP, this effect could be due to Coulombic repulsion between the bPRP–tannin soluble complexes decreasing the formation of aggregates, as bPRPs are highly charged at the acidic pH used in this study. Regarding gPRP, the presence of glycosylation has been reported to increase the solubility of the gPRP–tannin supramolecular edifices compared to nonglycosylated PRPs.⁸⁹ The presence of sugar chains may reduce the aggregation due to steric hindrance. Investigations on the affinity of TBSPs for tannins have revealed that bPRP and gPRP have a higher affinity for tannins than aPRP,⁹⁰ illustrating that a difference should be made between the TBSPs' affinity for tannin and their propensity to be precipitated. The same study highlighted the influence of the

tannin structure on TBSP–tannin affinity, with changes of the ranking of TBSP affinity as a function of tannin structure. Another study also reported a higher affinity of bPRPs for flavan-3-ol than aPRPs.⁹¹

Mucins, which play an important role in the lubrication of the oral cavity, are other salivary proteins prone to interact with tannins. The interaction between the mucin MUC5B and tannins is followed by a structural reorganization of the mucin network⁹² up to the formation of aggregates. MUC5B is the main salivary protein composing the mucosal pellicle, which is a thin layer of salivary proteins anchoring at the surface of the oral cells via interactions with the transmembrane mucin MUC1.⁹³ The aggregation of the mucosal pellicle by tannins leads to an increase of the friction forces at the surface of the oral mucosa.⁷⁸ This phenomenon is thought to be involved in the sensation of astringency and may induce the cleavage of the transmembrane MUC1, which is an external sensor of epithelial cells.⁷

Regarding the impact of the phenolic structure on tannin–protein interactions, the affinity of proanthocyanidins for proteins increases with the number of constitutive units and their level of galloylation.⁹⁴ The presence of a third hydroxyl group on the B-ring has also been reported to increase the affinity of tannins for proteins.⁹⁵ Indeed, the number of hydroxyl groups increases the stability of the tannin–protein interaction,⁸⁴ via the establishment of hydrogen bonds, while their positions are also important.⁹⁶ The establishment of interactions also seems to depend of the polyphenol spatial configuration.⁹⁷ For example, molecules with flavan-3-ol monomers linked via C4–C6 linkages have a higher affinity for proteins than their isomers with C4–C8 linkages.^{94,98} Moreover, *cis*-2,3-flavan-3-ols have a slightly higher affinity for proteins than *trans*-2,3-flavan-3-ols.⁹⁶

Persistence of Salivary Proteins–Polyphenols Complexes along the Digestive Tract. In order to investigate preabsorption events in the gastrointestinal tract, polyphenols are sometimes subjected to *in vitro* digestion, which includes an oral phase.⁹⁹ However, the most widely used and recognized protocols (e.g., ref 100) recommend the use of artificial saliva, which typically contains only amylase. Artificial saliva's protein composition is, therefore, far less complex than true saliva. The question of the resistance of complexes (specifically PRPs–tannins complexes) to digestion has, nevertheless, been addressed using *in vitro* approaches. Thus, insoluble complexes formed from basic PRPs and condensed tannins remained stable in conditions mimicking those in the stomach.¹⁰¹ Similar observations were reported for histatin–tannin aggregates, with most aggregates remaining insoluble under *in vitro* conditions mimicking those of the stomach and of the small intestine without bile salts.¹⁰² For both types of aggregates, the presence of bile salts in intestinal conditions partly solubilized the insoluble complexes. Whether tannins remained in the form of soluble complexes or were released was not investigated. More recently, it was shown that SP–tannins complexes with a low degree of polymerization were disrupted (or at least dissolved) by gastric digestion while complexes made of salivary proteins and tannin tetramers and above were significantly less solubilized, which the authors interpreted as being more resistant to gastric digestion.¹⁰³ Finally, an *in vivo* study showed a 3- to 4-fold increase of fecal excretion of a hydrolyzable tannin (pentagalloyl glucose) in PRPs-secreting rats compared to rats which were not secreting

PRPs. The authors, therefore, proposed that some PRP–tannins complexes may be excreted in feces.¹⁰⁴

■ IMPACT OF PP–SPS INTERACTIONS ON DIGESTIVE PROCESSES: EXPERIMENTAL EVIDENCE AND HYPOTHESES

Digestive Enzymes' Activities. Sugars' Metabolizing Enzymes. Some studies report interactions of polyphenols with salivary amylase. Tea, but not tea depleted in tannins, was shown to reduce amylase activity in saliva,¹⁰⁵ and isolated human salivary amylase was inhibited by a gallotannin.¹⁰⁶ A recent study reported that some flavonoids (ECG and EGCG) bind very rapidly, within seconds, to human salivary amylase.¹⁰⁷ In the earlier articles, the relevance of amylase inhibition was described in relation to the development of caries, but the downstream consequences on digestion were not envisaged. This is probably the case because salivary amylase was traditionally not considered as a direct contributor to starch digestion except within the oral cavity. However, it was evidenced that the kinetics of the gastric pH decrease was compatible with preservation of salivary amylase activity for at least 20 min.¹⁰⁸ In *in vitro* dynamic conditions, 80% of starch was released (under solubilized or hydrolyzed forms) during the first 20 min of gastric digestion in the presence of salivary amylase, vs approximately 10% when saliva was replaced by water. Taken together, these articles suggest that inhibition of salivary amylase by tannins may have an impact on starch digestion in the gastrointestinal tract. Pancreatic amylase or intestinal α -glucosidases are also inhibited by polyphenols or phenolic extracts from various plant species,^{109–111} which is rather considered as an opportunity to control glucose uptake and, therefore, manage type 2 diabetes in humans.¹¹² However, we found no report of the impact of tannin-binding salivary proteins (TBSPs) on the activities of such gastrointestinal enzymes. In contrast, it was shown that the brush border enzyme lactase was inhibited by EGCG but also that binding of EGCG to PRPs could alleviate this inhibition.¹¹³

Lipases. Similarly to sugar metabolizing enzymes, the general trend described is a reduction of pancreatic lipase activity by polyphenols.^{114,115} Gastric lipase has received less attention, but at least one study reports its inhibition by myricitrin-5-methyl ether, a star anise flavonoid.¹¹⁶ Again, the literature does not describe the impact of the formation of salivary proteins–PP complexes on lipase activity.

Proteases. Due to the importance of protein metabolism on muscle mass, much research has focused on the impact of dietary polyphenols (particularly condensed tannins) on protein digestibility in animals.^{117,118} The overall consensus is that diets rich in tannins have a negative impact on weight, but the underlying mechanisms (systemic or digestive) are not fully understood. Focusing on protein digestion, *in vitro* studies on gastric or gastrointestinal proteolysis sometimes differ in their conclusions,^{119–121} even within the same study, depending on the protein substrate.¹²² In fact, tannins may have different consequences on protein digestibility, as summarized by Butler.¹¹⁷ On one hand, tannins may denature dietary protein substrates and, thus, facilitate the action of proteases. Contrarily, binding of tannins to protein substrates may hinder accessibility of proteases to the cleavage sites. Finally, and more generally for all hydrolytic enzymes, binding of tannins to the enzymes themselves could either modify the tridimensional structure of their catalytic sites (allosteric inhibition) or preclude access to the catalytic site by steric hindrance

(competitive inhibition) and, thus, directly inhibit their activity. For example, phenolic extracts from various plant species can drastically inhibit the activity of isolated digestive enzymes, for example, the proteolytic enzymes pepsin, trypsin, and chymotrypsin.^{123,124} The activity of the brush border enzyme aminopeptidase N is also inhibited by some flavonoids of grape seeds.⁹⁹ In this context, salivary proteins (particularly PRPs) have long been considered as major actors in the protection against the detrimental impact of tannins through binding of those molecules before they can interact with proteolytic digestive enzymes. This hypothesis is partly supported by animal studies: in rodents, a tannin-rich diet induces weight loss attributed to reduced food digestibility, which is continuous in hamsters but reversible after 3 days in rats and mice. In parallel, feeding on tannins induces in rats and mice a dramatic increase in salivary proline-rich proteins (PRPs) within 3 days,⁷⁶ while this is not observed in hamsters.¹²⁵ Another study on rats demonstrated that diet supplementation in propanolol, which can suppress PRP production in saliva, aggravated the adverse effect of tannins on growth.¹²⁶ However, this study and others¹⁰⁴ suggested that the reduced apparent nitrogen digestibility induced by dietary tannins, and calculated using the nitrogen content in feces, may not be due to reduced digestibility of dietary protein but rather to increased nitrogen excretion related to higher secretion of PRPs. Shimada⁷⁵ even proposed that tannins should not be regarded as inhibitors of digestion but rather as toxins with systemic impacts on growth and metabolism. In fact, it is plausible that both effects exist, depending on the type and structure of tannins.

Oxidation of Nutrients. It is admitted that the gastric environment is suitable for oxidation of dietary nutrients, particularly lipid peroxidation. It was demonstrated that, on average (pooling salivas of 7 volunteers), saliva could limit lipid peroxidation of turkey meat in simulated gastric fluid and that this effect was greatly enhanced in the presence of red wine polyphenols.¹² Unfortunately, the impact of polyphenols alone was not evaluated in this study, so it is unknown whether the effect is of an additive or synergetic type. Depending on their structure, polyphenols may chelate Fe^{2+} ions, increasing their oxidation to Fe^{3+} ions in the presence of oxygen. This mechanism decreases the quantity of Fe^{2+} that can participate in the Fenton reaction and, thus, precludes the production of hydroxyl radicals.¹²⁷ This antioxidant activity of polyphenols may be limited by the presence of TBSP, which may compete with Fe^{2+} for the binding to polyphenols.¹²⁸ However, and in contrast to this hypothesis, a study evaluated the oxidant scavenging activities (OSAs) of various polyphenols and polyphenol-containing beverages and noted a very large intensification of the OSA in the presence of human saliva.¹²⁹ The authors suggested that salivary proteins help solubilize lipophilic polyphenols and, thereby, increase their availability. Whether this translates into limitation of oxidation in the gastric compartment is not described.

Physical Modification of the Gastric and Intestinal Mucus. The structure of the mucus layer, particularly in the small intestine, can modulate the rate of nutrient absorption. For example, dietary fibers (alginate) can combine with the intestinal mucus and delay the transport of lipid digestion products.¹³⁰ Various polyphenols can interact with oral or gastrointestinal mucins. For example, the oral mucins MUC5B and MUC7 aggregate in the presence of EGCG.⁹² Binding of EGCG to purified porcine gastric mucin was characterized and

indicated a multilayer binding process.¹³¹ Cross-linking of the GI purified mucins MUC5A and MUC2 by tea-derived galloylated polyphenols was also demonstrated.¹³² Using a cell-based model of oral epithelium lined by a mucosal pellicle,⁹³ it was shown that the presence of bPRPs greatly moderated the impact of EGCG on the formation of MUC5B aggregates.⁷⁸ In contrast, such a formal demonstration of the impact of salivary proteins in the gastrointestinal tract has to our knowledge not been performed. Nonetheless, some observations are in line with this hypothesis. For example, in pigs fed a diet rich in shredded acorns, the cardiac gland region of the stomach became dark-brownish, and fluorescent microscopy revealed that tannins bonded to the gastric mucin, but no tissue damage was observed. This relative innocuity of acorn tannins was accompanied by an accumulation of endogenous proline in the stomach. Although the origin of this proline was not precisely determined, the authors highlighted the possibility that it may correspond to salivary PRPs.¹³³ Therefore, PP–SP interactions would not prevent binding to gastric mucins but would limit the disruption of the mucin layer. If similar protective phenomena occur toward intestinal mucins, this would most likely preserve the physical structure of the mucus layer and, in turn, the diffusion of nutrients and the rates of nutrient absorption.

Transport of Polyphenols across the Intestinal Epithelial Barrier. Some studies have focused on the transport of polyphenols across the intestinal epithelium, in most cases using cell culture methods. However, studies often use polyphenols in their original form and not as complexes with dietary or salivary proteins. An interesting study investigated whether a basic PRP (IB4) modulated such transport in Caco-2 cells.¹³⁴ It was found that the presence of IB4 reduced the transport of a hydrolyzable tannin (pentagalloyl glucose) in both the apical to basolateral and basolateral to apical directions and that this was concomitant with the formation of insoluble tannins–IB4 complexes. The study was further extended to another salivary protein (histatin 5) and to two polyphenols.¹³⁵ The presence of histatin 5 considerably reduced the transport of pentagalloyl glucose across cells but hardly affected the transport of quercetin. The authors, therefore, concluded that some salivary proteins may act as scavenger molecules as far as in the intestine, to limit the uptake of tannins. Another article documented that the cytotoxic effect of EGCG was higher on intestinal HT29 (non-mucin-secreting) than on HT29-MTX-E12 (mucin-secreting) intestinal cells and overall much higher than the cytotoxicity of EC, which has lower ability to bind proteins. The results suggested a binding of EGCG to the mucus which prevented absorption of the noxious polyphenol. The presence of β -caseins further reduced the EGCG cytotoxicity, and the authors pointed out that β -casein may be considered as a simple model to replicate the effect of salivary PRPs due to common characteristics such as richness in proline residues.¹³⁶ Therefore, by analogy and as in the previously quoted articles, PRP–EGCG interactions would represent a protective mechanism to limit tannin bioavailability.

Modifications of the Chyme Physical Properties. One of the functions of saliva is to produce a food bolus that is lubricated and cohesive and, thus, safe to swallow. Saliva can also induce in-mouth emulsification of oil or fat¹³⁷ or flocculation of emulsions.¹³⁸ A recent study documented the impact of proteins modeling salivary constituents (α -amylase, mucin, or a combination of both) on the particle size of peanut




Figure 2. Schematic view of some consequences of salivary proteins–tannins complexation along the digestive tract. Created with BioRender.com.

olesome emulsions after *in vitro* oral and gastrointestinal digestion. Salivary constituents had no impact on the particle size distribution after the model oral phase but induced flocculation in gastric conditions, particularly in the presence of mucins.¹³ Given the high affinity of polyphenols for mucins as described above, it would be of interest to determine whether this effect is modified by the presence of polyphenols.

RESEARCH PROSPECTS

This rapid review of the literature highlighted that research on the consequences of polyphenols on digestive processes is active, and so is work on polyphenols–salivary proteins interactions. The impact of saliva on digestive processes is also an emerging field. However, only a limited number of studies have combined those topics. The few experimental works on the interactions salivary proteins–polyphenols (and even more specifically salivary proteins–tannins) all point at beneficial consequences on digestive events: such interactions would protect against the inhibition of digestive enzymes (amylase, lactase, and proteolytic enzymes), would limit the alteration of the gastric and intestinal mucosal integrity, and would reduce intestinal absorption of potentially toxic tannins (Figure 2).

Other potential consequences regarding oxidation of nutrients or the physical properties of the chyme in the gastric compartment (with, in turn, impact the digestibility or digestion kinetics) are only hypothetical and deserve further attention.

In any case, the studies focusing on the matter should carefully consider the ratio of the different interactors; that is, what is the proportion of polyphenols or tannins bound by salivary proteins, and how do these saliva proteins compete with proteins of the food matrix or with other host-derived proteins (in particular mucins of the gastrointestinal lining)? To our knowledge such data are not available in conditions

mimicking digestion of a mixed meal containing polyphenols. One should also consider that salivary proteins–tannins complexes formed in the oral cavity will be modified downstream during digestion. Studies on absorptive events (in particular, those using cellular models) could, therefore, take advantage of the now well-recognized *in vitro* digestion models to prepare the materials that may be applied to cells. It is thanks to those improvements that the combined effects of polyphenol and salivation on digestion will be adequately characterized. Such results would be of interest when formulating nutritional recommendations on tannin-containing foods, particularly for specific populations affected by hyposalivation due to pathologies, medical treatments, or aging.

AUTHOR INFORMATION

Corresponding Author

Martine Morzel – INRAE, Institut Agro, UMR STLO, F-35042 Rennes, France; orcid.org/0000-0002-3589-3641; Phone: +33 223 48 56 17; Email: martine.morzel@inrae.fr; Fax: +33 223 48 53 50

Authors

Francis Canon – Centre des Sciences du Goût et de l'Alimentation, UMR 1324 INRAE, UMR 6265 CNRS, Université de Bourgogne Franche-Comté, F-21000 Dijon, France; orcid.org/0000-0002-6140-5007

Sylvain Guyot – INRAE, UR BIA, F-35653 Le Rheu, France

Complete contact information is available at: <https://pubs.acs.org/10.1021/acs.jafc.2c01183>

Notes

The authors declare no competing financial interest.

REFERENCES

- (1) Khurana, S.; Venkataraman, K.; Hollingsworth, A.; Piche, M.; Tai, T. Polyphenols: Benefits to the Cardiovascular System in Health and in Aging. *Nutrients* **2013**, *5* (10), 3779–3827.
- (2) Burton-Freeman, B.; Brzeziński, M.; Park, E.; Sandhu, A.; Xiao, D.; Edirisinghe, I. A Selective Role of Dietary Anthocyanins and Flavan-3-Ols in Reducing the Risk of Type 2 Diabetes Mellitus: A Review of Recent Evidence. *Nutrients* **2019**, *11* (4), 841.
- (3) Baba, Y.; Inagaki, S.; Nakagawa, S.; Kaneko, T.; Kobayashi, M.; Takihara, T. Effect of Daily Intake of Green Tea Catechins on Cognitive Function in Middle-Aged and Older Subjects: A Randomized, Placebo-Controlled Study. *Molecules* **2020**, *25* (18), 4265.
- (4) Mcdougall, G. J.; Stewart, D. The Inhibitory Effects of Berry Polyphenols on Digestive Enzymes. *BioFactors* **2005**, *23* (4), 189–195.
- (5) Sun, L.; Wang, Y.; Miao, M. Inhibition of α -Amylase by Polyphenolic Compounds: Substrate Digestion, Binding Interactions and Nutritional Intervention. *Trends Food Sci. Technol.* **2020**, *104*, 190–207.
- (6) Duarte, L.; Gasaly, N.; Poblete-Aro, C.; Uribe, D.; Echeverria, F.; Gotteland, M.; Garcia-Diaz, D. F. Polyphenols and Their Anti-Obesity Role Mediated by the Gut Microbiota: A Comprehensive Review. *Rev. Endocr. Metab. Disord.* **2021**, *22* (2), 367–388.
- (7) Canon, F.; Belloir, C.; Bourillot, E.; Brignot, H.; Briand, L.; Feron, G.; Lesniewska, E.; Nivet, C.; Septier, C.; Schwartz, M.; Tournier, C.; Vargiolu, R.; Wang, M.; Zahouani, H.; Neiers, F. Perspectives on Astringency Sensation: An Alternative Hypothesis on the Molecular Origin of Astringency. *J. Agric. Food Chem.* **2021**, *69* (13), 3822–3826.
- (8) Assad-Bustillos, M.; Palier, J.; Rabesona, H.; Choiset, Y.; Della Valle, G.; Feron, G. Role of the Bolus Degree of Structure on the Protein Digestibility during *in Vitro* Digestion of a Pea Protein-fortified Sponge Cake Chewed by Elderly. *J. Texture Stud.* **2019**, *51*, 134–143.
- (9) Peyron, M.-A.; Sayd, T.; Sicard, J.; Mirade, P.-S.; Pinguet, J.; Chambon, C.; Santé-Lhoutellier, V. Deciphering the Protein Digestion of Meat Products for the Elderly by *in Vitro* Food Oral Processing and Gastric Dynamic Digestion, Peptidome Analysis and Modeling. *Food Funct.* **2021**, *12*, 7283.
- (10) Blanquet-Diot, S.; François, O.; Denis, S.; Hennequin, M.; Peyron, M.-A. Importance of Oral Phase in *in Vitro* Starch Digestibility Related to Wholegrain versus Refined Pastas and Mastication Impairment. *Food Hydrocoll.* **2021**, *112*, 106277.
- (11) Pedersen, A.; Sørensen, C.; Proctor, G.; Carpenter, G. Salivary Functions in Mastication, Taste and Textural Perception, Swallowing and Initial Digestion. *Oral Dis.* **2018**, *24* (8), 1399–1416.
- (12) Gorelik, S.; Kohen, R.; Ligumsky, M.; Kanner, J. Saliva Plays a Dual Role in Oxidation Process in Stomach Medium. *Arch. Biochem. Biophys.* **2007**, *458* (2), 236–243.
- (13) Wang, Q.; Gao, C.; Yang, N.; Nishinari, K. Effect of Simulated Saliva Components on the *in Vitro* Digestion of Peanut Oil Body Emulsion. *RSC Adv.* **2021**, *11* (49), 30520–30531.
- (14) Quideau, S.; Deffieux, D.; Douat-Casassus, C.; Pouysegou, L. Plant Polyphenols: Chemical Properties, Biological Activities, and Synthesis. *Angew. Chem.-Int. Ed.* **2011**, *50* (3), 586–621.
- (15) Andersen, O.; Markham, K. *Flavonoids, Chemistry, Biochemistry and Applications*; CRC Press: Boca Raton, FL, 2006.
- (16) Landete, J. Plant and Mammalian Lignans: A Review of Source, Intake, Metabolism, Intestinal Bacteria and Health. *FOOD Res. Int.* **2012**, *46* (1), 410–424.
- (17) Meng, X.; Zhou, J.; Zhao, C.; Gan, R.; Li, H. Health Benefits and Molecular Mechanisms of Resveratrol: A Narrative Review. *FOODS* **2020**, *9* (3), 340.
- (18) White, T. *The Chemistry of Vegetable Tannins - a Symposium*; Society of Leather Trade Chemists: Croydon, 1956; pp 7–29.
- (19) Swain, T.; Bate-Smith, E. C. *Flavonoids Compounds. Comparative Biochemistry*; Academic Press: New York, 1962; Vol. III, pp 755–809.
- (20) Girard, A.; Bean, S.; Tilley, M.; Adrianos, S.; Awika, J. Interaction Mechanisms of Condensed Tannins (Proanthocyanidins) with Wheat Gluten Proteins. *FOOD Chem.* **2018**, *245*, 1154–1162.
- (21) Leppä, M.; Laitila, J.; Salminen, J. Distribution of Protein Precipitation Capacity within Variable Proanthocyanidin Fingerprints. *Molecules* **2020**, *25* (21), 5002.
- (22) Smeriglio, A.; Barreca, D.; Bellocchio, E.; Trombetta, D. Proanthocyanidins and Hydrolysable Tannins: Occurrence, Dietary Intake and Pharmacological Effects. *Br. J. Pharmacol.* **2017**, *174* (11), 1244–1262.
- (23) Quideau, S. *Chemistry and Biology of Ellagitannins—An Underestimated Class of Bioactive Plant Polyphenols*; Quideau, S., Ed.; World Scientific: Singapore, 2009.
- (24) Meng, W.; Mu, T.; Sun, H.; Garcia-Vaquero, M. Phlorotannins: A Review of Extraction Methods, Structural Characteristics, Bioactivities, Bioavailability, and Future Trends. *Algal Res.-Biomass Biofuels Bioprod.* **2021**, *60*, 102484.
- (25) Tanaka, T.; Matsuo, Y.; Kouno, I. Chemistry of Secondary Polyphenols Produced during Processing of Tea and Selected Foods. *Int. J. Mol. Sci.* **2010**, *11* (1), 14–40.
- (26) Cheynier, V. Polyphenols in Foods Are More Complex than Often Thought. *Am. J. Clin. Nutr.* **2005**, *81* (1), 223S–229S.
- (27) Liu, X.; Le Bourvellec, C.; Guyot, S.; Renard, C. Reactivity of Flavanols: Their Fate in Physical Food Processing and Recent Advances in Their Analysis by Depolymerization. *Compr. Rev. FOOD Sci. FOOD Saf.* **2021**, *20* (5), 4841–4880.
- (28) Subramanian, N.; Venkatesh, P.; Ganguli, S.; Sinker, V. Role of Polyphenol Oxidase and Peroxidase in the Generation of Black Tea Theaflavins. *J. Agric. FOOD Chem.* **1999**, *47* (7), 2571–2578.
- (29) Nicolas, J. J.; Richard-Forget, F. C.; Goupy, P. M.; Amiot, M.; Aubert, S. Y. Enzymatic Browning Reactions in Apple and Apple Products. *Crit. Rev. Food Sci. Nutr.* **1994**, *34* (2), 109–157.
- (30) Takemoto, M.; Takemoto, H. Synthesis of Theaflavins and Their Functions. *Molecules* **2018**, *23* (4), 918.
- (31) Poupard, P.; Sanoner, P.; Baron, A.; Renard, C. M. G. C.; Guyot, S. Characterization of Procyanidin B2 Oxidation Products in an Apple Juice Model Solution and Confirmation of Their Presence in Apple Juice by High-Performance Liquid Chromatography Coupled to Electrospray Ion Trap Mass Spectrometry: LC-ESI-MS/MS of Procyanidin B2 Oxidation Products. *J. Mass Spectrom.* **2011**, *46* (11), 1186–1197.
- (32) Kondo, K.; Kurihara, M.; Fukuhara, K.; Tanaka, T.; Suzuki, T.; Miyata, N.; Toyoda, M. Conversion of Procyanidin B-Type (Catechin Dimer) to A-Type: Evidence for Abstraction of C-2 Hydrogen in Catechin during Radical Oxidation. *TETRAHEDRON Lett.* **2000**, *41* (4), 485–488.
- (33) Vernhet, A.; Carrillo, S.; Poncet-Legrand, C. Condensed Tannin Changes Induced by Autooxidation: Effect of the Initial Degree of Polymerization and Concentration. *J. Agric. FOOD Chem.* **2014**, *62* (31), 7833–7842.
- (34) Rentzsch, M.; Schwarz, M.; Winterhalter, P. Pyranoanthocyanins - an Overview on Structures, Occurrence, and Pathways of Formation. *Trends Food Sci. Technol.* **2007**, *18* (10), 526–534.
- (35) Zanchi, D.; Konarev, P. V.; Tribet, C.; Baron, A.; Svergun, D. I.; Guyot, S. Rigidity, Conformation, and Solvation of Native and Oxidized Tannin Macromolecules in Water-Ethanol Solution. *J. Chem. Phys.* **2009**, *130* (24), 245103.
- (36) Castillo-Fraire, C. M.; Brandão, E.; Poupard, P.; Le Quére, J.-M.; Salas, E.; de Freitas, V.; Guyot, S.; Soares, S. Interactions between Polyphenol Oxidation Products and Salivary Proteins: Specific Affinity of CQA Dehydrodimers with Cystatins and P-B Peptide. *Food Chem.* **2021**, *343*, 128496.
- (37) Bermudez-Soto, M.; Tomas-Barberan, F.; Garcia-Conesa, M. Stability of Polyphenols in Chokeberry (*Aronia Melanocarpa*) Subjected to *in Vitro* Gastric and Pancreatic Digestion. *Food Chem.* **2007**, *102* (3), 865–874.
- (38) Neilson, A.; Hopf, A.; Cooper, B.; Pereira, M.; Bomser, J.; Ferruzzi, M. Catechin Degradation with Concurrent Formation of

- Homo- and Heterocatechin Dimers during in Vitro Digestion. *J. Agric. FOOD Chem.* **2007**, *55* (22), 8941–8949.
- (39) Thomas-Valdes, S.; Theoduloz, C.; Jimenez-Aspee, F.; Schmeda-Hirschmann, G. Effect of Simulated Gastrointestinal Digestion on Polyphenols and Bioactivity of the Native Chilean Red Strawberry (*Fragaria Chilensis* Ssp. *Chilensis* f. *Patagonica*). *Food Res. Int.* **2019**, *123*, 106–114.
- (40) Martini, S.; Conte, A.; Tagliazucchi, D. Bioaccessibility, Bioactivity and Cell Metabolism of Dark Chocolate Phenolic Compounds after in Vitro Gastro-Intestinal Digestion. *J. Funct. FOODS* **2018**, *49*, 424–436.
- (41) Nemeth, K.; Plumb, G.; Berrin, J.; Juge, N.; Jacob, R.; Naim, H.; Williamson, G.; Swallow, D.; Kroon, P. Deglycosylation by Small Intestinal Epithelial Cell Beta-Glucosidases Is a Critical Step in the Absorption and Metabolism of Dietary Flavonoid Glycosides in Humans. *Eur. J. Nutr.* **2003**, *42* (1), 29–42.
- (42) Gonzalez-Sarrias, A.; Espin, J.; Tomas-Barberan, F. Non-Extractable Polyphenols Produce Gut Microbiota Metabolites That Persist in Circulation and Show Anti-Inflammatory and Free Radical-Scavenging Effects. *TRENDS FOOD Sci. Technol.* **2017**, *69*, 281–288.
- (43) Selma, M.; Espin, J.; Tomas-Barberan, F. Interaction between Phenolics and Gut Microbiota: Role in Human Health. *J. Agric. FOOD Chem.* **2009**, *57* (15), 6485–6501.
- (44) Serra, A.; Macia, A.; Romero, M.; Reguant, J.; Ortega, N.; Motilva, M. Metabolic Pathways of the Colonic Metabolism of Flavonoids (Flavonols, Flavones and Flavanones) and Phenolic Acids. *FOOD Chem.* **2012**, *130* (2), 383–393.
- (45) Clifford, M.; van der Hooft, J.; Crozier, A. Human Studies on the Absorption, Distribution, Metabolism, and Excretion of Tea Polyphenols. *Am. J. Clin. Nutr.* **2013**, *98* (6), 1619S–1630S.
- (46) Bazzocco, S.; Mattila, I.; Guyot, S.; Renard, C.; Aura, A. Factors Affecting the Conversion of Apple Polyphenols to Phenolic Acids and Fruit Matrix to Short-Chain Fatty Acids by Human Faecal Microbiota in Vitro. *Eur. J. Nutr.* **2008**, *47* (8), 442–452.
- (47) Deprez, S.; Brezillon, C.; Rabot, S.; Philippe, C.; Mila, I.; Lapiere, C.; Scalbert, A. Polymeric Proanthocyanidins Are Catabolized by Human Colonic Microflora into Low-Molecular-Weight Phenolic Acids. *J. Nutr.* **2000**, *130* (11), 2733–2738.
- (48) Takagaki, A.; Nanjo, F. Catabolism of (+)-Catechin and (–)-Epicatechin by Rat Intestinal Microbiota. *J. Agric. FOOD Chem.* **2013**, *61* (20), 4927–4935.
- (49) Le Bourvellec, C.; Boas, P.; Lepercq, P.; Comtet-Marre, S.; Auffret, P.; Ruiz, P.; Bott, R.; Renard, C.; Dufour, C.; Chatel, J.; Mosoni, P. Procyanidin-Cell Wall Interactions within Apple Matrices Decrease the Metabolization of Procyanidins by the Human Gut Microbiota and the Anti-Inflammatory Effect of the Resulting Microbial Metabolome In Vitro. *Nutrients* **2019**, *11* (3), DOI: 10.3390/nu11030664.
- (50) Cerda, B.; Periago, P.; Espin, J.; Tomas-Barberan, F. Identification of Urolithin A as a Metabolite Produced by Human Colon Microflora from Ellagic Acid and Related Compounds. *J. Agric. FOOD Chem.* **2005**, *53* (14), 5571–5576.
- (51) Larrosa, M.; Gonzalez-Sarrias, A.; Garcia-Conesa, M.; Tomas-Barberan, F.; Espin, J. Urolithins, Ellagic Acid-Derived Metabolites Produced by Human Colonic Microflora, Exhibit Estrogenic and Antiestrogenic Activities. *J. Agric. FOOD Chem.* **2006**, *54* (5), 1611–1620.
- (52) Humphrey, S. P.; Williamson, R. T. A Review of Saliva: Normal Composition, Flow, and Function. *J. Prosthet. Dent.* **2001**, *85* (2), 162–169.
- (53) Dawes, C.; Pedersen, A. M. L.; Villa, A.; Ekström, J.; Proctor, G. B.; Vissink, A.; Aframian, D.; McGowan, R.; Aliko, A.; Narayana, N.; Sia, Y. W.; Joshi, R. K.; Jensen, S. B.; Kerr, A. R.; Wolff, A. The Functions of Human Saliva: A Review Sponsored by the World Workshop on Oral Medicine VI. *Arch. Oral Biol.* **2015**, *60* (6), 863–874.
- (54) Morzel, M.; Jeannin, A.; Lucchi, G.; Truntzer, C.; Pecqueur, D.; Nicklaus, S.; Chambon, C.; Ducoroy, P. Human Infant Saliva Peptidome Is Modified with Age and Diet Transition. *J. Proteomics* **2012**, *75* (12), 3665–3673.
- (55) Vandenberghe-Descamps, M.; Labouré, H.; Prot, A.; Septier, C.; Tournier, C.; Feron, G.; Sulmont-Rossé, C. Salivary Flow Decreases in Healthy Elderly People Independently of Dental Status and Drug Intake. *J. Texture Stud.* **2016**, *47* (4), 353–360.
- (56) Dawes, C. Effects of Diet on Salivary Secretion and Composition. *J. Dent. Res.* **1970**, *49* (6), 1263–1272.
- (57) Chambon, C.; Neyraud, E.; Sayd, T.; Bros, P.; Di Biagio, R.; Hyvrier, F.; Féart, C.; André, P.; Rodriguez-Artalejo, F.; Lopez-Garcia, E.; Garcia-Esquinas, E.; Gomez-Cabrero, D.; Proctor, G.; Morzel, M. The Salivary Proteome Reflects Some Traits of Dietary Habits in Diabetic and Non-Diabetic Older Adults. *Eur. J. Nutr.* **2021**, *60* (8), 4331–4344.
- (58) Proctor, G. B.; Shaalan, A. M. Disease-Induced Changes in Salivary Gland Function and the Composition of Saliva. *J. Dent. Res.* **2021**, *100* (11), 1201–1209.
- (59) Neyraud, E.; Heinzlerling, C. I.; Bult, J. H. F.; Mesmin, C.; Dransfield, E. Effects of Different Tastants on Parotid Saliva Flow and Composition. *Chemosens. Percept.* **2009**, *2* (2), 108–116.
- (60) Watanabe, S.; Dawes, C. The Effects of Different Foods and Concentrations of Citric Acid on the Flow Rate of Whole Saliva in Man. *Arch. Oral Biol.* **1988**, *33* (1), 1–5.
- (61) Messana, I.; Inzitari, R.; Fanali, C.; Cabras, T.; Castagnola, M. Facts and Artifacts in Proteomics of Body Fluids. What Proteomics of Saliva Is Telling Us? *J. Sep. Sci.* **2008**, *31* (11), 1948–1963.
- (62) Schenkels, L. C. P. M.; Veerman, E. C. I.; Nieuw Amerongen, A. V. Biochemical Composition of Human Saliva in Relation To Other Mucosal Fluids. *Crit. Rev. Oral Biol. Med.* **1995**, *6* (2), 161–175.
- (63) Bennick, A. Salivary Proline-Rich Proteins. *Mol. Cell. Biochem.* **1982**, *45* (2), DOI: 10.1007/BF00223503.
- (64) Vitorino, R.; Barros, A.; Caseiro, A.; Domingues, P.; Duarte, J.; Amado, F. Towards Defining the Whole Salivary Peptidome. *PROTEOMICS - Clin. Appl.* **2009**, *3* (5), 528–540.
- (65) Manconi, B.; Castagnola, M.; Cabras, T.; Olanas, A.; Vitali, A.; Desiderio, C.; Sanna, M. T.; Messana, I. The Intriguing Heterogeneity of Human Salivary Proline-Rich Proteins. *J. Proteomics* **2016**, *134*, 47–56.
- (66) Dawes, C. Salivary Flow Patterns and the Health of Hard and Soft Oral Tissues. *J. Am. Dent. Assoc.* **2008**, *139*, 18S–24S.
- (67) Richardson, C. T.; Feldman, M. Salivary Response to Food in Humans and Its Effect on Gastric Acid Secretion. *Am. J. Physiol.-Gastrointest. Liver Physiol.* **1986**, *250* (1), G85–G91.
- (68) Méjean, C.; Morzel, M.; Neyraud, E.; Issanchou, S.; Martin, C.; Bozonnet, S.; Urbano, C.; Schlich, P.; Hercberg, S.; Péneau, S.; Feron, G. Salivary Composition Is Associated with Liking and Usual Nutrient Intake. *PLoS One* **2015**, *10* (9), No. e0137473.
- (69) Henskens, Y. M. C.; van der Velden, V.; Veerman, E. C. I.; Nieuw Amerongen, A. V. Protein, Albumin and Cystatin Concentrations in Saliva of Healthy Subjects and of Patients with Gingivitis or Periodontitis. *J. Periodontol. Res.* **1993**, *28*, 43–48.
- (70) Chiba, F. Y.; Sumida, D. H.; Moimaz, S. A. S.; Chaves Neto, A. H.; Nakamune, A. C. M. S.; Garbin, A. J. L.; Garbin, C. A. S. Periodontal Condition, Changes in Salivary Biochemical Parameters, and Oral Health-related Quality of Life in Patients with Anorexia and Bulimia Nervosa. *J. Periodontol.* **2019**, *90* (12), 1423–1430.
- (71) Neyraud, E.; Tremblay-Franco, M.; Gregoire, S.; Berdeaux, O.; Canlet, C. Relationships between the Metabolome and the Fatty Acid Composition of Human Saliva; Effects of Stimulation. *Metabolomics* **2013**, *9* (1), 213–222.
- (72) Aleksiewicz, M. B.; Kjeldal, H. Ø.; Klenø, T. G. Identification of Stool Proteins in C57BL/6J Mice by Two-Dimensional Gel Electrophoresis and MALDI-TOF Mass Spectrometry. *Biomarkers* **2005**, *10* (1), 29–40.
- (73) Clauss, M.; Gehrke, J.; Hatt, J.-M.; Dierenfeld, E. S.; Flach, E. J.; Hermes, R.; Castell, J.; Streich, W. J.; Fickel, J. Tannin-Binding Salivary Proteins in Three Captive Rhinoceros Species. *Comp. Biochem. Physiol. A. Mol. Integr. Physiol.* **2005**, *140* (1), 67–72.

- (74) McArthur, C.; Sanson, G. D.; Beal, A. M. Salivary Proline-Rich Proteins in Mammals: Roles in Oral Homeostasis and Counteracting Dietary Tannin. *J. Chem. Ecol.* **1995**, *21* (6), 663–691.
- (75) Shimada, T. Salivary Proteins as a Defense Against Dietary Tannins. *J. Chem. Ecol.* **2006**, *32*, 1149–1163.
- (76) Mehansho, H.; Hagerman, A.; Clements, S.; Butler, L.; Rogler, J.; Carlson, D. M. Modulation of Proline-Rich Protein Biosynthesis in Rat Parotid Glands by Sorghums with High Tannin Levels. *Proc. Natl. Acad. Sci. U. S. A.* **1983**, *80* (13), 3948–3952.
- (77) Glendinning, J. I. Effect of Salivary Proline-Rich Proteins on Ingestive Responses to Tannic Acid in Mice. *Chem. Senses* **1992**, *17* (1), 1–12.
- (78) Ployon, S.; Morzel, M.; Belloir, C.; Bonnotte, A.; Bourillot, E.; Briand, L.; Lesniewska, E.; Lherminier, J.; Aybeke, E.; Canon, F. Mechanisms of Astringency: Structural Alteration of the Oral Mucosal Pellicle by Dietary Tannins and Protective Effect of BPRPs. *Food Chem.* **2018**, *253*, 79–87.
- (79) Canon, F.; Paté, F.; Meudec, E.; Marlin, T.; Cheynier, V.; Giuliani, A.; Sarni-Manchado, P. Characterization, Stoichiometry, and Stability of Salivary Protein-Tannin Complexes by ESI-MS and ESI-MS/MS. *Anal. Bioanal. Chem.* **2009**, *395* (8), 2535–2545.
- (80) Canon, F.; Ployon, S.; Mazauric, J.-P.; Sarni-Manchado, P.; Réfrégiers, M.; Giuliani, A.; Cheynier, V. Binding Site of Different Tannins on a Human Salivary Proline-Rich Protein Evidenced by Dissociative Photoionization Tandem Mass Spectrometry. *Tetrahedron* **2015**, *71* (20), 3039–3044.
- (81) Canon, F.; Paté, F.; Cheynier, V.; Sarni-Manchado, P.; Giuliani, A.; Pérez, J.; Durand, D.; Li, J.; Cabane, B. Aggregation of the Salivary Proline-Rich Protein IB5 in the Presence of the Tannin EgCG. *Langmuir* **2013**, *29* (6), 1926–1937.
- (82) Luck, G.; Liao, H.; Murray, N. J.; Grimmer, H. R.; Warminski, E. E.; Williamson, M. P.; Lilley, T. H.; Haslam, E. Polyphenols, Astringency and Proline-Rich Proteins. *Phytochemistry* **1994**, *37* (2), 357–371.
- (83) Canon, F.; Ballivian, R.; Chirof, F.; Antoine, R.; Sarni-Manchado, P.; Lemoine, J.; Dugourd, P. Folding of a Salivary Intrinsically Disordered Protein upon Binding to Tannins. *J. Am. Chem. Soc.* **2011**, *133* (20), 7847–7852.
- (84) Canon, F.; Giuliani, A.; Paté, F.; Sarni-Manchado, P. Ability of a Salivary Intrinsically Unstructured Protein to Bind Different Tannin Targets Revealed by Mass Spectrometry. *Anal. Bioanal. Chem.* **2010**, *398* (2), 815–822.
- (85) Charlton, A. J.; Baxter, N. J.; Lilley, T. H.; Haslam, E.; McDonald, C. J.; Williamson, M. P. Tannin Interactions with a Full-Length Human Salivary Proline-Rich Protein Display a Stronger Affinity than with Single Proline-Rich Repeats. *FEBS Lett.* **1996**, *382* (3), 289–292.
- (86) Yan, Q.; Bennick, A. Identification of Histatins as Tannin-Binding Proteins in Human Saliva. *Biochem. J.* **1995**, *311* (1), 341–347.
- (87) Shraberg, J.; Rick, S.; Rannulu, N.; Cole, R. A Study of Procyanidin Binding to Histatin 5 Using Electrospray Ionization Tandem Mass Spectrometry (ESI-MS/MS) and Molecular Simulations. *Phys. Chem. Chem. Phys.* **2015**, *17* (18), 12247–12258.
- (88) Soares, S.; Vitorino, R.; Osório, H.; Fernandes, A.; Venâncio, A.; Mateus, N.; Amado, F.; de Freitas, V. Reactivity of Human Salivary Proteins Families Toward Food Polyphenols. *J. Agric. Food Chem.* **2011**, *59* (10), 5535–5547.
- (89) Sarni-Manchado, P.; Canals-Bosch, J.-M.; Mazerolles, G.; Cheynier, V. Influence of the Glycosylation of Human Salivary Proline-Rich Proteins on Their Interactions with Condensed Tannins. *J. Agric. Food Chem.* **2008**, *56* (20), 9563–9569.
- (90) Soares, S.; Mateus, N.; de Freitas, V. Interaction of Different Classes of Salivary Proteins with Food Tannins. *Food Res. Int.* **2012**, *49* (2), 807–813.
- (91) Ramos-Pineda, A. M.; García-Estévez, I.; Soares, S.; de Freitas, V.; Dueñas, M.; Escribano-Bailón, M. T. Synergistic Effect of Mixture of Two Proline-Rich-Protein Salivary Families (APRP and BPRP) on the Interaction with Wine Flavanols. *Food Chem.* **2019**, *272*, 210–215.
- (92) Davies, H. S.; Pudney, P. D. A.; Georgiades, P.; Waigh, T. A.; Hodson, N. W.; Ridley, C. E.; Blanch, E. W.; Thornton, D. J. Reorganisation of the Salivary Mucin Network by Dietary Components: Insights from Green Tea Polyphenols. *PLoS One* **2014**, *9* (9), e108372.
- (93) Ployon, S.; Belloir, C.; Bonnotte, A.; Lherminier, J.; Canon, F.; Morzel, M. The Membrane-Associated MUC1 Improves Adhesion of Salivary MUC5B on Buccal Cells. Application to Development of an in Vitro Cellular Model of Oral Epithelium. *Arch. Oral Biol.* **2016**, *61*, 149–155.
- (94) Ricardo-da-Silva, J.; Cheynier, V.; Souquet, J.; Moutounet, M.; Cabanis, J.; Bourzeix, M. Interaction of Grape Seed Procyanidins with Various Proteins in Relation to Wine Fining. *J. Sci. Food Agric.* **1991**, *57* (1), 111–125.
- (95) Siebert, K. J. Effects of Protein-Polyphenol Interactions on Beverage Haze, Stabilization, and Analysis. *J. Agric. Food Chem.* **1999**, *47* (2), 353–362.
- (96) Bacon, J. R.; Rhodes, M. J. C. Development of a Competition Assay for the Evaluation of the Binding of Human Parotid Salivary Proteins to Dietary Complex Phenols and Tannins Using a Peroxidase-Labeled Tannin. *J. Agric. Food Chem.* **1998**, *46* (12), 5083–5088.
- (97) McManus, J.; Davis, K.; Beart, J.; Gaffney, S.; Lilley, T.; Haslam, E. Polyphenol Interactions. 1. Introduction - Some Observations on the Reversible Complexation of Polyphenols with Proteins and Polysaccharides. *J. Chem. Soc.-Perkin Trans. 2* **1985**, *9*, 1429–1438.
- (98) Ezaki-Furuichi, E.; Nonaka, G.; Nishioka, I.; Hayashi, K. Affinity of Procyanidins (Condensed Tannins) from the Bark of *Rhaphiolepis Umbellata* for Proteins. *Agric. Biol. Chem.* **1987**, *51* (1), 115–120.
- (99) Laurent, C.; Besançon, P.; Caporiccio, B. Flavonoids from a Grape Seed Extract Interact with Digestive Secretions and Intestinal Cells as Assessed in an in Vitro Digestion/Caco-2 Cell Culture Model. *Food Chem.* **2007**, *100* (4), 1704–1712.
- (100) Brodtkorb, A.; Egger, L.; Alminger, M.; Alvito, P.; Assunção, R.; Ballance, S.; Bohn, T.; Bourliew-Lacanal, C.; Boutrou, R.; Carrière, F.; Clemente, A.; Corredig, M.; Dupont, D.; Dufour, C.; Edwards, C.; Golding, M.; Karakaya, S.; Kirkhus, B.; Le Feunteun, S.; Lesmes, U.; Macierzanka, A.; Mackie, A. R.; Martins, C.; Marze, S.; McClements, D. J.; Ménard, O.; Minekus, M.; Portmann, R.; Santos, C. N.; Souchon, I.; Singh, R. P.; Vegarud, G. E.; Wickham, M. S. J.; Weitschies, W.; Recio, I. INFOGEST Static in Vitro Simulation of Gastrointestinal Food Digestion. *Nat. Protoc.* **2019**, *14*, 991.
- (101) Lu, Y.; Bennick, A. Interaction of Tannin with Human Salivary Proline-Rich Proteins. *Arch. Oral Biol.* **1998**, *43* (9), 717–728.
- (102) Naurato, N.; Wong, P.; Lu, Y.; Wroblewski, K.; Bennick, A. Interaction of Tannin with Human Salivary Histatins. *J. Agric. Food Chem.* **1999**, *47* (6), 2229–2234.
- (103) Soares, S.; Brandão, E.; Mateus, N.; de Freitas, V. Interaction between Red Wine Procyanidins and Salivary Proteins: Effect of Stomach Digestion on the Resulting Complexes. *RSC Adv.* **2015**, *5* (17), 12664–12670.
- (104) Skopec, M. M.; Hagerman, A. E.; Karasov, W. H. Do Salivary Proline-Rich Proteins Counteract Dietary Hydrolyzable Tannin in Laboratory Rats? *J. Chem. Ecol.* **2004**, *30* (9), 1679–1692.
- (105) Zhang, J.; Kashket, S. Inhibition of Salivary Amylase by Black and Green Teas and Their Effects on the Intraoral Hydrolysis of Starch. *Caries Res.* **1998**, *32* (3), 233–238.
- (106) Kandra, L.; Gyémánt, G.; Zajác, Á.; Batta, G. Inhibitory Effects of Tannin on Human Salivary α -Amylase. *Biochem. Biophys. Res. Commun.* **2004**, *319* (4), 1265–1271.
- (107) D'Costa, A. S.; Bordenave, N. Inhibition of Starch Digestion by Flavonoids: Role of Flavonoid-Amylase Binding Kinetics. *Food Chem.* **2021**, *341*, 128256.

- (108) Freitas, D.; Le Feunteun, S.; Panouillé, M.; Souchon, I. The Important Role of Salivary α -Amylase in the Gastric Digestion of Wheat Bread Starch. *Food Funct.* **2018**, *9* (1), 200–208.
- (109) Rasouli, H.; Hosseini-Ghazvini, S. M.-B.; Adibi, H.; Khodarahmi, R. Differential α -Amylase/ α -Glucosidase Inhibitory Activities of Plant-Derived Phenolic Compounds: A Virtual Screening Perspective for the Treatment of Obesity and Diabetes. *Food Funct.* **2017**, *8* (5), 1942–1954.
- (110) Tan, Y.; Chang, S. K. C. Digestive Enzyme Inhibition Activity of the Phenolic Substances in Selected Fruits, Vegetables and Tea as Compared to Black Legumes. *J. Funct. Foods* **2017**, *38*, 644–655.
- (111) Yang, X.; Kong, F. Effects of Tea Polyphenols and Different Teas on Pancreatic α -Amylase Activity in Vitro. *LWT - Food Sci. Technol.* **2016**, *66*, 232–238.
- (112) Ayua, E. O.; Nkhata, S. G.; Namaumbo, S. J.; Kamau, E. H.; Ngoma, T. N.; Aduol, K. O. Polyphenolic Inhibition of Enterocytic Starch Digestion Enzymes and Glucose Transporters for Managing Type 2 Diabetes May Be Reduced in Food Systems. *Heliyon* **2021**, *7* (2), e06245.
- (113) Naz, S.; Siddiqi, R.; Dew, T. P.; Williamson, G. Epigallocatechin-3-Gallate Inhibits Lactase but Is Alleviated by Salivary Proline-Rich Proteins. *J. Agric. Food Chem.* **2011**, *59* (6), 2734–2738.
- (114) Lunagariya, N. A.; Patel, N. K.; Jagtap, S. C.; Bhutani, K. K. Inhibitors of Pancreatic Lipases: State of the Art and Clinical Perspectives. *EXCLI J.* **2014**, *13*, 897–921.
- (115) Rahim, A. T. M. A.; Takahashi, Y.; Yamaki, K. Mode of Pancreatic Lipase Inhibition Activity in Vitro by Some Flavonoids and Non-Flavonoid Polyphenols. *Food Res. Int.* **2015**, *75*, 289–294.
- (116) Kamoun, J.; Rahier, R.; Sellami, M.; Koubaa, I.; Mansuelle, P.; Lebrun, R.; Berlioz-Barbier, A.; Fiore, M.; Alvarez, K.; Abousalham, A.; Carrière, F.; Aloulou, A. Identification of a New Natural Gastric Lipase Inhibitor from Star Anise. *Food Funct.* **2019**, *10* (1), 469–478.
- (117) Butler, L. G. Effects of Condensed Tannin on Animal Nutrition. In *Chemistry and Significance of Condensed Tannins*; Hemingway, R. W., Karchesy, J. J., Branham, S. J., Eds.; Springer US: Boston, MA, 1989; pp 391–402.
- (118) Mole, S.; Butler, L. G.; Iason, G. Defense against Dietary Tannin in Herbivores: A Survey for Proline Rich Salivary Proteins in Mammals. *Biochem. Syst. Ecol.* **1990**, *18* (4), 287–293.
- (119) Lamothe, S.; Azimy, N.; Bazinet, L.; Couillard, C.; Britten, M. Interaction of Green Tea Polyphenols with Dairy Matrices in a Simulated Gastrointestinal Environment. *Food Funct.* **2014**, *5* (10), 2621–2631.
- (120) Tagliacruzchi, D.; Verzelloni, E.; Conte, A. Effect of Some Phenolic Compounds and Beverages on Pepsin Activity during Simulated Gastric Digestion. *J. Agric. Food Chem.* **2005**, *53* (22), 8706–8713.
- (121) van der Burg-Koorevaar, M. C. D.; Miret, S.; Duchateau, G. S. M. J. E. Effect of Milk and Brewing Method on Black Tea Catechin Bioaccessibility. *J. Agric. Food Chem.* **2011**, *59* (14), 7752–7758.
- (122) Donmez, O.; Mogol, B. A.; Gokmen, V.; Tang, N.; Andersen, M. L.; Chatterton, D. E. W. Modulation of Gastrointestinal Digestion of Beta-Lactoglobulin and Micellar Casein Following Binding by (–)-Epigallocatechin-3-Gallate (EGCG) and Green Tea Flavanols. *Food Funct.* **2020**, *11*, 6038–6053.
- (123) Oh, H.; Hoff, J. E. Effect of Condensed Grape Tannins on the In Vitro Activity of Digestive Proteases and Activation of Their Zymogens. *J. Food Sci.* **1986**, *51* (3), 577–580.
- (124) Gonçalves, R.; Mateus, N.; de Freitas, V. Biological Relevance of the Interaction between Procyanidins and Trypsin: A Multi-technique Approach. *J. Agric. Food Chem.* **2010**, *58* (22), 11924–11931.
- (125) Mehansho, H.; Ann, D. K.; Butler, L. G.; Rogler, J.; Carlson, D. M. Induction of Proline-Rich Proteins in Hamster Salivary Glands by Isoproterenol Treatment and an Unusual Growth Inhibition by Tannins. *J. Biol. Chem.* **1987**, *262* (25), 12344–12350.
- (126) Mole, S.; Rogler, J. C.; Morell, C. J.; Butler, L. G. Herbivore Growth Reduction by Tannins: Use of Waldbauer Ratio Techniques and Manipulation of Salivary Protein Production to Elucidate Mechanisms of Action. *Biochem. Syst. Ecol.* **1990**, *18* (2–3), 183–197.
- (127) Schwartz, M.; Neiers, F.; Feron, G.; Canon, F. The Relationship Between Salivary Redox, Diet, and Food Flavor Perception. *Front. Nutr.* **2021**, *7*, 612735.
- (128) Delimont, N. M.; Rosenkranz, S. K.; Haub, M. D.; Lindshield, B. L. Salivary Proline-Rich Protein May Reduce Tannin-Iron Chelation: A Systematic Narrative Review. *Nutr. Metab.* **2017**, *14* (1), 47.
- (129) Ginsburg, I.; Koren, E.; Shalish, M.; Kanner, J.; Kohen, R. Saliva Increases the Availability of Lipophilic Polyphenols as Antioxidants and Enhances Their Retention in the Oral Cavity. *Arch. Oral Biol.* **2012**, *57* (10), 1327–1334.
- (130) Mackie, A.; Bajka, B.; Rigby, N. Roles for Dietary Fibre in the Upper GI Tract: The Importance of Viscosity. *Food Res. Int.* **2016**, *88*, 234–238.
- (131) Zhao, Y.; Chen, L.; Yakubov, G.; Aminiafshar, T.; Han, L.; Lian, G. Experimental and Theoretical Studies on the Binding of Epigallocatechin Gallate to Purified Porcine Gastric Mucin. *J. Phys. Chem. B* **2012**, *116* (43), 13010–13016.
- (132) Georgiades, P.; Pudney, P. D. A.; Rogers, S.; Thornton, D. J.; Waigh, T. A. Tea Derived Galloylated Polyphenols Cross-Link Purified Gastrointestinal Mucins. *PLoS One* **2014**, *9* (8), e105302.
- (133) Cappai, M. G.; Wolf, P.; Pinna, W.; Kamphues, J. Pigs Use Endogenous Proline to Cope with Acorn (*Quercus Pubescens* Willd.) Combined Diets High in Hydrolysable Tannins. *Livest. Sci.* **2013**, *155* (2–3), 316–322.
- (134) Cai, K.; Hagerman, A. E.; Minto, R. E.; Bennick, A. Decreased Polyphenol Transport across Cultured Intestinal Cells by a Salivary Proline-Rich Protein. *Biochem. Pharmacol.* **2006**, *71* (11), 1570–1580.
- (135) Cai, K.; Bennick, A. Effect of Salivary Proteins on the Transport of Tannin and Quercetin across Intestinal Epithelial Cells in Culture. *Biochem. Pharmacol.* **2006**, *72* (8), 974–980.
- (136) D’Agostino, E. M.; Rossetti, D.; Atkins, D.; Ferdinando, D.; Yakubov, G. E. Interaction of Tea Polyphenols and Food Constituents with Model Gut Epithelia: The Protective Role of the Mucus Gel Layer. *J. Agric. Food Chem.* **2012**, *60* (12), 3318–3328.
- (137) Glumac, M.; Qin, L.; Chen, J.; Ritzoulis, C. Saliva Could Act as an Emulsifier during Oral Processing of Oil/Fat. *J. Texture Stud.* **2019**, *50* (1), 83–89.
- (138) Vingerhoeds, M. H.; Blijdenstein, T. B. J.; Zoet, F. D.; van Aken, G. A. Emulsion Flocculation Induced by Saliva and Mucin. *Food Hydrocoll.* **2005**, *19* (5), 915–922.