

The outward Shaker channel OsK5.2 improves plant salt tolerance by contributing to control of both leaf transpiration and  $K^+$  secretion into xylem sap

Running head: OsK5.2  $K^+$  channel contribution to salt tolerance

**Jing Zhou, Thanh Hao Nguyen<sup>1</sup>, Dorsaf Hmidi, Doan Trung Luu, Hervé Sentenac, Anne-Aliénor Véry\***

BPMP, Univ Montpellier, CNRS, INRAE, Institut Agro, Montpellier, France

<sup>1</sup>Present address: University of Glasgow, Institute of Molecular Cell & System Biology, Bower Building, Glasgow G12 8QQ, UK

\*For correspondence. E-mail [anne-alienor.very@cnrs.fr](mailto:anne-alienor.very@cnrs.fr)

## **Funding**

This work was supported in part by a grant from the China Scholarship Council (to J.Z.), by a doctoral fellowship from the French Embassy in Vietnam (to T.H.N.), and by an ANR-DFG grant (ANR-20-CE92-0005 to A.A.V.).

## **Acknowledgments**

We are grateful to Emmanuel Guiderdoni, Christian Chaine, Eve Lorenzini, Christophe Périn and Remy Michel (UMR AGAP, Montpellier) for the rice mutant line amplification, and to Aurélie Putois (UMR IATE, Montpellier) for assistance with picture acquisitions at the Scanning Electron Microscope.

## Abstract

Soil salinity constitutes a major environmental constraint to crop production worldwide. Leaf  $K^+/Na^+$  homeostasis, which involves regulation of transpiration, and thus of the xylem sap flow, and control of the ionic composition of the ascending sap, is a key determinant of plant salt tolerance. Here, we show, using a reverse genetics approach, that the outwardly-rectifying  $K^+$ -selective channel OsK5.2, which is involved in both  $K^+$  release from guard cells for stomatal closure in leaves and  $K^+$  secretion into the xylem sap in roots, is a strong determinant of rice salt tolerance (plant biomass production and shoot phenotype under saline constraint). *OsK5.2* expression was up-regulated in shoots from the onset of the saline treatment, and OsK5.2 activity in guard cells led to fast decrease in transpirational water flow and therefore reduced  $Na^+$  translocation to shoots. In roots, upon saline treatment, OsK5.2 activity in xylem sap  $K^+$  loading was maintained, and even transiently increased, outperforming the negative effect on  $K^+$  translocation to shoots resulting from the reduction in xylem sap flow. Thus, the overall activity of OsK5.2 in shoots and roots, which both reduces  $Na^+$  translocation to shoots and benefits shoot  $K^+$  nutrition, strongly contributes to leaf  $K^+/Na^+$  homeostasis.

## Keywords (5-10)

Outward  $K^+$  channel, Shaker channel, salt tolerance, rice,  $K^+/Na^+$  homeostasis, transpirational flux, xylem sap, root-to-shoot translocation, *Tos17* insertion mutants

46

47 **1 | INTRODUCTION**

48 High soil salinity is a widespread environmental constraint over the world that causes  
49 substantial restrictions in production and quality of a majority of crops, including cereals.  
50 Understanding how plants cope with high salinity in the environment is thus an issue of great  
51 agricultural importance. Rice is rated as a salt-sensitive cereal (Munns & Tester, 2008; Zeng  
52 & Shannon, 2000), and salinity levels have increased in rice fields, particularly due to the  
53 climate change and sea level rise, which strongly challenge rice culture in coastal regions.

54 The adverse effects of high soil salinity on plant growth are mainly related to the  
55 decrease in osmotic potential of the soil and to ionic toxicity of  $\text{Na}^+$  in leaves (Munns &  
56 Tester, 2008). Studies on the latter phenomenon have revealed that genes encoding  $\text{Na}^+$   
57 transport systems correspond to major quantitative trait loci (QTLs) of salt tolerance (Hauser  
58 & Horie, 2010). In return, such findings have spurred research efforts in this domain of  
59 membrane transport biology, highlighting also that  $\text{Na}^+$  detrimental effects are counteracted  
60 by the plant's ability to take up the essential macronutrient  $\text{K}^+$  and control its  $\text{K}^+$  nutritional  
61 status in presence of high external  $\text{Na}^+$  concentrations (Maathuis & Amtmann, 1999).  $\text{K}^+$  is  
62 involved in vital functions such as enzyme activation, the cytoplasmic pH homeostasis,  
63 control of cell membrane potential and cell turgor-driven movements (Marschner, 2011;  
64 Nieves-Cordones, Al Shiblawi & Sentenac, 2016). Upon salt stress, the massive influx of  
65 positively charged  $\text{Na}^+$  ions causes cell membrane depolarization, which reduces the driving  
66 force for  $\text{K}^+$  uptake and even in some cases leads to channel-mediated root  $\text{K}^+$  losses  
67 (Jayakannan, Bose, Babourina, Rengel, & Shabala, 2013; Rubio, Nieves-Cordones, Horie, &

Shabala, 2020). Thus, plant exposure to high salinity is inevitably accompanied by chronic  $K^+$  deficiency, which affects the leaf  $K^+$  to  $Na^+$  content ratio, whose maintenance to a high value is a key determinant of salt tolerance (Hauser & Horie, 2010; Maathuis & Amtmann, 1999).

$K^+$  and  $Na^+$  ions taken up by root cells can migrate to stelar tissues and be translocated to leaves by the upward flow of sap in the xylem vessels. Control of the ionic composition of xylem sap, involving membrane ion transport processes in parenchyma cells along the sap ascent pathway, is thus a major determinant of salt tolerance, together with control of the flux of xylem sap, which is driven by leaf transpiration and hence dependent on the level of stomatal aperture, or driven by the so-called root pressure, resulting from increased osmotic pressure in the xylem vessels due to increased solute concentration in the sap in absence of significant plant transpiration (Jeschke, 1984; Marschner, 2011). Therefore, transport systems contributing to  $Na^+$  or  $K^+$  secretion/retrieval into/from the xylem sap or to regulation of stomatal aperture can contribute to processes that play crucial roles in salt tolerance.

In various plant species,  $Na^+$  transporters from the HKT family have been shown to contribute to  $Na^+$  retrieval from the xylem sap and loading into the xylem parenchyma cells bordering the vessels, *i.e.* to the so-called "sap desalinization" process (Hauser & Horie, 2010). In rice, the *HKT* transporter genes identified as involved in this process are *OsHKT1;5*, which is mainly expressed in root xylem parenchyma cells and corresponds to the major salt-tolerance QTL *SKCI* (Ren et al., 2005), *OsHKT1;4*, expressed in both root and basal leaf xylem tissues (Suzuki et al., 2016; Khan et al., 2020), and *OsHKT1;1* expressed in both xylem and phloem and thereby also involved in  $Na^+$  recirculation from leaves to roots within

the phloem sap, thereby making an additional contribution to promoting Na<sup>+</sup> accumulation in roots compared to leaves (Campbell et al., 2017; Wang et al., 2015).

Compared with the large number of studies focused on the Na<sup>+</sup> transporters controlling Na<sup>+</sup> translocation to leaves and thereby contributing to maintain the ratio of leaf K<sup>+</sup> to Na<sup>+</sup> contents to a high value, less attention has been paid to the K<sup>+</sup> transport mechanisms that operate under saline conditions and ensure efficient K<sup>+</sup> supply to leaves. Current knowledge in this domain essentially concerns K<sup>+</sup> transport systems involved in root K<sup>+</sup> uptake, and mainly high-affinity K<sup>+</sup> transporters from the HAK/KUP/KT family, AtHAK5 in Arabidopsis and OsHAK1, OsHAK5, OsHAK16 and OsHAK21 in rice (Chen et al., 2015; Feng et al., 2019; Nieves-Cordones, Alemán, Martínez & Rubio, 2010; Shen et al., 2015; Yang et al., 2014).

In rice, the outwardly rectifying Shaker K<sup>+</sup> channel OsK5.2 is involved both in K<sup>+</sup> secretion into the xylem sap for translocation toward the shoots and in control of stomatal aperture and leaf transpiration by driving K<sup>+</sup> efflux from guard cells for stomatal closure (Nguyen et al., 2017). OsK5.2 is homologous to two outwardly rectifying K<sup>+</sup> channels from Arabidopsis, SKOR and GORK, respectively involved in K<sup>+</sup> secretion into the xylem sap (Gaymard et al., 1998) and K<sup>+</sup> efflux from guard cells for stomatal closure (Hosy et al., 2003). Gathering the two functions of its Arabidopsis counterparts, OsK5.2 emerged as a good model to assess the level of contribution of these functions to the control of Na<sup>+</sup> and K<sup>+</sup> delivery to shoots upon saline conditions and salt tolerance. This has been achieved in the present study by phenotyping *osk5.2* knock-out (KO) mutant plants subjected to saline conditions. We found that the lack of functional *OsK5.2* expression does result in increased

plant sensitivity to salt stress and analyzed the bases of the salt sensitive phenotype.

## **2 | MATERIALS AND METHODS**

### **2.1 | Plant growth and salt treatment**

The selection from *Tos17*-insertion lines of *osk5.2* mutant and corresponding wild-type (WT) plants in the background of rice Nipponbare cultivar (*Oryza sativa* L. ssp. japonica cv. Nipponbare) has been previously described (Nguyen et al., 2017). Rice seeds were germinated on a raft floating on deionized water for one week. The seedlings were then hydroponically grown on half-strength Yoshida medium for one week, and thereafter on Yoshida medium (0.5 mM (NH<sub>4</sub>)<sub>2</sub>SO<sub>4</sub>, 1.6 mM MgSO<sub>4</sub>, 1.2 mM Ca(NO<sub>3</sub>)<sub>2</sub>, 0.7 mM KNO<sub>3</sub>, 0.8 mM KH<sub>2</sub>PO<sub>4</sub>, 60 μM Na<sub>2</sub>FeEDTA, 20 μM MnSO<sub>4</sub>, 0.32 μM (NH<sub>4</sub>)<sub>6</sub>Mo<sub>7</sub>O<sub>24</sub>, 1.4 μM ZnSO<sub>4</sub>, 1.6 μM CuSO<sub>4</sub>, 45.2 μM H<sub>3</sub>BO<sub>3</sub>, and pH adjusted to 5.5 with H<sub>2</sub>SO<sub>4</sub>). Five-week-old rice plants were subjected to salt treatment by supplementing the hydroponic Yoshida medium with 50 mM NaCl for up to 14 days (and also 100 mM for 7 days in growth assays). The rice plants were grown in a growth chamber (70% relative humidity, light intensity 130 photon μmol.m<sup>-2</sup>.s<sup>-1</sup>, 29°C/25°C 12 h/12 h day/night).

### **2.2 | RNA extraction and quantitative real time PCR experiments**

Five-week-old Nipponbare plants grown on Yoshida medium were either supplemented with 50 mM NaCl for 14 days and then transferred back to standard Yoshida medium for three days, or further grown during this period on Yoshida medium (control batch). Total RNAs were extracted from samples collected at same times from salt-treated or control plant

batches using the RNeasy plus mini kit with gDNA eliminator (Qiagen, Germany). First-strand cDNAs were synthesized from 3  $\mu$ g of RNAs using SuperScript III reverse transcriptase (Invitrogen) and used as template for qRT-PCR experiments. qRT-PCR analyses were performed using the Lightcycler480 system (Roche diagnostics) and SYBR *Premix Ex Taq* (Takara) in a total volume of 10  $\mu$ l, which contained 2  $\mu$ l of cDNA, 3  $\mu$ l of forward and reverse primer mixture (1  $\mu$ M), and 5  $\mu$ l of SYBR *Premix Ex Taq*. Reactions were performed with three independent biological replicates, each one with three technical replicates (PCR program: 95°C for 30 sec; 45 cycles of 95°C for 10 sec, 60°C for 10 sec, and 72°C for 15 sec; followed by a melt cycle from 60°C to 95°C).  $C_T$  (cycle threshold) values were obtained from amplification data using a threshold of 0.37. The *OsK5.2* absolute number of copies was calculated according to standard curves obtained by successive dilutions with known quantities of *OsK5.2*, and then normalization using the  $C_T$  values of three housekeeping genes (ubiquitin-like protein gene *SMT3*, PP2A-interactor gene *Tip41* and elongation factor gene *EF1beta*) as described in Khan et al. (2020). The sequences of the primers used for qRT-PCR experiments are provided in Table S1.

### 2.3 | $\text{Na}^+$ and $\text{K}^+$ assays in tissues and xylem sap

Five-week-old plants hydroponically grown as described above were supplemented or not with 50 mM NaCl for 14 days. Excised root systems and shoots were periodically collected during the salt treatment, both from salt treated and control plants. The roots (rinsed in deionized water) and shoots were dried (60°C for 3 days) and weighed. Ions were extracted from the tissues in 0.1 N HCl for 3 days and assayed by flame spectrophotometry (SpectrAA

220FS, Varian).

Xylem sap was collected through natural exudation under control condition from de-topped plants (3 cm above the root system). Upon salt treatment, xylem sap was obtained through pressurization. The root system of the de-topped plants was placed into a pressure chamber (Boursiac *et al.*, 2005) filled with hydroponic medium containing 50 mM NaCl, and sealed with a silicon dental paste (PRESIDENT Light Body, Coltene, Switzerland). The first few drops (2  $\mu$ l) were discarded to avoid the contamination that results from injured cells. Twenty  $\mu$ l of sap samples were then collected using a micro-pipette, transferred into 0.2 ml Eppendorf tubes kept on ice, and diluted in 0.1 N HCl for Na<sup>+</sup> and K<sup>+</sup> assay (flame spectrophotometry).

## 2.4 | Leaf transpiration

Intact plants were transferred into a multipotometer device (Nguyen *et al.*, 2017) one day before transpiration rate measurement. The root system of each intact plant was inserted into a 50 ml syringe filled with hydroponic medium through the rubber plunger, and sealed with a silicon dental paste. Each syringe was connected to a graduated 1-ml serological plastic pipette via a thin silicone tube. The pipettes were refilled with the same medium with proper time intervals, making sure that no air bubble was present in the system. A camera took photographs of the set of pipettes every two minutes in order to record the changes in water level in the pipettes. Image capture was started after 3 h under light condition and maintained for 30 minutes. Plants were then exposed to darkness for 5 h under continuous recording. The rate of decrease in the water volume in the pipettes was used to calculate the mean



transpiration rate of plants. After transpiration rate measurements, each plant was placed on a paper sheet of known dimension, the blades being thoroughly positioned with small pieces of double-sided tape in order to fix the abaxial face on the paper sheet and to avoid blade superposition. A picture of the plant was then taken and the whole area of leaf tissues was determined using imageJ software ([imagej.nih.gov/ij](http://imagej.nih.gov/ij)) for transpiration rate value normalization.

### 3 | RESULTS

#### 3.1 | *osk5.2* mutant plants exhibit increased sensitivity to salinity

The two *osk5.2* KO mutant lines ASJA08 and ASHF06 (Nguyen et al., 2017) and the corresponding wild type (WT) plants displayed a very similar development and phenotype when growth occurred in absence of salt stress (Figure 1). This was no longer the case when the plants were subjected to saline conditions (6-week-old plants subjected to 50 or 100 mM NaCl for 7 days). The mutant plants then displayed a severe reduction in biomass production when compared with that of plants from the same mutant lines but grown in control conditions, by about 25-35% with 50 mM NaCl and 45-50% with 100 mM NaCl (Figure 1a, b). Conversely, the corresponding reduction observed for the WT plants appeared weak (~2% with 50 mM NaCl and less than 10-15% with 100 mM NaCl) and not statistically significant (Figure 1b). While the biomass of mutant plants in control conditions was similar to that of corresponding wild type plants, the length of leaves and roots were significantly reduced in mutant plants, only slightly in leaves (by ~10%) but more in roots (by 15-20%)(Figure S1). The saline treatments however, especially stunted leaf growth in the mutant plants, increasing

the leaf length difference between mutant and wild type plants to 20% with 100 mM NaCl (Figure S1). The saline treatments had no differential effect on root length between mutant and wild type plants, on the other hand (Figure S1). Shoot water content decreased slightly with the most severe salt treatment but, whatever the treatment, no significant difference in shoot or root water content was found between mutant and wild-type plants (Figure S2). More “dried” leaf tips in the mutant than the WT plants were however observed under the most severe salt treatment (Figure 1a). Altogether, these analyses indicated that the lack of *OsK5.2* functional expression resulted in reduced tolerance to saline conditions.

### **3.2 | *OsK5.2* transcript accumulation under salt stress**

Real-time qRT-PCR analyses revealed that *OsK5.2* was expressed in both roots and leaves (Figure 2). In plants grown in control condition, *OsK5.2* transcripts were 4 folds more abundant in roots than in leaves. When the plants were subjected to saline conditions (50 mM NaCl), a change in the relative expression of *OsK5.2* between roots and shoots was observed leading to balanced expression in the plant. In leaves, the accumulation of *OsK5.2* transcripts was rapidly (from one day after salt treatment) up-regulated by about 3 folds, and remained high during 7 days (Figure 2b). The leaf level of *OsK5.2* transcripts appeared to decrease with longer exposure to the saline conditions but, after 14 days of salt treatment, it was still about 1.5-fold that observed in control conditions (Figure 2b). Recovery from salt stress for 1 to 3 days further decreased the leaf level of *OsK5.2* transcripts, down to that observed in leaves from control plants. In roots, as compared with leaves, the accumulation of *OsK5.2* transcripts showed less variation in response to the saline treatment and tended to slightly

decrease throughout the saline treatment period, a down-regulation by about 25% being statistically significant at days 3 and 14 of the treatment (Figure 2a). After 1 day of recovery from salt stress, the root level of *OsK5.2* transcripts recovered to same expression level of control plants and then remained stable for at least 2 days.

### **3.3 | *osk5.2* mutant plants display larger transpirational water loss than WT plants under salt stress**

Stomatal aperture measurement is hardly feasible in rice due to the small size of rice stomata and the dumbbell-shaped guard cells, which lead to extremely thin aperture of the stomatal pore (Figure S3; Huang et al., 2009; Matsuda et al., 2016). Absence of noticeable effect of *OsK5.2* loss-of-function on stomatal shape was found on the abaxial side of the flag leaf using scanning electron microscopy (Figure S3). The role of *OsK5.2* in control of leaf transpiration under salt stress was then investigated by adding 50 mM NaCl into the hydroponics solution of 5-week-old *osk5.2* mutant and WT plants, and performing measurements of the steady-state rates of plant water loss periodically for 14 days (on days 1, 3, 7 and 14 of the salt treatment) both under light and in dark conditions using a multipotometer (Figure 3). The data obtained with the 2 *osk5.2* mutant lines ASJA08 and ASHF06 led to the same conclusions. The salt treatment strongly decreased the rate of transpirational water loss in the *osk5.2* mutant and the corresponding WT plants, by up to 50 to 55 % in the light and 30 to 40% in the dark conditions (Figure 3). The kinetics of reduction of transpiration rate upon NaCl exposure was clearly slower in *osk5.2* mutant compared with WT plants both in light and dark conditions. During the two weeks of salt treatment, a higher

rate of transpirational water loss was consistently observed in *osk5.2* mutant as compared with WT plants in light and dark conditions, and the difference was highly significant in most of the analyzed time points (Figure 3). The greatest difference in transpiration rate between WT and *osk5.2* mutant plants occurred after one day of treatment owing to the slower response to NaCl exposure in *osk5.2* mutant plants.

### **3.4 | Na<sup>+</sup> and K<sup>+</sup> concentrations in xylem sap under salt stress and translocation fluxes towards the shoots**

Xylem sap samples were collected from de-topped plants subjected to the same protocol of salt treatment as that used in the experiment described by Figure 3: 2 weeks in 50 mM NaCl hydroponics solution applied to 5-week-old plants previously grown under control conditions. The concentrations of K<sup>+</sup> and Na<sup>+</sup> determined in sap samples (Figure 4) and the transpiration rates recorded under the same experimental conditions (Figure 3) were used to estimate the transpiration-driven fluxes of K<sup>+</sup> and Na<sup>+</sup> (Figure 5) arriving in shoots under such conditions.

Under control condition, K<sup>+</sup> concentrations measured in xylem sap were close to 11 mM (7-folds the K<sup>+</sup> concentration in the hydroponic medium) in WT plants, and were 30 to 40% lower in the two *osk5.2* mutant lines (as previously reported; Nguyen et al., 2017). The xylem sap K<sup>+</sup> concentration in both WT and *osk5.2* mutant plants displayed transient variations upon exposure to the saline conditions (Figure 4a). The concentrations observed in the *osk5.2* mutant plants remained lower than those displayed by the corresponding WT plants by about 40% to 50% over the entire duration of the salt treatment (Figure 4a). With respect to Na<sup>+</sup>, the concentration of this cation in the xylem sap was extremely low, in the submillimolar

range, in the absence of salt treatment (Figure 4b). Exposure to 50 mM NaCl led to the loading of a large amount of  $\text{Na}^+$  into the xylem sap with no significant difference between WT and *osk5.2* mutant plants (Figure 4b). The  $\text{Na}^+$  concentrations measured in the xylem sap of the two types of plants were close to that in the hydroponic medium (50 mM) after one day of NaCl supplementation, and remained fairly stable during the two weeks of salt treatment. Altogether, these results indicated that the lack of *OsK5.2* functional expression constitutively resulted in a large reduction in xylem sap  $\text{K}^+$  concentration, by *ca.* 40-50%, but did not affect xylem sap  $\text{Na}^+$  concentration. As a result, the  $\text{K}^+/\text{Na}^+$  xylem sap concentration ratios, computed from the data provided by Figure 4a and 4b, appear consistently higher in WT than in *osk5.2* mutant plants (Figure 4c).

The estimated  $\text{K}^+$  flux arriving in shoots during the light period in these experimental conditions, obtained by integrating the data from Figure 3a and 4a (Figure 5a), is decreased by the salt treatment in both the *osk5.2* mutant and WT plants. It is lower in both *osk5.2* mutant lines than in the corresponding WT plants, except at day 1 of the salt treatment due to the sharp decrease in transpiration rate displayed by WT plants at this time point. The differences between the WT and mutant plants under salt treatment are in the 30 to 50% range from day 3 of the salt treatment, *i.e.*, similar to those observed between the two types of plants in control conditions (Figure 5a).

The estimated flux of  $\text{Na}^+$  arriving in the shoots of WT and *osk5.2* mutant plants is very low under control conditions (this cation being then present as trace contaminant in the hydroponics solution). A marked increase in  $\text{Na}^+$  flux is observed in all genotypes from the first day of exposure to NaCl (Figure 5b). The flux is larger in *osk5.2* mutant than in WT

plants, by ca. 20 to 40% during the first week of the salt treatment, which essentially reflects the difference in transpiration rate between the two types of plants during this period (see Figure 3).

### 3.5 | *osk5.2* mutant plants accumulate less K<sup>+</sup> and more Na<sup>+</sup> under salt stress

K<sup>+</sup> and Na<sup>+</sup> contents were determined in roots and shoots of *osk5.2* and WT plants subjected to the same salt treatment protocol as that previously used.

In all genotypes, root and shoot K<sup>+</sup> contents, and thus whole plant K<sup>+</sup> contents, decreased with the duration of the salt treatment (Figure 6a, b, respectively). In shoots, the decrease was clearly more pronounced in the *osk5.2* mutant lines, when compared with the corresponding WT plants, and the relative difference in shoot K<sup>+</sup> contents between the mutant and WT plants increased with the duration of the salt treatment (Figure 6b): the difference was in the range of 10-20% at the beginning of the treatment (in the absence of NaCl addition and at day 1 of the salt treatment), and reached 40-45% after two weeks of treatment (Figure 6b). In roots, the impact of lacking *OsK5.2* functional expression on K<sup>+</sup> contents appeared much weaker than that observed in shoots, mostly not statistically significant (Figure 6a).

Regarding Na<sup>+</sup>, the contents of this cation were very low in all plants, whatever their genotype, in the absence of salt treatment (Figure 7). The salt treatment increased both the root and shoot (and thus the whole plant) contents of this cation, from the first day of treatment and over the two weeks of treatment, in all plant genotypes (Figure 7). Under all conditions except the longest duration of the salt treatment (in other words, under control conditions and during the first week of salt exposure), significantly higher Na<sup>+</sup> contents were

found in the *osk5.2* mutants than in the corresponding WT plants, for both mutant lines, by more than 20% in roots (Figure 7a) and 35% in shoots (Figure 7b). The relative differences between the mutant and WT plants were weaker at the last time point of salt treatment (after 14 days), and the differences were no longer statistically significant (except for the roots of one mutant line). At this time,  $\text{Na}^+$  levels in shoots exceeded those in roots in all genotypes (Figure 7a, b). Thus,  $\text{Na}^+$  accumulation was higher in *osk5.2* mutant plants than in the corresponding WT plants until the late stage of  $\text{Na}^+$  plant invasion, when the level of  $\text{Na}^+$  in shoots had become higher than that in roots.

$\text{K}^+/\text{Na}^+$  content ratios were calculated from the data displayed by Figures 6 and 7. The ratios were significantly lower in *osk5.2* mutant plants than in WT plants both in shoots and roots under all conditions except in roots of one mutant line at the last time point of salt treatment (Figure 8). In salt-stressed leaves, the relative reduction in  $\text{K}^+/\text{Na}^+$  content ratios observed in the *osk5.2* mutant plants as compared with the corresponding WT plants, was in the range of 35-55% (Figure 8). Thus, *OsK5.2* lack of functional expression strongly impaired  $\text{K}^+/\text{Na}^+$  homeostasis in leaves under salt stress.

## 4 | DISCUSSION

### 4.1 | *OsK5.2*, a model for assessing the involvement of $\text{K}^+$ channels in the control of $\text{K}^+$ and $\text{Na}^+$ translocation to the shoots under salt stress

Mechanisms that control long distance transport of  $\text{Na}^+$  and  $\text{K}^+$  in the plant vasculature contribute to maintaining the shoot  $\text{K}^+/\text{Na}^+$  content ratio at a high value, which is a key determinant of salt tolerance (Munns and Tester, 2008; Maathuis, Ahmad and Patishtan, 2014;

Ismail and Horie, 2017; Wu, Zhang, Giraldo and Shabala, 2018).

Regarding  $\text{Na}^+$ , clear evidence has been obtained, in rice, as well as in Arabidopsis and various other species, that  $\text{Na}^+$  transporters belonging to the HKT family are involved in desalinization of the ascending xylem sap (Hauser & Horie, 2010). The  $\text{H}^+/\text{Na}^+$  antiport system SOS1 has been suggested to also contribute to this function when the concentration of  $\text{Na}^+$  in the xylem sap reaches very high values (Maathuis et al., 2014).

Regarding the mechanisms controlling  $\text{K}^+$  translocation to shoots, outwardly rectifying channels belonging to the Shaker family (SKOR in Arabidopsis and OsK5.2 in rice; see below), have been shown to mediate  $\text{K}^+$  secretion into the xylem sap under normal conditions (Gaymard et al., 1998; Nguyen et al., 2017) but their contribution to this function under salt stress remains poorly documented. Furthermore,  $\text{K}^+$  secretion into the xylem sap may have to be active, under some environmental conditions (Wu, Zhang, Giraldo and Shabala, 2018), which would exclude channel-mediated (passive) contribution to this function in such conditions. Also, active  $\text{H}^+$ -coupled  $\text{K}^+$  transports mediated by HAK/KUP/KT transporters (see below) or involving a NRT1/PTR family member, NRT1;5 (Li et al., 2017), have been shown to contribute to  $\text{K}^+$  translocation towards the shoots. In rice, OsHAK1 and OsHAK5, which are thought to be endowed with  $\text{H}^+$ - $\text{K}^+$  symport activity (Véry et al., 2014), have been shown to play a role in  $\text{K}^+$  translocation towards the shoots under saline conditions (Chen et al., 2015; Yang et al., 2014). The mechanisms that underlie these contributions remain however to be specified.  $\text{H}^+$ - $\text{K}^+$  symport activity in parenchyma cells bordering the xylem vessels would result in  $\text{K}^+$  retrieval from the xylem sap since the pH gradient between the sap and the cytoplasm is inwardly directed and thus favors  $\text{K}^+$  influx into the cells. It has thus



been hypothesized that such  $H^+$ - $K^+$  symporters may allow  $K^+$  acquisition within the stele by parenchyma cells, and that this would result in a higher concentration of  $K^+$  in xylem-adjacent cells, and thus in an outwardly-directed  $K^+$  electrochemical gradient that would allow (passive) channel-mediated  $K^+$  release into the sap (Yang et al., 2014).

At the leaf surface, control of stomatal aperture (and hence of transpiration rate) provides another type of contribution to salt tolerance. Exposure to saline conditions has been shown to rapidly result in stomatal closure, which limits the flux of xylem sap, and thus the rate of  $Na^+$  translocation to shoots (Fricke et al., 2006; Huang et al., 2009; Hedrich and Shabala, 2018). Such a control is however likely to also affect the rate of  $K^+$  translocation to shoots, and thus its contribution to shoot  $K^+/Na^+$  homeostasis should benefit from mechanisms allowing to counteract the depressive effect of the reduction in volumetric flow of xylem sap on  $K^+$  translocation (see § 4.4).

The two outwardly rectifying Shaker  $K^+$  channels in Arabidopsis, SKOR and GORK, are involved in xylem sap  $K^+$  loading (Gaymard et al., 1998), and in guard cell  $K^+$  release-mediated stomatal closure (Hosy et al., 2003), respectively. The roles of SKOR and GORK in Arabidopsis salt tolerance remain, however, poorly documented. *OsK5.2*, which is, like SKOR and GORK, an outwardly rectifying Shaker channel, had been previously shown to be involved both in xylem sap  $K^+$  loading and in guard cell  $K^+$  release-mediated stomatal closure (Nguyen et al., 2017). The expression level of *OsK5.2* is fairly maintained in roots under saline conditions, and even increased in shoots in these conditions (Figure 2). This channel has thus been used as a model in the present report to investigate xylem sap  $K^+$  loading under salt stress, i.e., whether it can be channel mediated or requires active transport

systems (e.g.,  $H^+/K^+$  antiporters?), and the involvement of transpiration rate control in salt tolerance.

#### 4.2 | $K^+$ Secretion into the xylem sap under salt stress

Saline conditions weakly affected the expression level of *OsK5.2* in roots (Figure 2a), as shown for its counterpart *SKOR* in Arabidopsis roots (Pilot et al., 2003). In line with this rather stable expression, the contribution of *OsK5.2* to  $K^+$  secretion into the xylem sap (estimated from the difference in sap concentration between the WT and *osk5.2* mutant plants; Figure 4a) did not appear to be much modified by the salt treatment. This contribution even tended to slightly increase during the first week of the treatment, which may be due to increased driving force for  $K^+$  secretion under conditions of salt-induced membrane depolarization (Jayakannan et al., 2013; Mian et al., 2011).

Reliable measurements of both the membrane potential and the apoplastic  $K^+$  concentration of stelar cells are difficult to obtain. However, the fact that *OsK5.2* can contribute to  $K^+$  secretion under salt stress provides definitive evidence that passive (since channel-mediated) secretion of  $K^+$  can occur in stelar cells of rice plants facing saline conditions. This conclusion, which does not exclude a contribution of active  $K^+$  transport mechanisms to  $K^+$  secretion under saline conditions, also means that other channels besides *OsK5.2*, either  $K^+$ -selective and belonging to the Shaker family (Véry et al., 2014) or poorly  $K^+$ -selective like NSCC channels identified in stelar cells by patch clamp experiments (Wegner & de Boer, 1997), could also contribute to  $K^+$  secretion into xylem sap under such conditions.

### 4.3 | Reduction of the volumetric flux of xylem sap under salt stress

Exposure to saline conditions is known to rapidly result in reduced stomatal aperture and plant transpiration (Fricke et al., 2006; Hedrich & Shabala, 2018; Robinson, Véry, Sanders, & Mansfield, 1997). In the present study, the transpiration rate was similarly reduced in WT and *osk5.2* mutant plants at the end of the salt treatment, by about 50% under light conditions and 30% under dark conditions (Figure 3). The kinetics of the reduction in transpiration rate was however more rapid in WT than in *osk5.2* mutant plants.

Upon an increase in external medium salinity, abscisic acid (ABA) produced in response to the resulting osmotic stress is rapidly directed to guard cells, where it is expected to activate the PYR/PYL/RCAR-ABI1 PP2C phosphatase-OST1 SnRK kinase signaling pathway, leading to guard cell anion channel activation and stomatal closure (Hedrich & Shabala, 2018). The actual contribution of guard cell anion channels to the triggering of stomatal closure upon salt stress has however been little investigated so far. Likewise, the role in stomatal closure upon salt stress of the K<sup>+</sup> outward channels acting as downstream effectors (Pandey et al., 2007; Schroeder et al., 2001) was still poorly documented. Indeed, although extensive analyses have concerned the integrated involvement of transport systems in regulation of guard cell turgor (Jezek & Blatt, 2017), little information is related to high salinity conditions (Lebaudy et al., 2008; Thiel & Blatt, 1991; Véry, Robinson, Mansfield, & Sanders, 1998). Here, our data reveal the important role of channel-mediated outward K<sup>+</sup> transport activity in guard cells in control of transpiration rate upon salt stress. The whole plant transpiration data (Figure 3) suggest that OsK5.2 activity in stomata contributed to the

reduction in stomatal aperture over the entire duration of the salt treatment. Furthermore, our data indicate that this activity is of particular importance at the onset of salt stress by allowing a more rapid reduction in the rate of transpiration (Figure 3).

#### **4.4 | Na<sup>+</sup> and K<sup>+</sup> translocation to shoots by the xylem sap and K<sup>+</sup>/Na<sup>+</sup> shoot homeostasis**

An apoplastic pathway strongly contributing to Na<sup>+</sup> entry into the root and radial migration to the root vasculature (the so-called bypass flow across the root to the xylem) has been evidenced in rice in the presence of high Na<sup>+</sup> concentrations (Faiyue, Al-Azzawi, & Flowers, 2012; Flam-Shepherd et al., 2018; Yeo, 1998). In our experimental conditions, the concentration of Na<sup>+</sup> in the xylem sap in both WT and *osk5.2* mutant plants was quite similar to that in the hydroponic medium (50 mM) (Figure 4). Also, the xylem sap K<sup>+</sup>/Na<sup>+</sup> concentration ratio decreased very rapidly down to values lower than 0.1 after one day of salt treatment, and was then more than 10 times lower than the K<sup>+</sup>/Na<sup>+</sup> root content ratio (Figure S4). This indicates that the former ratio (in the xylem sap) was not likely to reflect the corresponding ratio in the root symplasm. Altogether, these results support the hypothesis that the bypass flow of Na<sup>+</sup> was the major determinant of the migration of this cation towards the xylem vasculature. In such conditions, the flux of Na<sup>+</sup> translocated to the shoot becomes proportional to the volumetric flow of xylem sap. Since the Na<sup>+</sup> concentration of the xylem sap was similar in the mutant and WT plants, the larger rate of transpiration under salt stress in the mutant plants (Figure 3) due to impaired control of stomatal aperture is the major determinant of the difference in Na<sup>+</sup> translocation rate between the two types of plants (Figure 5b). Thus, OsK5.2-dependent control of transpiration rate results in a reduction of

Na<sup>+</sup> translocation towards the shoots.

Despite the positive effect on salt stress tolerance of a reduction in xylem sap volumetric flow, which decreases the flux of translocated Na<sup>+</sup> to shoots, a reduction in xylem sap volumetric flow is also likely to have negative impact on the rate of K<sup>+</sup> translocation to shoots (Figure 5a). Nevertheless, due to the dual role of OsK5.2 in controlling transpiration rate and K<sup>+</sup> secretion into xylem sap, WT plants expressing *OsK5.2* showed a smaller reduction in the rate of K<sup>+</sup> translocation to shoots compared to mutant plants. In other words, although OsK5.2 activity in stomata decreases the transpiration rate (Figure 3) and thus has a negative effect on K<sup>+</sup> translocation to shoots, the contribution of OsK5.2 to K<sup>+</sup> loading into the xylem sap (Figure 4a) outperforms the "negative" effect resulting from reduced xylem volumetric flow (Figure 5a). This conclusion supports the hypothesis that the beneficial effect, in terms of control of Na<sup>+</sup> translocation to shoots and tolerance to salinity, of the reduction in transpiration rate upon salt stress is likely to integrate the plant ability to increase, or at least maintain, the rate of K<sup>+</sup> secretion into the xylem sap.

Due to the beneficial effects of the overall OsK5.2 activity, the ratio of the K<sup>+</sup> to Na<sup>+</sup> translocation rates towards the shoots (identical to the xylem sap K<sup>+</sup>/Na<sup>+</sup> concentration ratio; Figure 4c) is larger in WT than in *osk5.2* mutant plants. This is probably the main reason why the kinetics of the decrease in shoot K<sup>+</sup>/Na<sup>+</sup> content ratio is slower in WT than in mutant plants (Figure 8b), and why the overall activity of OsK5.2 contributes to salt tolerance (Figure 1).

#### 4.5 | Contribution of a K<sup>+</sup> channel to plant salt tolerance

Salt stress does not only strongly increase shoot  $\text{Na}^+$  content but also leads to severe  $\text{K}^+$  deficiency (Hauser & Horie, 2010; Marschner, 2011). Since insuring efficient root  $\text{K}^+$  uptake from soil appears as the primary way to insure shoot  $\text{K}^+$  feeding, most studies aiming at identifying salt tolerance determinants among  $\text{K}^+$  transport systems have focused on root uptake systems. Exposure to high salinity can substantially depolarize root periphery cells and make passive  $\text{K}^+$  uptake through inwardly rectifying  $\text{K}^+$  channels thermodynamically impossible (Rubio et al., 2020). High-affinity HAK/KUP/KT transporters, expected to rely on pH gradients created by the  $\text{H}^+$ -ATPase pump to energize inward  $\text{K}^+$  fluxes through  $\text{H}^+$ - $\text{K}^+$  symport mechanism, are therefore considered as the main  $\text{K}^+$  transport systems taking part in root  $\text{K}^+$  uptake under high saline conditions. Several HAK transporters have been shown to be involved in root  $\text{K}^+$  uptake and thereby to contribute to plant salt tolerance: AtHAK5 in Arabidopsis (Nieves-Cordones et al., 2010), and OsHAK1, OsHAK5, OsHAK16 and OsHAK21 in rice (Chen et al., 2015; Feng et al., 2019; Shen et al., 2015; Yang et al., 2014). KO mutations in these different genes have been shown to result in reduced  $\text{K}^+$  uptake and root  $\text{K}^+$  content, and probably as a consequence, also in reduced  $\text{K}^+$  translocation to shoots and often reduced shoot  $\text{K}^+$  contents. Such defects could be observed upon salt stress but also in absence of saline treatment, and resulted in reduced plant growth in all conditions (Chen et al., 2015; Feng et al., 2019; Nieves-Cordones et al., 2010; Shen et al., 2015; Yang et al., 2014). Increased plant  $\text{Na}^+$  uptake was noted in some mutants (Shen et al., 2015), which could originate from higher root cell polarization (Nieves-Cordones et al., 2017). Also supporting the importance of HAK-mediated plant  $\text{K}^+$  uptake in salt tolerance, transcript level variations in the *OsHAK1* gene between rice subspecies have been found to contribute to the

difference in their salt tolerance (Chen et al., 2015).

It is also well known that transport systems from the H<sup>+</sup>/cation antiporter families, involved in K<sup>+</sup> and Na<sup>+</sup> intracellular compartmentalization, are major contributors to salt tolerance (van Zelm et al., 2020) through their roles in Na<sup>+</sup> compartmentalization and turgor regulation, but also through indirect contributions to K<sup>+</sup> homeostasis. For instance, increased activity (due to overexpression in transgenic plants) of the antiporter AtNHX1 from *Arabidopsis* or LeNHX2 from tomato has been shown to result in improved root K<sup>+</sup> uptake and higher K<sup>+</sup> contents in all tissues. Such effects, which are beneficial to salt tolerance, have been proposed to result from a decrease in cytosolic K<sup>+</sup> concentration that these transport systems would generate by compartmentalizing K<sup>+</sup>, which would lead to increased expression and/or activity of high affinity K<sup>+</sup> transporters involved in root K<sup>+</sup> uptake (Leidi et al., 2010; Huertas et al., 2013). Altogether, these studies provide evidence of strong interactions between K<sup>+</sup> uptake, compartmentalization and translocation to shoots.

Other K<sup>+</sup> transport-mediated mechanisms of plant salt tolerance and in particular mechanisms involving K<sup>+</sup> channels, were reported but have not yet been deciphered. Transcriptional regulation of a few K<sup>+</sup> channel genes, especially the strong up-regulation of the inward Shaker regulatory subunit *AtKCI* in leaves (Pilot et al., 2003), suggests a role of inward K<sup>+</sup> channels in salt tolerance, which has not been determined so far. Recently, reverse genetics analyses suggested that the weak inwardly rectifying channel gene *OsAKT2*, which is expressed in phloem, indirectly controls root Na<sup>+</sup> uptake, thereby contributing to maintenance of leaf K<sup>+</sup>/Na<sup>+</sup> homeostasis upon salt stress (Tian et al., 2021).

Here, we showed that KO mutation in the outward Shaker K<sup>+</sup> channel gene *OsK5.2* leads

to increased salt sensitivity. Lack of *OsK5.2* functional expression was found to result in impaired growth in plants subjected to saline conditions but not in plants grown in standard conditions (Figure 1), in contrast to what has been reported in KO mutant plants impaired in root  $K^+$  uptake, which mostly showed growth defects even in absence of salt stress (see above). *OsK5.2* is involved in control of transpiration rate and in  $K^+$  secretion into the xylem sap, and these two functions together underlie its contribution to salt tolerance. It is also worth to note that lack of *OsK5.2* activity results also in impaired net  $K^+$  uptake under saline conditions since both plant growth and whole plant  $K^+$  contents were lower in *osk5.2* mutant plants compared with the corresponding WT plants (Figure 1 and Figure 6). Together with the defects in  $K^+$  translocation to shoots under salt stress that have been reported in mutant plants impaired in root  $K^+$  uptake or in  $K^+$  intracellular compartmentalization (see above), the reduction in  $K^+$  uptake resulting from lack of *OsK5.2* channel activity provides evidence that the three functions, uptake, compartmentalization and translocation, are especially intensively coordinated under saline conditions. In conclusion, the present results highlight  $K^+$ -channel-mediated mechanisms of salt tolerance, and provide a new possible target for plant breeders towards the improvement of tolerance to salt stress in rice.

#### CONFLICT OF INTEREST

The authors declare no competing interests

#### AUTHOR CONTRIBUTIONS

A.-A.V., H.S., T.H.N. and J.Z. conceived the original research plans; A.-A.V., and D.T.L.



supervised the experiments; D.H., T.H.N. and J.Z. performed the experiments. A.-A.V., H.S., D.H., T.H.N. and J.Z. analyzed the data; A.-A.V., H.S. and J.Z. wrote the first draft of the manuscript. All authors contributed to the article and approved the submitted version.

#### DATA AVAILABILITY STATEMENT

The data that support the findings of this study are available from the corresponding author upon reasonable request

#### ORCID

D Hmidi: 0000-0002-5039-4357

DT Luu: 0000-0001-9765-2125

H Sentenac: 0000-0003-3641-4822

A-A Véry: 0000-0003-1961-5243

T-H Nguyen: 0000-0003-0347-2946

#### FIGURE LEGENDS

**FIGURE 1** Effect of *OsK5.2* loss of function on rice plant phenotype in control and salt stress conditions. Comparison of growth phenotype (a) and dry weight (b) between corresponding wild-type and *osk5.2* mutant plants (black and white bars, respectively) issued from ASJA08 or ASHF06 lines (left and right panels, respectively) under control and salt treatment. Six-week-old plants grown on hydroponic Yoshida medium were supplemented or not during

the last 7 days with 50 or 100 mM NaCl. Enlargements of leaf tips in 100 mM NaCl treatment are shown in inserts at the right for the four genotypes. Scale bars = 10 cm in (a). Means  $\pm$  SE,  $n = 10$  in (b). Double stars above the bars denote statistically significant differences between wild-type and *osk5.2* mutant plants (Student's  $t$  test,  $P \leq 0.01$ ).

**FIGURE 2** Effect of saline conditions on *OsK5.2* transcript levels in roots and leaves. Five-week-old rice plants cv Nipponbare hydroponically grown on Yoshida medium were supplemented or not with 50 mM NaCl for 14 days. Salt-treated plants were thereafter allowed to recover for 3 days on standard Yoshida medium. Expression data in roots (a) and leaves (b) were determined by real-time quantitative RT-PCR. Means  $\pm$  SE ( $n = 3$  biological replicates under salt treatment after 1, 3, 7 and 14 days and recovery, and  $n = 4$  under control treatment, one plant in control condition sampled at each time of salt treatment). Different letters indicate statistically significant differences (one-way ANOVA, Duncan's test,  $P \leq 0.05$ ).

**FIGURE 3** Steady-state transpiration rates in wild-type and *osk5.2* mutant plants under control and salt treatment conditions. Five-week-old plants hydroponically grown on Yoshida medium were supplemented or not with 50 mM NaCl for 14 days. Left and right panels: *osk5.2* mutant plants ( $\circ$ ) issued from ASJA08 or ASHF06 lines, respectively, and the corresponding wild-type plants ( $\bullet$ ). Transpiration was measured after 1, 3, 7 and 14 days of salt treatment (and at the same times for the plants maintained in control conditions). (a) and (b): steady-state transpiration rates in light (panel a;  $\sim 3$  h after light was switched on) and in

dark (panel b; ~5 h after light was switched off) conditions. Steady-state transpiration rate was determined by dividing the average plant rate of water loss at steady-state (means of 3 values) by the total surface of the plant aerial parts. Means  $\pm$  SE;  $n = 9$  under salt treatment after 1, 3, 7, 14 days, and  $n = 12$  under control conditions (3 plants in control condition sampled at each time of salt treatment). Single and double stars denote statistically significant differences between wild-type and *osk5.2* mutant plants (Student's *t* test,  $P \leq 0.05$  and  $P \leq 0.01$ , respectively).

**FIGURE 4** Xylem sap  $K^+$  and  $Na^+$  concentrations in wild-type and *osk5.2* mutant plants under control and salt treatment conditions. Five-week-old plants hydroponically grown on Yoshida medium were supplemented or not with 50 mM NaCl for 14 days. Left and right panels: *osk5.2* mutant plants ( $\circ$ ) issued from ASJA08 or ASHF06 lines, respectively, and the corresponding wild-type plants ( $\bullet$ ). Xylem sap exudates were collected after 1, 3, 7 and 14 days of salt treatment (and at the same times for the plants maintained in control conditions). (a) and (b):  $K^+$  (a) and  $Na^+$  (b) concentrations assayed in the collected xylem sap samples. (c)  $K^+/Na^+$  concentration ratios deduced from (a) and (b). Means  $\pm$  SE;  $n = 9$  under salt treatment after 1, 3, 7, 14 days, and  $n = 12$  under control conditions (3 plants in control condition sampled at each time of salt treatment). Single and double stars denote statistically significant differences between wild-type and *osk5.2* mutant plants (Student's *t* test,  $P \leq 0.05$  and  $P \leq 0.01$ , respectively).

**FIGURE 5**  $K^+$  and  $Na^+$  fluxes arriving at light in leaves of wild-type and *osk5.2* mutant plants

under control and salt treatment conditions. (a) and (b): normalized values (expressed per leaf surface) of  $K^+$  (a) and  $Na^+$  (b) fluxes assessed by multiplying the mean transpiration rates (data from Figure 3) by the corresponding  $K^+$  or  $Na^+$  concentrations in xylem sap (data from Figure 4). Left and right panels: *osk5.2* mutant plants ( $\circ$ ) and corresponding wild-type plants ( $\bullet$ ) issued from ASJA08 (left) or ASHF06 (right) lines. Means  $\pm$  SE;  $n = 9$  under salt treatment after 1, 3, 7 and 14 days, and  $n = 12$  under control conditions. See legends to Figures 3 and 4. Single and double stars denote statistically significant differences between the wild-type and *osk5.2* mutant plants (Student's  $t$  test,  $P \leq 0.05$  and  $P \leq 0.01$ , respectively).

**FIGURE 6** Root and Shoot  $K^+$  contents in wild-type and *osk5.2* mutant plants under control and salt treatment conditions. Five-week-old plants hydroponically grown on Yoshida medium were supplemented or not with 50 mM NaCl for 14 days. Left and right panels: *osk5.2* mutant plants ( $\circ$ ) and the corresponding wild-type plants ( $\bullet$ ) issued from ASJA08 (left) or ASHF06 (right) lines. Roots and shoots were sampled after 1, 3, 7 and 14 days of salt treatment (and at the same times for the plants maintained in control conditions). (a), (b) and (c):  $K^+$  contents in roots, shoots and whole plant, respectively. Means  $\pm$  SE;  $n = 15$  to 20 for the wild-type and mutant plants issued from the ASJA08 line, and 9 to 12 for wild-type and mutant plants issued from the ASHF06 line. Single and double stars denote statistically significant differences between the wild-type and *osk5.2* mutant plants (Student's  $t$  test,  $P \leq 0.05$  and  $P \leq 0.01$ , respectively).

**FIGURE 7** Root and Shoot  $Na^+$  contents in wild-type and *osk5.2* mutant plants under control

and salt treatment conditions. Same plants as in Figure 6. (a), (b) and (c): Na<sup>+</sup> contents in roots, shoots and whole plant, respectively. Means ± SE; *n* = 15 to 20 for the wild-type and mutant plants issued from the ASJA08 line, and 9 to 12 for wild-type and mutant plants issued from the ASHF06 line. Single and double stars denote statistically significant differences between the wild-type and *osk5.2* mutant plants (Student's *t* test, *P* ≤ 0.05 and *P* ≤ 0.01, respectively).

**FIGURE 8** Root and shoot K<sup>+</sup>/Na<sup>+</sup> content ratio in wild-type and *osk5.2* mutant plants under control and salt treatment conditions. Same experiment as in Figures 6 and 7. K<sup>+</sup>/Na<sup>+</sup> content ratio: K<sup>+</sup> content from Figure 6 divided by the corresponding Na<sup>+</sup> content from Figure 7. (a) and (b): K<sup>+</sup>/Na<sup>+</sup> content ratio in roots and shoots. Left and right panels: *osk5.2* mutant plants (○) and corresponding wild-type plants (●) issued from ASJA08 (left) or ASHF06 (right) lines. Means ± SE; *n* = 15 to 20 for the wild-type and mutant plants issued from the ASJA08 line, and 9 to 12 for wild-type and mutant plants issued from the ASHF06 line. Single and double stars denote statistically significant differences between the wild-type and *osk5.2* mutant plants (Student's *t* test, *P* ≤ 0.05 and *P* ≤ 0.01, respectively).

**REFERENCES**

Campbell, M. T., Bandillo, N., Al Shiblawi, F. R. A., Sharma, S., Liu, K., Du, Q., . . . Walia, H. (2017). Allelic variants of OsHKT1;1 underlie the divergence between indica and japonica subspecies of rice (*Oryza sativa*) for root sodium content. *PLoS Genetics*, 13(6),

e1006823.

Chen, G., Hu, Q., Luo, L., Yang, T., Zhang, S., Hu, Y., . . . Xu, G. (2015). Rice potassium transporter OsHAK1 is essential for maintaining potassium-mediated growth and functions in salt tolerance over low and high potassium concentration ranges. *Plant, Cell & Environment*, 38(12), 2747-2765.

Faiyue, B., Al-Azzawi, M. J., & Flowers, T. J. (2012). A new screening technique for salinity resistance in rice (*Oryza sativa* L.) seedlings using bypass flow. *Plant, Cell & Environment*, 35(6), 1099-1108.

Feng, H., Tang, Q., Cai, J., Xu, B., Xu, G., & Yu, L. (2019). Rice OsHAK16 functions in potassium uptake and translocation in shoot, maintaining potassium homeostasis and salt tolerance. *Planta*, 250(2), 549-561.

Flam-Shepherd, R., Huynh, W. Q., Coskun, D., Hamam, A. M., Britto, D. T. & Kronzucker, H. J. (2018). Membrane fluxes, bypass flows, and sodium stress in rice: the influence of silicon. *Journal of Experimental Botany*, 69(7), 1679–1692.

Fricke, W., Akhiyarova, G., Wei, W., Alexandersson, E., Miller, A., Kjellbom, P. O., . . . Volkov, V. (2006). The short-term growth response to salt of the developing barley leaf. *Journal of Experimental Botany*, 57(5), 1079-1095.

Gaymard, F., Pilot, G., Lacombe, B., Bouchez, D., Bruneau, D., Boucherez, J., . . . Sentenac, H. (1998). Identification and disruption of a plant shaker-like outward channel involved in K<sup>+</sup> release into the xylem sap. *Cell*, 94(5), 647-655.

Hauser, F., & Horie, T. (2010). A conserved primary salt tolerance mechanism mediated by HKT transporters: a mechanism for sodium exclusion and maintenance of high K<sup>+</sup>/Na<sup>+</sup>

- ratio in leaves during salinity stress. *Plant, cell & environment*, 33(4), 552-565.
- Hedrich, R., & Shabala, S. (2018). Stomata in a saline world. *Current opinion in plant biology*, 46, 87-95.
- Hosy, E., Vavasseur, A., Mouline, K., Dreyer, I., Gaymard, F., Porée, F., . . . Sentenac, H. (2003). The Arabidopsis outward K<sup>+</sup> channel GORK is involved in regulation of stomatal movements and plant transpiration. *Proceedings of the National Academy of Sciences USA*, 100(9), 5549-5554.
- Huang, X.-Y., Chao, D.-Y., Gao, J.-P., Zhu, M.-Z., Shi, M., & Lin, H.-X. (2009). A previously unknown zinc finger protein, DST, regulates drought and salt tolerance in rice via stomatal aperture control. *Genes & Development*, 23, 1805–1817.
- Huertas, R., Rubio, L., Cagnac, O., García-Sánchez, M. J., Alché, J. de D, Venema, K., ... Rodríguez-Rosales, M. P. (2013). The K<sup>+</sup>/H<sup>+</sup> antiporter LeNHX2 increases salt tolerance by improving K<sup>+</sup> homeostasis in transgenic tomato. *Plant, Cell and Environment*, 36, 2135–2149.
- Ismail, A. M., & Horie, T. (2017). Genomics, physiology, and molecular breeding approaches for improving salt tolerance. *Annual Review of Plant Biology*, 68, 405-434.
- Jayakannan, M., Bose, J., Babourina, O., Rengel, Z., & Shabala, S. (2013). Salicylic acid improves salinity tolerance in Arabidopsis by restoring membrane potential and preventing salt-induced K<sup>+</sup> loss via a GORK channel. *Journal of Experimental Botany*, 64(8), 2255-2268.
- Jeschke, W. D. (1984). Effects of transpiration on potassium and sodium fluxes in root cells and the regulation of ion distribution between roots and shoots of barley seedlings.

*Journal of plant physiology*, 117(3), 267-285.

Jezek, M., & Blatt, M. R. (2017). The membrane transport system of the guard cell and its integration for stomatal dynamics. *Plant Physiology*, 174(2), 487-519.

Khan, I., Mohamed, S., Regnault, T., Mieulet, D., Guiderdoni, E., Sentenac, H., & Véry, A.-A. (2020). Constitutive contribution by the rice OsHKT1;4 Na<sup>+</sup> transporter to xylem sap desalinization and low Na<sup>+</sup> accumulation in young leaves under low as high external Na<sup>+</sup> conditions. *Frontiers in Plant Science*, 11, 1130.

Lebaudy, A., Vavasseur, A., Hosy, E., Dreyer, I., Leonhardt, N., Thibaud, J.-B., Véry, A.-A., Simonneau, T., & Sentenac, H. (2008). Plant adaptation to fluctuating environment and biomass production are strongly dependent on guard cell potassium channels. *Proceedings of the National Academy of Sciences, USA*, 105, 5271–5276.

Leidi, E. O., Barragán, V., Rubio, L., El-Hamdaoui, A., Ruiz, M. T., Cubero, B., ... Pardo, J. M. (2010). The AtNHX1 exchanger mediates potassium compartmentation in vacuoles of transgenic tomato. *The Plant Journal*, 61(3), 495-506.

Maathuis, F. J., & Amtmann, A. (1999). K<sup>+</sup> nutrition and Na<sup>+</sup> toxicity: the basis of cellular K<sup>+</sup>/Na<sup>+</sup> ratios. *Annals of Botany*, 84(2), 123-133.

Maathuis, F. J., Ahmad, I., & Patishtan, J. (2014). Regulation of Na<sup>+</sup> fluxes in plants. *Frontiers in plant science*, 5, 467.

Marschner, H. (2011). *Marschner's mineral nutrition of higher plants*: Academic press.

Matsuda, S., Takano, S., Sato, M., Furukawa, K., Nagasawa, H., Yoshikawa, S., ...Kato, K. (2016). Rice stomatal closure requires guard cell plasma membrane ATP-binding cassette transporter RCN1/OsABCG5. *Molecular Plant*, 9, 417–427.



- 706 Mian, A., Oomen, R. J. F. J., Isayenkov, S., Sentenac, H., Maathuis, F. J. M., & Véry, A.-A.  
707 (2011) Over-expression of an Na<sup>+</sup>- and K<sup>+</sup>-permeable HKT transporter in barley  
708 improves salt tolerance. *The Plant Journal*, 68, 468–479.
- 709 Munns, R., & Tester, M. (2008). Mechanisms of salinity tolerance. *Annu. Rev. Plant Biol.*, 59,  
710 651-681.
- 711 Nguyen, T. H., Huang, S., Meynard, D., Chaine, C., Michel, R., Roelfsema, M. R. G., . . .  
712 Véry, A.-A. (2017). A dual role for the OsK5.2 ion channel in stomatal movements and  
713 K<sup>+</sup> loading into xylem sap. *Plant Physiology*, 174(4), 2409-2418.
- 714 Nieves-Cordones, M., Alemán, F., Martínez, V. & Rubio, F. (2010). The Arabidopsis  
715 thaliana HAK5 K<sup>+</sup> transporter is required for plant growth and K<sup>+</sup> acquisition from low  
716 K<sup>+</sup> solutions under saline conditions. *Molecular Plant*, 3(2), 326–333.
- 717 Nieves-Cordones, M., Al Shiblawi, F. R., & Sentenac, H. (2016). Roles and transport of  
718 sodium and potassium in plants. In *The alkali metal ions: Their role for life* (pp.  
719 291-324). Springer, Cham.
- 720 Nieves-Cordones, M., Mohamed, S., Tanoi, K., Kobayashi, N., Takagi, K., Vernet, A., ...  
721 Véry, A.-A. (2017). Production of low-Cs<sup>+</sup> rice plants by inactivation of the K<sup>+</sup>  
722 transporter OsHAK1 with the CRISPR-Cas system. *The Plant Journal*, 92, 43-56.
- 723 Pandey, S., Zhang, W., & Assmann, S. M. (2007). Roles of ion channels and transporters in  
724 guard cell signal transduction. *FEBS letters*, 581(12), 2325-2336.
- 725 Pilot, G., Gaymard, F., Mouline, K., Chérel, I., & Sentenac, H. (2003). Regulated expression  
726 of Arabidopsis Shaker K<sup>+</sup> channel genes involved in K<sup>+</sup> uptake and distribution in the  
727 plant. *Plant Molecular Biology*, 51, 773–787.

- Ren, Z.-H., Gao, J.-P., Li, L.-G., Cai, X.-L., Huang, W., Chao, D.-Y., . . . Lin, H.-X. (2005). A rice quantitative trait locus for salt tolerance encodes a sodium transporter. *Nature genetics*, 37(10), 1141-1146.
- Robinson, M. F., Véry, A.-A., Sanders, D., & Mansfield, T. A. (1997). How can stomata contribute to salt tolerance? *Annals of botany*, 80(4), 387-393.
- Rubio, F., Nieves-Cordones, M., Horie, T., & Shabala, S. (2020). Doing ‘business as usual’ comes with a cost: evaluating energy cost of maintaining plant intracellular  $K^+$  homeostasis under saline conditions. *New Phytologist*, 225(3), 1097-1104.
- Schroeder, J. I., Allen, G. J., Hugouvieux, V., Kwak, J. M., & Waner, D. (2001). Guard cell signal transduction. *Annual review of plant biology*, 52(1), 627-658.
- Shen, Y., Shen, L., Shen, Z., Jing, W., Ge, H., Zhao, J., & Zhang, W. (2015). The potassium transporter OsHAK21 functions in the maintenance of ion homeostasis and tolerance to salt stress in rice. *Plant, Cell & Environment*, 38(12), 2766-2779.
- Suzuki, K., Yamaji, N., Costa, A., Okuma, E., Kobayashi, N. I., Kashiwagi, T., . . . Horie, T. (2016). OsHKT1;4-mediated  $Na^+$  transport in stems contributes to  $Na^+$  exclusion from leaf blades of rice at the reproductive growth stage upon salt stress. *BMC Plant Biology*, 16(1), 1-15.
- Thiel, G. & Blatt, M. R. (1991). The mechanism of ion permeation through  $K^+$  channels of stomatal guard cells: voltage-dependent block by  $Na^+$ . *Journal of Plant Physiology* 138, 326–334.
- Tian, Q., Shen, L., Luan, J., Zhou, Z., Guo, D., Shen, Y., Jing, W., Zhang, B., Zhang, Q., & Zhang, W. (2021). Rice shaker potassium channel OsAKT2 positively regulates salt

tolerance and grain yield by mediating K<sup>+</sup> redistribution. *Plant, Cell & Environment*,  
1–15. <https://doi.org/10.1111/pce.14101>

Véry, A. A., Robinson, M. F., Mansfield, T. A., & Sanders, D. (1998). Guard cell cation  
channels are involved in Na<sup>+</sup>-induced stomatal closure in a halophyte. *The Plant  
Journal*, 14(5), 509-521.

Véry, A. A., Nieves-Cordones, M., Daly, M., Khan, I., Fizames, C., & Sentenac, H. (2014).  
Molecular biology of K<sup>+</sup> transport across the plant cell membrane: what do we learn  
from comparison between plant species?. *Journal of plant physiology*, 171(9), 748-769.

Wang, R., Jing, W., Xiao, L., Jin, Y., Shen, L., & Zhang, W. (2015). The rice high-affinity  
potassium transporter1;1 is involved in salt tolerance and regulated by an MYB-type  
transcription factor. *Plant Physiology*, 168(3), 1076-1090.

Wu, H., Zhang, X., Giraldo, J. P., & Shabala, S. (2018). It is not all about sodium: revealing  
tissue specificity and signalling roles of potassium in plant responses to salt stress. *Plant  
and soil*, 431(1), 1-17.

Wegner, L. H., & de Boer, A. H. (1997). Properties of two outward-rectifying channels in  
root xylem parenchyma cells suggest a role in K<sup>+</sup> homeostasis and long-distance  
signaling. *Plant Physiology*, 115(4), 1707-1719.

Yang, T., Zhang, S., Hu, Y., Wu, F., Hu, Q., Chen, G., . . . Xu, G. (2014). The role of a  
potassium transporter OsHAK5 in potassium acquisition and transport from roots to  
shoots in rice at low potassium supply levels. *Plant Physiology*, 166(2), 945-959.

Yeo, A. (1998). Molecular biology of salt tolerance in the context of whole-plant physiology.  
*Journal of Experimental Botany*, 49(323), 915-929.

van Zelm, E., Zhang, Y., & Testerink, C. (2020). Salt tolerance mechanisms of plants. *Annual Review of Plant Biology*, 71, 403-433.

Zeng, L., & Shannon, M. C. (2000). Salinity effects on seedling growth and yield components of rice. *Crop Science*, 40(4), 996-1003.

#### SUPPORTING INFORMATION

Additional supporting information may be found online in the Supporting Information section at the end of this article.


**Table S1:** Primers used for qRT-PCR experiments

**Figure S1** Effect of salt treatment on shoot and root length of wild-type (black bars) and *osk5.2* mutant (white bars) plants issued from ASJA08 (top) or ASHF06 (bottom) lines.


**Figure S2** Shoot and root water contents of wild-type and *osk5.2* mutant plants under control and salt treatment conditions.

**FIGURE S3** Scanning electron micrographs showing stomatal shape on the abaxial flag leaf side from *osk5.2* mutant (right column) and corresponding wild-type (left column) plants issued from ASJA08 (top) or ASHF06 (bottom) lines.


**Figure S4** The ionic composition of the xylem sap does not reflect the K<sup>+</sup> and Na<sup>+</sup> relative contents of the roots.




**FIGURE 1** Effect of *Osk5.2* loss of function on rice plant phenotype in control and salt stress conditions. Comparison of growth phenotype (a) and dry weight (b) between corresponding wild-type and *osk5.2* mutant plants (black and white bars, respectively) issued from ASJA08 or ASHF06 lines (left and right panels, respectively) under control and salt treatment. Six-week-old plants grown on hydroponic Yoshida medium were supplemented or not during the last 7 days with 50 or 100 mM NaCl. Enlargements of leaf tips in 100 mM NaCl treatment are shown in inserts at the right for the four genotypes. Scale bars = 10 cm in (a). Means  $\pm$  SE,  $n = 10$  in (b). Double stars above the bars denote statistically significant differences between wild-type and *osk5.2* mutant plants (Student's  $t$  test,  $P \leq 0.01$ ).



**FIGURE 2** Effect of saline conditions on *OsK5.2* transcript levels in roots and leaves. Five-week-old rice plants cv Nipponbare hydroponically grown on Yoshida medium were supplemented or not with 50 mM NaCl for 14 days. Salt-treated plants were thereafter allowed to recover for 3 days on standard Yoshida medium. Expression data in roots (a) and leaves (b) were determined by real-time quantitative RT-PCR. Means  $\pm$  SE ( $n = 3$  biological replicates under salt treatment after 1, 3, 7 and 14 days and recovery, and  $n = 4$  under control treatment sampled at each time of salt treatment). Different letters indicate statistically significant differences (One-way ANOVA, Duncan's test,  $P \leq 0.05$ ).




**FIGURE 3** Steady-state transpiration rates in wild-type and *osk5.2* mutant plants under control and salt treatment conditions. Five-week-old plants hydroponically grown on Yoshida medium were supplemented or not with 50 mM NaCl for 14 days. Left and right panels: *osk5.2* mutant plants (○) issued from ASJA08 or ASHF06 lines, respectively, and the corresponding wild-type plants (●). Transpiration was measured after 1, 3, 7 and 14 days of salt treatment (and at the same times for the plants maintained in control conditions). (a) and (b): steady-state transpiration rates in light (panel a; ~3 h after light was switched on) and in dark (panel b; ~5 h after light was switched off) conditions. Steady-state transpiration rate was determined by dividing the average plant rate of water loss at steady-state (means of 3 values) by the total surface of the plant aerial parts. Means  $\pm$  SE;  $n = 9$  under salt treatment after 1, 3, 7, 14 days, and  $n = 12$  under control conditions (3 plants in control condition sampled at each time of salt treatment). Single and double stars denote statistically significant differences between wild-type and *osk5.2* mutant plants (Student's  $t$  test,  $P \leq 0.05$  and  $P \leq 0.01$ , respectively).



**FIGURE 4** Xylem sap  $K^+$  and  $Na^+$  concentrations in wild-type and *osk5.2* mutant plants under control and salt treatment conditions. Five-week-old plants hydroponically grown on Yoshida medium were supplemented or not with 50 mM NaCl for 14 days. Left and right panels: *osk5.2* mutant plants (○) issued from ASJA08 or ASHF06 lines, respectively, and the corresponding wild-type plants (●). Xylem sap exudates were collected after 1, 3, 7 and 14 days of salt treatment (and at the same times for the plants maintained in control conditions). (a) and (b):  $K^+$  (a) and  $Na^+$  (b) concentrations assayed in the collected xylem sap samples. (c)  $K^+/Na^+$  concentration ratios deduced from (a) and (b). Means  $\pm$  SE;  $n = 9$  under salt treatment after 1, 3, 7, 14 days, and  $n = 12$  under control conditions (3 plants in control condition sampled at each time of salt treatment). Single and double stars denote statistically significant differences between wild-type and *osk5.2* mutant plants (Student's  $t$  test,  $P \leq 0.05$  and  $P \leq 0.01$ , respectively).







**FIGURE 5** K<sup>+</sup> and Na<sup>+</sup> fluxes arriving at light in leaves of wild-type and *osk5.2* mutant plants under control and salt treatment conditions. (a) and (b): normalized values (expressed per leaf surface) of K<sup>+</sup> (a) and Na<sup>+</sup> (b) fluxes assessed by multiplying the mean transpiration rates (data from Figure 3) by the corresponding K<sup>+</sup> or Na<sup>+</sup> concentrations in xylem sap (data from Figure 4). Left and right panels: *osk5.2* mutant plants (○) and corresponding wild-type plants (●) issued from ASJA08 (left) or ASHF06 (right) lines. Means ± SE; *n* = 9 under salt treatment after 1, 3, 7 and 14 days, and *n* = 12 under control conditions. See legends to Figures 3 and 4. Single and double stars denote statistically significant differences between the wild-type and *osk5.2* mutant plants (Student's *t* test, *P* ≤ 0.05 and *P* ≤ 0.01, respectively).

## ASJA08 line


## ASHF06 line



**FIGURE 6** Root and Shoot  $K^+$  contents in wild-type and *osk5.2* mutant plants under control and salt treatment conditions. Five-week-old plants hydroponically grown on Yoshida medium were supplemented or not with 50 mM NaCl for 14 days. Left and right panels: *osk5.2* mutant plants ( $\circ$ ) and the corresponding wild-type plants ( $\bullet$ ) issued from ASJA08 (left) or ASHF06 (right) lines. Roots and shoots were sampled after 1, 3, 7 and 14 days of salt treatment (and at the same times for the plants maintained in control conditions). (a), (b) and (c):  $K^+$  contents in roots, shoots and whole plant, respectively. Means  $\pm$  SE;  $n = 15$  to 20 for the wild-type and mutant plants issued from the ASJA08 line, and 9 to 12 for wild-type and mutant plants issued from the ASHF06 line. Single and double stars denote statistically significant differences between the wild-type and *osk5.2* mutant plants (Student's  $t$  test,  $P \leq 0.05$  and  $P \leq 0.01$ , respectively).



**FIGURE 7** Root and Shoot Na<sup>+</sup> contents in wild-type and *osk5.2* mutant plants under control and salt treatment conditions. Same plants as in Figure 6. (a), (b) and (c): Na<sup>+</sup> contents in roots, shoots and whole plant, respectively. Means  $\pm$  SE;  $n = 15$  to 20 for the wild-type and mutant plants issued from the ASJA08 line, and 9 to 12 for wild-type and mutant plants issued from the ASHF06 line. Single and double stars denote statistically significant differences between the wild-type and *osk5.2* mutant plants (Student's  $t$  test,  $P \leq 0.05$  and  $P \leq 0.01$ , respectively).




**FIGURE 8** Root and shoot  $K^+/Na^+$  content ratio in wild-type and *osk5.2* mutant plants under control and salt treatment conditions. Same experiment as in Figures 6 and 7.  $K^+/Na^+$  content ratio:  $K^+$  content from Figure 6 divided by the corresponding  $Na^+$  content from Figure 7. (a) and (b):  $K^+/Na^+$  content ratio in roots and shoots. Left and right panels: *osk5.2* mutant plants ( $\circ$ ) and corresponding wild-type plants ( $\bullet$ ) issued from ASJA08 (left) or ASHF06 (right) lines. Means  $\pm$  SE;  $n = 15$  to 20 for the wild-type and mutant plants issued from the ASJA08 line, and 9 to 12 for wild-type and mutant plants issued from the ASHF06 line. Single and double stars denote statistically significant differences between the wild-type and *osk5.2* mutant plants (Student's  $t$  test,  $P \leq 0.05$  and  $P \leq 0.01$ , respectively).


**Table S1:** Primers used for qRT-PCR experiments.

Gene	Primer name	Sequence (5'-3')	Amplicon length (bp)
<i>OsK5.2</i>	qPCR-OsK52-F358	TTTGGCTTCTTCAGGGGGCT	470 *
	qPCR-OsK52-R808	TCTCCCAACTGCAAGCTCCC	
	qPCR-OsK52-F467	CTGACACCTACCGCATGGTT	D-164
	qPCR-OsK52-R611	TCGTGTCAACCGAATCCACA	
<i>SMT3</i>	SMT3B303F	GGGAGGAGGACAAGAAGC	273 *
	SMT3B303R	CTCCAGTCTGGTGGAGCAT	
	QRT94SMT3-F	CCTCAAGGTCAAGGGACAGG	D-94
	QRT94SMT3-R	ACGGTCACAATAGGCGTTCA	
<i>Tip41</i>	Tip41B389F	TGGTTTTTGGGGAGAGTTTC	389 *
	Tip41B389R	TGAAATGCCATTATCGGCTA	
	QRT101Tip41-F	TGGGAGTGATGCTTTGGTTC	D-101
	QRT101Tip41-R	CAAGGTCAATCCGATCCTCA	
<i>eEF-1-beta2</i>	eEFB2Pcs	TGGTGAGGAGACTGAAGAGG	418 *
	eEFB2Pcas	CTTCCGGATTTTTCTTTTATC	
	qPCR-eEF1b2-F413	TGGGAAGTCCTCAGTGTTGC	D-136
	qPCR-eEF1b2-R549	GTAACCAACCGGGACAAGCT	

“D”, pair of primers for transcript level quantification, that anneal on cDNA fragments derived from two exons. “\*” refers to big amplicons that were used as a template for the amplification of calibration standards.




**Figure S1** Effect of salt treatment on shoot and root length of wild-type (black bars) and *osk5.2* mutant (white bars) plants issued from ASJA08 (top) or ASHF06 (bottom) lines. Five-week-old rice plants hydroponically grown on Yoshida medium were supplemented or not with 50 or 100 mM NaCl for 7 days. Means  $\pm$  SE ( $n = 10$  plants). One or 2 stars above bars corresponding to mutant plants indicate statistically significant differences with respect to the corresponding wild-type plants (Student's  $t$  test,  $P < 0.05$  and  $< 0.01$ , respectively).




**Figure S2** Shoot and root water contents of wild-type and *osk5.2* mutant plants under control and salt treatment conditions. Five-week-old rice plants hydroponically grown on Yoshida medium were supplemented or not with 50 or 100 mM NaCl for 7 days. Top and bottom panels: *osk5.2* mutant plants (white bars) and corresponding wild-type plants (black bars) issued from ASJA08 (top) or ASHF06 (bottom) lines. Means  $\pm$  SE ( $n = 10$ ). Absence of star above the bars indicates no statistically significant difference between the wild-type and *osk5.2* mutant plants (Student's  $t$  test,  $P > 0.05$ ).

WT

ASJA08 line




ASHF06 line



**FIGURE S3** Scanning electron micrographs showing stomatal shape on the abaxial flag leaf side from *osk5.2* mutant (right column) and corresponding wild-type (left column) plants issued from ASJA08 (top) or ASHF06 (bottom) lines. Plants were grown in control conditions (Yoshida hydroponic medium). A ca. 1 cm-length portion of leaf, cut from an illuminated plant, was stuck on a thin layer of OCT cryo-embedding matrix (Carlroth) and quickly frozen at -14°C. Observations were performed on 30 stomata from 3 plants per genotype using a benchtop Phenom Pro X scanning electron microscope (Phenom World, Eindhoven, The Netherlands) using a backscattered electron detector with an acceleration voltage of 15 kV in image mode. GC: guard cell; SC: subsidiary cell.





**FIGURE S4** The ionic composition of the xylem sap does not reflect the K<sup>+</sup> and Na<sup>+</sup> relative contents of the roots. Plants were subjected to a saline treatments (50 mM NaCl for 2 weeks; x axis). Y axis: xylem sap K<sup>+</sup>/Na<sup>+</sup> concentration ratio (data from Figure 4c) reported to the root K<sup>+</sup>/Na<sup>+</sup> content ratio (data from Figure 8a). Left and right panels: *osk5.2* mutant plants (○) and corresponding wild-type plants (●) issued from ASJA08 (left) or ASHF06 (right) lines. Means ± SE; *n* = 9 under salt treatment after 1, 3, 7 and 14 days, and *n* = 12 under control conditions. See legends to Figures 4 and 8.