



HAL
open science

A Perspective on Plant Phenomics: Coupling Deep Learning and Near-Infrared Spectroscopy

François Vasseur, Denis Cornet, Grégory Beurier, Julie Messier, Lauriane Rouan, Justine Bresson, Martin Ecartot, Mark Stahl, Simon Heumos, Marianne Gérard, et al.

► To cite this version:

François Vasseur, Denis Cornet, Grégory Beurier, Julie Messier, Lauriane Rouan, et al.. A Perspective on Plant Phenomics: Coupling Deep Learning and Near-Infrared Spectroscopy. *Frontiers in Plant Science*, 2022, 13, 10.3389/fpls.2022.836488 . hal-03683630

HAL Id: hal-03683630

<https://hal.inrae.fr/hal-03683630>

Submitted on 31 May 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License



A Perspective on Plant Phenomics: Coupling Deep Learning and Near-Infrared Spectroscopy

François Vasseur^{1*}, Denis Cornet^{2,3}, Grégory Beurier^{2,3}, Julie Messier⁴, Lauriane Rouan^{2,3}, Justine Bresson¹, Martin Ecartot³, Mark Stahl⁵, Simon Heumos^{6,7}, Marianne Gérard¹, Hans Reijnen¹, Pascal Tillard⁸, Benoît Lacombe⁸, Amélie Emanuel^{1,8}, Justine Floret^{1,9}, Aurélien Estarague¹, Stefania Przybylska¹, Kevin Sartori¹, Lauren M. Gillespie¹, Etienne Baron¹, Elena Kazakou¹⁰, Denis Vile⁹ and Cyrille Violle¹

¹CEFE, Univ Montpellier, CNRS, EPHE, IRD, Montpellier, France, ²CIRAD, UMR AGAP Institut, Montpellier, France, ³UMR AGAP Institut, Univ Montpellier, CIRAD, INRAE, Institut Agro, Montpellier, France, ⁴Department of Biology, University of Waterloo, Waterloo, ON, Canada, ⁵Center for Plant Molecular Biology (ZMBP), University of Tübingen, Tübingen, Germany, ⁶Quantitative Biology Center (QBiC), University of Tübingen, Quantitative Biology Center (QBiC), University of Tübingen, Germany, ⁷Biomedical Data Science, Department of Computer Science, University of Tübingen, Tübingen, Germany, ⁸BPMP, Univ Montpellier, CNRS, INRAE, Montpellier, France, ⁹LEPSE, Univ Montpellier, INRAE, Institut Agro, Montpellier, France, ¹⁰CEFE, Univ Montpellier, CNRS, EPHE, Institut Agro, IRD, Université Paul Valéry Montpellier, Montpellier, France

OPEN ACCESS

Edited by:

Alexis Joly,
Research Center Inria Sophia
Antipolis Méditerranée, France

Reviewed by:

Roxana Vidican,
University of Agricultural Sciences
and Veterinary Medicine of Cluj-
Napoca, Romania
Satoru Tsuchikawa,
Nagoya University, Japan

*Correspondence:

François Vasseur
francois.vasseur@cefe.cnrs.fr

Specialty section:

This article was submitted to
Technical Advances in Plant Science,
a section of the journal
Frontiers in Plant Science

Received: 03 January 2022

Accepted: 09 March 2022

Published: 20 May 2022

Citation:

Vasseur F, Cornet D, Beurier G,
Messier J, Rouan L, Bresson J,
Ecartot M, Stahl M, Heumos S,
Gérard M, Reijnen H, Tillard P,
Lacombe B, Emanuel A, Floret J,
Estarague A, Przybylska S, Sartori K,
Gillespie LM, Baron E, Kazakou E,
Vile D and Violle C (2022) A
Perspective on Plant Phenomics:
Coupling Deep Learning and Near-
Infrared Spectroscopy.
Front. Plant Sci. 13:836488.
doi: 10.3389/fpls.2022.836488

The trait-based approach in plant ecology aims at understanding and classifying the diversity of ecological strategies by comparing plant morphology and physiology across organisms. The major drawback of the approach is that the time and financial cost of measuring the traits on many individuals and environments can be prohibitive. We show that combining near-infrared spectroscopy (NIRS) with deep learning resolves this limitation by quickly, non-destructively, and accurately measuring a suite of traits, including plant morphology, chemistry, and metabolism. Such an approach also allows to position plants within the well-known CSR triangle that depicts the diversity of plant ecological strategies. The processing of NIRS through deep learning identifies the effect of growth conditions on trait values, an issue that plagues traditional statistical approaches. Together, the coupling of NIRS and deep learning is a promising high-throughput approach to capture a range of ecological information on plant diversity and functioning and can accelerate the creation of extensive trait databases.

Keywords: *Arabidopsis thaliana*, near-infrared spectroscopy (NIRS), multivariate analysis, machine learning, functional traits, metabolomics, trait-based ecology

INTRODUCTION

In trait-based ecology, the comparison of plant phenotype across multiple species aims at identifying general trends of variation to describe the biodiversity of plant forms and functions (Grime, 1988; Keddy, 1992; Díaz et al., 2016; Garnier et al., 2016). Ecological strategies are characterized qualitatively and quantitatively from the measurement of key functional traits, i.e., morphological, physiological, and phenological parameters that determine plant growth and reproduction (Violle et al., 2007). However, our understanding of plant diversity with comparative approaches is impeded by three main limitations. First, measuring the traits that describe ecological strategies on many individuals remains laborious. Second, intraspecific trait variability and plasticity to

the environment still remain largely unconnected to traditional cross-species studies (but see Albert et al., 2010; Anderegg et al., 2018). Third, we need to clarify if and how plant (“soft”) traits used classically to describe ecological strategies are connected to plant metabolism and physiology (“hard” traits).

The development of near-infrared spectroscopy (NIRS) has provided a unique, fast, and reliable tool enabling the collection of a myriad of traits non-destructively (Foley et al., 1998; Cozzolino et al., 2001; Pasquini, 2018; Silva-Perez et al., 2018). NIRS measures the light reflected from a sample after irradiating it with wavelengths ranging from visible (VIS, 400–700 nm), near-infrared (NIR, 700–1,100 nm), to shortwave infrared (SWIR, 1,100–2,500 nm). This provides a signature of the physical and chemical characteristics of the sample (Box 1). NIRS has been widely used for determining chemical traits in various fields. For instance, it is extensively used to characterize chemical products in pharmaceutical, agricultural, and food sectors (Shepherd and Walsh, 2007; Wójcicki, 2015; Biancolillo and Marini, 2018; Pasquini, 2018). In plant science, NIRS takes an increasingly important place as a high-throughput, cost-efficient method for the characterization of biodiversity (Arslan et al., 2018; Silva-Perez et al., 2018; Burnett et al., 2021; Kothari et al., 2021). For instance, it is widely used to predict differences in leaf palatability, digestibility, and decomposability—through lignin and fiber content—between species (Birth and Hecht, 1987; Andrés et al., 2005). The advantages of this method are numerous: spectral measurements are extremely rapid, taking only a few seconds, a single spectral measurement simultaneously captures multiple diverse plant traits (Petit Bon et al., 2020), minimal or no sample preparation is required, and the measurements are non-destructive which allows to track trait changes over time and avoids interfering with the organism.

BOX 1 | Principle of near-infrared spectroscopy (NIRS) for plant characterization

The leaf spectral reflectance is based on the low reflectivity in the visible part of the spectrum (400–700 nm), due to a strong absorption by photosynthetic pigments, and the high reflectivity in the near infrared (700–1,100 nm) produced by a high scattering of light by the leaf mesophyll tissues (Knippling, 1970). For instance, in the SWIR part of the spectrum (1,100–2,500 nm), the reflectance intensity is affected by the water, cellulose, protein, and lignin content of plant tissues (Rascher et al., 2010). Healthy leaves emit radiation in the thermal infrared band ($\approx 10\mu\text{m}$) according to their temperature, because of their high water content (emissivity between 0.97 and 0.99). The leaves appear green because the green light band (550 nm) is reflected relatively efficiently when compared with the blue, yellow, and red bands, which are absorbed by photoactive pigments. This absorption at different wavelength produces a spectrum of light reflectance (Figure 1), which can be treated as a “signal” of the leaf physical and chemical properties.

The physical association between leaf properties and light reflectance is particularly useful to investigate leaf composition, functioning, and diversity. Different leaves will have different spectral signatures depending on their structure and chemical composition. For example, leaf nitrogen concentrations are associated with wavelengths absorbed by chlorophyll a and b in the visible part of the spectrum (400–700 nm), the spectral red edge (700–760 nm), and proteins in the SWIR (1,300–2,500 nm; Gitelson and Merzlyak, 1997; Kokaly, 2001). In the SWIR (SWIR; 700–1,300 nm), structures such as palisade cell density are important determinants of the spectral reflectance because of the very low effective photon penetration distance at these wavelengths.

While NIRS data are simple to acquire and rapidly generate a very large amount of information, they also require extensive post-processing, *via* chemiometric and multivariate statistical

BOX 2 | The promise of deep learning to analyze NIRS

Chemometrics, the science of extracting information from chemical systems, faces multiple challenges while studying the link between samples near infrared spectra and desired traits. Partial least square regression (PLSR), the reference method, is well designed to address many of these but still exhibit some weakness that native properties and tools associated with deep learning may allow to tackle.

First, the inherent and unwanted variability associated with the spectrometric measure result in a highly noisy signal (e.g., environmental, machine, or sample variability). To deal with it, reference methods rely on pretreatment (i.e., filtering) of the signal and removal of the spectral outliers. Pretreatment works by removing noise and linearizing the response of a variable. As the number of possible pretreatments increases with research progress, finding the optimal solution empirically becomes more and more a challenge. Moreover, pretreatment development is optimized for its suitability to filter spectra in combination with a reference model (i.e., PLSR in most situations). This may result in suboptimal solution and loss of information. Conversely, some deep learning algorithms are particularly efficient in filtering input signals. Cui and Fearn (2018) illustrated how the convolutional layer can continually tune the variables in the filter, until it finds the best form of preprocessing. This means the spectroscopic preprocessing done by the convolutional layer is more flexible and saves a lot of effort when building new calibrations. In addition, because PLSR is highly sensitive to outliers, a common technique aims to remove spectra based on distance metrics and arbitrary threshold (Wadoux et al., 2021). This normative procedure could lead to the loss of informative spectra, particularly while dealing with small datasets.

In deep learning, many techniques (e.g., robust loss function and early stopping) allow us to deal with noise. This includes original signal noise but also artificially added noise allowing for data augmentation and regularization in order to minimize overfitting and increase robustness. Deep learning allows keeping all spectra without questioning the representativeness of a highly complex and spatially explicit signal (i.e., the spectra) based on a simple global distance.

Second, wavelength range and resolution of the actual spectrometer allow for a highly multivariate signal. Reference methods to deal with this often imply dimension reduction (e.g., PLSR), leading to a loss of information. The ability of multilayer networks trained with gradient descent to learn complex high dimensional non-linear mappings makes them obvious candidates. Indeed, deep networks proved to have the theoretical guarantee that they can avoid the curse of dimensionality for many problems (Poggio et al., 2017). Among deep learning algorithms, convolutional neural networks (CNN) are known to efficiently take care of variable selection. And as already mentioned, deep learning algorithms come with multiple useful techniques to deal with the overfitting risk (e.g., batch normalization, dropout, early stopping, and noise generation).

Finally, one of the main challenges for chemometrics is to infer traits values based on the reflectance of a limited number of chemical bonds that may be shared by multiple compounds (i.e., molecules). Therefore, the prediction of the compound's content relies on a large number of very indirect relationships between reflectance values. This is already true for chemical composition (e.g., most carbohydrates exhibit only limited differences in their chemical composition) but it is even more obvious while working on functional traits resulting from multiple physiochemical changes. Moreover, the functional properties potentially arise from non-linear relationship or threshold, compensatory, moderator, and mediator effects at tissue, molecule, and chemical bounds scales. These multiscale non-linear relationships are hardly modeled by reference methods such as partial least squares regression (PLSR). The multilayer nature of deep learning algorithms allows for the identification of multiscale patterns and easily tackles non-linearity.




FIGURE 1 | Leaf reflectance as a function of light wavelength. All spectrum available in the database used to analyze the ability of spectral reflectance to predict trait values and plant categories are represented here and colored according to the experiment they come from (see **Supplementary Table S1** and **Supplementary Material** for details about experiments, conditions, as well as number of spectra per experiment). Colored lines represent the mean absorbance spectra, light grey lines represent the median absorbance spectra, dark shaded area represents spectra with absorbance ranging between the 5 and 95th percentiles, and light shaded area represents the entire absorbance range covered by the spectra.

analyses. Usually, spectral information can be exploited through the development of calibration models relating spectra and reference trait data. Calibration models are built with a representative subsample of a complete data set, in terms of the range of spectral variation treated (Foley et al., 1998). After building and validating models linking plant spectra to independently measured traits in the calibration dataset, the trait values of new samples are predicted from their spectra using these models. For that, different statistical methods are commonly used to predict trait data from spectra, including partial least squares regression (PLSR; Wold et al., 1983), principal components analysis (Dreccer et al., 2014), and 2D correlation plots (Darvishzadeh et al., 2008). However, the performance of these methods, and especially PLSR, in estimating plant traits has been shown to vary significantly across species and growth conditions (Fu et al., 2020). In recent years, machine learning approaches have become widespread in multiple fields due to their better predictive performance. Machine learning and more particularly deep learning techniques—specific machine learning

algorithms using a series of neural networks (**Box 2**)—are promising methods to improve the statistical analysis of high-throughput data (Mishra and Passos, 2021).

Spectral predictions of functional traits have been used to screen interspecific diversity across individual leaves, canopies, and biomes (Doughty et al., 2011; Roelofsen et al., 2014; Serbin et al., 2016; Wu et al., 2016; Chavana-Bryant et al., 2017). Yet, investigating intraspecific variability is crucial to connect global trait diversity to the underlying mechanisms of selection, genetic differentiation, and evolutionary adaptation (Violle et al., 2014). In this context, the model species *Arabidopsis thaliana* is an interesting model to test the predictive power of plant diversity with NIRS. Indeed, this species exhibits a large range of functional trait variation across its geographic range (Lasky et al., 2012; May et al., 2017; Price et al., 2018; Takou et al., 2018; Sartori et al., 2019), and hundreds of natural ecotypes have been fully sequenced to examine the genetic determinism of this variation (1001 Genomes Consortium, 2016). Ecological studies have taken advantage of this feature to examine the evolution of plant strategies

in response to climate (Price et al., 2018; Vasseur et al., 2018a,b; Exposito-Alonso et al., 2019; Exposito-Alonso, 2020; Lorts and Lasky, 2020). In addition, this model species has been widely used to examine metabolic and physiological features (Chan et al., 2010; Tohge et al., 2018; Wu et al., 2018). Advanced molecular techniques—“omics” approaches—allow accurate quantification of transcriptome, proteome, metabolome profiles, and fluxome (Beale et al., 2016).

On the one hand, we need to increase sample size across species, genotypes, and environments to obtain sufficient statistical power for broad generalization and predictions. On the other hand, the time-consuming careful methods required to measure physiologically meaningful (“hard”) traits limit studies to small sample sizes. We argue that a promising avenue to avoid this trade-off between generality and feasibility is to combine NIRS and deep learning computation. In this perspective article, we document how NIRS and deep learning paves the way for a quick and accurate quantification of plant trait diversity, ecological strategies, and physiological adaptation. In addition to examples from the literature, we compiled 21,032 spectra and 108 trait measurements from published and unpublished datasets (**Supplementary Table S1**) across 5,683 *Arabidopsis thaliana* plants grown in various conditions. Using this database and examples from the literature, we first show that NIRS can accurately predict leaf functional traits and identify major plant ecological strategies. Second, we show that NIRS predicts the growth conditions and the plant phenotypic response to stress. Finally, we provide evidence that NIRS can give access to new traits and functions, notably those related to plant life history, physiology, and metabolism.

NIRS QUANTIFIES FUNCTIONAL TRAIT VARIABILITY AND SUMMARIZES PLANT ECOLOGICAL STRATEGIES

A key goal of trait-based ecology is to determine the physiological mechanisms of plant adaptation to the environment through the measurement of multiple traits related to resource-use, growth, development, and phenology. Recent efforts based on analyzing interspecific trait diversity have revealed functional tradeoffs at both local and global scales (Messier et al., 2016), which suggests that plant diversity is shaped by universal constraints. For instance, Díaz et al. (2016) recently analyzed more than 45,000 plant species and demonstrated that their diversity falls along two main phenotypic dimensions: one related to plant size, which affects competitive ability and dispersal; the other related to leaf anatomy, chemical composition, and longevity. This second phenotypic dimension, called the leaf economics spectrum (Wright et al., 2004), trades off traits positively related to nutrient retention—such as leaf dry matter content (LDMC), leaf nitrogen content (LNC), and leaf life span—with traits positively related to carbon acquisition—such as specific leaf area (SLA) and leaf photosynthetic rate. Importantly, the same trade-off has been observed within species (Vasseur et al., 2012; Anderegg et al., 2018; Sartori et al., 2019).

Different theories have been proposed to categorize plant phenotypic diversity into ecological strategies related to plant

adaptation to the environment. Among these theories, Grime (1974) proposed that the quantitative variation in plant strategies is expected to result from their adaptation to contrasting levels of resource availability and disturbance. Following this hypothesis, plant strategies can thus be classified through a combination of three main axes, competitors (C), stress-tolerators (S), and ruderals (R; Grime, 1977, 1988). The “CSR” model suggests that the evolution of plant strategies is driven by trade-offs between resource capture and conservation, space occupancy, longevity, and dispersal. For instance, C-type plants invest resources into the growth of large organs to outcompete neighbors, S-type plants invest resources to conserve nutrients and protect tissues from stress damages, while R-type plants invest resources into rapid reproduction and propagule dispersal in highly disturbed environments. The CSR strategies are often depicted in a triangle with the primary types occupying the corners and intermediate forms, composed of a combination of types (e.g., “SR” and “CS”), arrayed within the triangle. The quantitative variations between CSR strategies are expected to result from plant adaptation to contrasting levels of abiotic stresses and disturbance. CSR variation has also been reported within species—notably in *A. thaliana*—and explained by evolutionary adaptation to the environment (May et al., 2017; Vasseur et al., 2018b). However, measuring through destructive methods, the numerous traits that enable the quantification of ecological strategies within—and *a fortiori* across—species remains an obstacle for the large-scale analysis of plant populations, which therefore limits our ability to temporally follow the relationships between plant traits, strategies, and the environment.

Using convolutional neural network (CNN; Box 2, **Supplementary Material, Supplementary Table S2**) to analyze our database of spectra and traits in *A. thaliana*, we show that most leaf traits were accurately predicted (**Table 1**). For instance, only leaf relative water content (RWC) and the leaf isotopic ratio of nitrogen ($\delta^{15}\text{N}$) had validation R^2 below 0.65 (**Table 1**). Yet, previous studies showed that $\delta^{15}\text{N}$ can be predicted with NIRS (Kleinebecker et al., 2009). Here, correlations between measured and predicted values were the highest for leaf traits associated with the leaf economics spectrum (SLA, LDMC, and LNC, all $r^2 > 0.85$; **Table 1**). Importantly, for these traits, the correlations calculated from the predicted data were the same as those calculated from the direct measurements ($p > 0.05$; **Figures 1A–C**). Previous studies showed that SLA can be accurately measured with NIRS from the level of individual leaves to the level of the tree canopies (Curran, 1989; Lymburner et al., 2000; Asner and Martin, 2008; Asner et al., 2009; Jacquemoud et al., 2009; Kokaly et al., 2009; Ecarnot et al., 2013; Singh et al., 2015; Serbin et al., 2016). Other LES traits have been shown to be well predicted by NIRS (Ecarnot et al., 2013; Kattenborn et al., 2017, 2019). In addition, LNC, another LES trait, can also be predicted using light reflectance at the individual leaf and at canopy levels (Sims and Gamon, 2002). Other traits related to resource-use and conservation can be predicted with spectroscopy, such as leaf age and photosynthetic capacity (Doughty et al., 2011; Chavana-Bryant et al., 2017). Thus, NIRS can provide estimates of integrated properties, such as trait covariations, whole-plant traits, and

TABLE 1 | Prediction accuracy for functional traits.

| Variable | n | Calibration | | Validation | | | |
|---|-------|-------------|----------------|------------|-------|-------|------|
| | | SD | R ² | RMSE | Bias | Slope | RPD |
| LDMC (mg g ⁻¹) | 2,932 | 52.73 | 0.86 | 16.10 | 0.38 | 1.06 | 3.28 |
| SLA (mm ² mg ⁻¹) | 3,423 | 20.90 | 0.85 | 7.47 | 0.14 | 1.01 | 2.80 |
| LNC (%) | 1,961 | 2.18 | 0.93 | 0.53 | -0.06 | 0.97 | 4.12 |
| Leaf thickness (μm) | 4,143 | 178.08 | 0.89 | 69.49 | 2.79 | 1.02 | 2.56 |
| RWC (%) | 1,421 | 22.06 | 0.17 | 4.52 | 0.40 | 1.27 | 4.88 |
| LCC (%) | 1,960 | 4.78 | 0.65 | 1.17 | 0.03 | 0.86 | 4.10 |
| δ ¹³ C | 1,222 | 1.59 | 0.83 | 0.62 | -0.04 | 0.95 | 2.56 |
| δ ¹⁵ N | 1,223 | 3.76 | 0.28 | 1.83 | -0.13 | 0.82 | 2.06 |
| Plant lifespan (days) | 1,403 | 10.55 | 0.17 | 8.01 | -1.31 | 0.86 | 1.32 |
| Plant growth rate (mg d ⁻¹) | 701 | 0.01 | 0.53 | 0.00 | 0.00 | 0.96 | 1.94 |
| C score (%) | 2,905 | 10.25 | 0.88 | 3.28 | -0.02 | 1.03 | 3.13 |
| S score (%) | 2,905 | 11.64 | 0.75 | 2.57 | 0.19 | 1.11 | 4.53 |
| R score (%) | 2,905 | 17.03 | 0.87 | 4.79 | 0.33 | 0.99 | 3.55 |

LDMC, leaf dry matter content; SLA, specific leaf area; LNC, leaf nitrogen content; RWC, relative water content; LCC, leaf carbon content; δ¹³C, fraction of ¹³C isotope; and δ¹⁵N, fraction of ¹⁵N isotope. CSR scores were estimated from leaf traits by the algorithm from Pierce et al. (2017). n is the total number of leaves used for modelling from our database that are associated with both trait and spectra measurements. All predictions have been obtained from convolutional neural network (CNN) models (see **Supplementary Material** for details). SD, standard deviation; RMSE, root mean square deviation; and RPD, relative percent difference.

strategies. Accordingly, traits such as plant growth rate and water use efficiency (estimated by δ¹³C; Farquhar et al., 1989) were also well predicted in *A. thaliana* ($r^2=0.53$ and 0.83 , respectively; **Table 1**). By contrast, predictive performance was lower for plant life span here ($r^2=0.17$), although previous studies showed that spectral profiles are able to capture key differences in plant life history (Ustin et al., 2004). Plant ecological strategies depicted by CSR scores were highly predicted in our database (**Table 1**; **Figure 1D**), as were CSR intermediate categories (e.g., SR, R/SR, S/SC, and CS), with a prediction accuracy estimated at 70% (**Table 2**).

Consistent with previous studies (Le, 2020; Barradas et al., 2021; Mishra and Passos, 2021), our results show that NIRS coupled with deep learning is a powerful tool to assess phenotypic variations in plants. Using 15 functional and metabolomic traits, we show that deep learning methods outperform classical analytical techniques such as PLSR (**Supplementary Table S3**). Moreover, deep learning approaches have numerous advantages compared to PLSR (Box 2). In particular, it does not require preprocessing of the data (cleaning and standardization of the spectra and removal of outliers), which often depends on the user's choice and differs from one dataset to another. Importantly, analyzing the spectral signature of plants with deep learning allows determining with reasonable accuracy the plant genotype. For instance here, genotype identity was correctly predicted for 64% of the tested accessions (**Table 2**), as previously observed in maize (Rincent et al., 2018). Such estimation opens promising avenues as an alternative to expensive sequencing technologies, as well as to combine genomics with phenomics.

MEASURING PLANT RESPONSES TO THE ENVIRONMENT WITH NIRS

Large-scale comparisons of ecological strategies have been performed with large databases of trait values measured on many species

under various conditions, from lab benches to greenhouse, common garden, and field conditions (Kattge et al., 2020). Although these trait databases are used to interpret plant adaptation to the environment, they surprisingly contain very little information about the response of the measured plant properties (demographics, growth rate, and traits) to the environment (Salguero-Gómez et al., 2018). Indeed, comparative approaches generally focus on interspecific variation, considering a mean trait value per species and neglecting intraspecific variability and phenotypic plasticity (but see Albert et al., 2010, 2011). For instance, CSR strategies, which should reflect environmental specialization and specific stress resistance, still remain largely unconnected to the plant evolutionary responses to biotic and abiotic stresses (Takou et al., 2018).

Spectral measurements are widely used to design screening protocols for plant drought responses (Shepherd and Walsh, 2007; Barradas et al., 2021; Burnett et al., 2021). For example, Cabrera-Bosquet et al. (2011) used spectra to accurately predict genotypic differences in the kernel and leaf water content in maize grown under different water treatments. In addition, spectral measurement is a promising method for detecting the severity of damage caused by pathogens, especially for those leaving no visible signs (Spinelli et al., 2004; Sabatier and Rutherford, 2013). Indeed, healthy plants interact (absorb, reflect, emit, transmit, and fluoresce) with electromagnetic radiation in a manner different from that of infected or damaged plants (Li et al., 2014).

To further explore the potential of NIRS as a predictive tool of plant stress level, we used experimental data included in our database (**Supplementary Material**) from 30 genotypes of *Arabidopsis thaliana* subjected to water deficit combined with either high or low (freezing) temperatures (Estarague et al., 2021). All plant individuals were measured for leaf NIRS in the course of the treatment, and survival was visually recorded after the treatment. Both measured and CNN-predicted survival rates varied from 14 to 80% depending on the genotype, with an estimated accuracy of survival prediction of 91% in an external validation dataset (**Table 2**; **Figure 2A**). Importantly, spectral measurements





TABLE 2 | Prediction accuracy for five plant categories.

| | Calibration accuracy (%) | Validation accuracy (%) |
|---------------------|--------------------------|-------------------------|
| Survival (2) | 0.988 | 0.915 |
| Genotypes (10) | 0.831 | 0.640 |
| Indoor/Outdoor (2) | 0.998 | 1.000 |
| CSR categories (11) | 0.980 | 0.700 |
| Treatment (2) | 0.955 | 0.714 |

Plant survival has two categories (dead or alive), which were measured according to the protocol described in Estarague et al. (2021). Genotypes have 10 categories corresponding to the 10 natural accessions used here. Indoor/outdoor represents whether a plant has been grown in a greenhouse or growth chamber (indoor) or in a common garden (outdoor) across all the experiments included in the database used here. CSR categories are the intermediate CSR classes estimated from leaf traits by the algorithm from Pierce et al. (2017), such as R/SR, S/SC, RS, and C/CSR (see **Supplementary Material**). Treatment has two categories (control and water stress) from the dedicated experiments included in the database (see **Supplementary Material**). All predictions have been obtained from CNN models.

were taken during the treatment, before individuals started showing visible signs of death (Estarague et al., 2021). This suggests that NIRS is a powerful tool to estimate stress effects leading to plant death early on, even before any visible sign of adverse effects.

Convolutional neural network models were able to accurately predict the environmental treatment in which plants were grown (control vs. water stress; prediction accuracy = 71%, **Table 2**). More surprisingly, CNN models reached 100% accuracy to predict if a plant was grown indoor (growth chamber and greenhouse) vs. outdoor (common garden; **Figure 2B**; **Table 2**). This result not only demonstrates the capacity of NIRS and deep learning to characterize the environmental conditions in which plants are cultivated but also suggests that plants grown indoor and outdoor have very contrasted spectral signatures. In turn, these questions our ability to draw conclusions about plant adaptation in natural conditions from experiments led in controlled conditions (growth chamber and greenhouse).



METABOLOMICS AS A NEW PHENOTYPIC DIMENSION: FUTURE PERSPECTIVES FOR THE CHARACTERIZATION OF PLANT ECOLOGICAL STRATEGIES

A broader screening of the metabolic pathways involved in the physiological adaptation of plants to contrasting environments is a promising avenue for ecology in the future. So far, large comparative approaches remain limited by the type and availability of traits collected from the literature and organized into shared databases (Kattge et al., 2020). This constraint reduces our ability to fully understand the drivers of phenotypic diversity, as well as to identify new and ecologically meaningful axes of phenotypic variation. In this perspective, NIRS allows us to detect a large variety of commonly measured chemical compounds such as phosphorus (P)—a key element of the leaf economics spectrum—and base cations [calcium (Ca), potassium (K), and magnesium (Mg)], and other micronutrients (Cozzolino et al., 2001; Ortiz-Monasterio et al., 2007; Galvez-Sola et al., 2015; Ercioglu et al., 2018; de Oliveira et al., 2019; Yu et al., 2019; Prananto et al., 2021). This opens new avenues to link resource-use strategies with plant elemental composition, fluxes, stoichiometry, and beyond, with nutrient cycling in ecosystems (Ustin et al., 2004). In addition, studies have shown that not only LNC but also chlorophyll *a* and *b* can be predicted using reflectance and transmittance of light from individual leaves and at canopy level (Sims and Gamon, 2002).

Using quantitative measurements of 67 metabolites with GC-MS and LC-MS (**Supplementary Material**), we investigated whether NIRS can estimate variations in foliar content of sugars (e.g., glucose and fructose), hormones (e.g., salicylic acid, auxin, and abscisic acid), and secondary metabolites (e.g., phenolic compounds and glucosinolates). Our results show that prediction accuracy (estimated in an external dataset; **Supplementary Material**) was highly variable between metabolites. For instance, validation r^2 ranged from 0% for the poorest predictions (see examples in **Table 3**) to 85% for the highest (dihydro caffeoyl glucuronide; **Table 3**). For sugars, the best predictions were obtained for fructose, cellobiose, mannose, and raffinose (**Table 3**). Among hormones, only auxin (IAA) and jasmonic acid (JA) were satisfactorily predicted by NIRS (**Table 3**), while other hormones were very poorly predicted (for instance, $r^2 < 0.10$). Glucosinolates are a class of metabolites produced by the Brassicaceae family, which are involved in plant defense against herbivores (Ratzka et al., 2002). Many of them showed relatively high prediction accuracy (e.g., glucoraphenin and neoglucobrassicin with $r^2 > 0.70$; **Table 3**), which paves the way for predicting plant responses to herbivore attack on many individuals at low cost. Finally, many other secondary metabolites showed substantial prediction accuracy (e.g., $r^2 > 50\%$; **Table 3**), although prediction accuracy was very variable between metabolites. More studies are needed to fully explore the potential of NIRS and deep learning to predict leaf chemistry and metabolisms. However, applying NIRS—coupled with deep learning computation—for high-throughput phenotyping of plants from cellular level to whole-plant level is perhaps the most exciting perspective of this approach.

TABLE 3 | Prediction accuracy for 67 metabolites.

| | Variable | Caibration validation | | | | | |
|-----------------------------|---|-----------------------|----------------|---------|---------|----------|-------|
| | | SD | R ² | RMSE | Bias | Slope | RPD |
| Sugars | Glucose | 6764.56 | 0.14 | 1621.88 | -4.49 | 0.95 | 4.17 |
| | Fructose | 10240.92 | 0.56 | 1316.93 | 352.08 | 1.17 | 7.78 |
| | Sucrose | 11380.72 | 0.00 | 2086.69 | 538.48 | -12.55 | 5.45 |
| | Fucose | 28.65 | 0.03 | 1.90 | 0.37 | 0.75 | 15.04 |
| | Isomaltose | 26.02 | 0.16 | 6.58 | 1.44 | 1.41 | 3.95 |
| | Cellobiose | 157.51 | 0.39 | 73.21 | 19.87 | 1.85 | 2.15 |
| | Arabinose | 37.57 | 0.00 | 51.42 | 9.39 | 100.65 | 0.73 |
| | Galactose | 293.66 | 0.18 | 304.29 | 82.21 | 1.11 | 0.97 |
| | Inositol | 911.06 | 0.31 | 136.28 | 23.17 | 1.29 | 6.69 |
| | Maltose | 58.40 | 0.02 | 57.31 | 19.37 | 0.86 | 1.02 |
| | Mannose | 219.79 | 0.42 | 35.78 | 12.77 | 2.19 | 6.14 |
| | Raffinose | 644.65 | 0.57 | 457.00 | 112.77 | 1.12 | 1.41 |
| | Rhamnose | 68.56 | 0.02 | 95.56 | 17.09 | -1150.74 | 0.72 |
| | Ribose | 32.35 | 0.00 | 42.17 | 13.41 | 138.61 | 0.77 |
| | Palatinose | 236.89 | 0.00 | 294.60 | 36.80 | -5.60 | 0.80 |
| | Melezitose | 15.62 | 0.38 | 7.47 | 1.31 | 1.26 | 2.09 |
| | Melibiose | 200.00 | 0.09 | 264.69 | 47.47 | 0.69 | 0.76 |
| Trehalose | 176.00 | 0.00 | 146.34 | 23.78 | -1.69 | 1.20 | |
| Xylose | 35.75 | 0.13 | 7.09 | 1.54 | 1.32 | 5.04 | |
| Hormones | ABA | 12.54 | 0.06 | 11.25 | 1.43 | 0.57 | 1.12 |
| | IAA | 21.37 | 0.26 | 18.16 | 1.84 | 0.95 | 1.18 |
| | JA | 337.70 | 0.29 | 197.91 | 31.53 | 1.03 | 1.71 |
| | SA | 799.00 | 0.00 | 495.41 | 147.44 | -10.54 | 1.61 |
| | CMLX | 7277.61 | 0.02 | 8086.67 | 2421.27 | 63.66 | 0.90 |
| Glucosinolates | Glucosylsin | 28.79 | 0.10 | 27.76 | 3.95 | 1.05 | 1.04 |
| | Glucobrassicin | 1462.69 | 0.15 | 914.32 | 210.01 | 0.76 | 1.60 |
| | Glucorucin | 12.22 | 0.39 | 5.88 | 0.51 | 0.86 | 2.08 |
| | Gluconapin | 5005.90 | 0.00 | 4703.53 | 2123.30 | 0.43 | 1.06 |
| | Gluconasturtiin | 94.36 | 0.00 | 91.73 | 12.46 | 0.63 | 1.03 |
| | Glucoraphanin | 1308.98 | 0.00 | 1166.48 | 250.14 | 0.22 | 1.12 |
| | Glucoraphenin | 1.78 | 0.74 | 0.62 | 0.07 | 0.91 | 2.88 |
| | Epigallocatechin | 210.86 | 0.27 | 163.05 | 2.91 | 0.83 | 1.29 |
| | Progoitrin | 666.26 | 0.01 | 564.65 | 135.83 | 0.38 | 1.18 |
| | Epiprogoitrin | 6316.22 | 0.09 | 5944.42 | 1814.64 | 0.74 | 1.06 |
| | Isobutyl | 473.57 | 0.03 | 356.50 | 56.56 | 0.67 | 1.33 |
| | Glucosinalbin | 10.35 | 0.00 | 7.96 | 1.28 | 2.52 | 1.30 |
| | Sinigrin | 4445.20 | 0.07 | 4259.39 | 1571.86 | 1.04 | 1.04 |
| | Hexyl | 49.96 | 0.00 | 45.61 | 12.28 | 0.53 | 1.10 |
| | Butyl | 5.49 | 0.51 | 3.20 | -0.24 | 1.07 | 1.72 |
| | Neoglucobrassicin Peak1 | 265.97 | 0.73 | 273.80 | 59.08 | 1.86 | 0.97 |
| | Neoglucobrassicin Peak2 | 1051.25 | 0.06 | 254.92 | 24.16 | 0.41 | 4.12 |
| | X3MTP | 47.48 | 0.51 | 9.63 | 0.36 | 1.41 | 4.93 |
| | X5MTP | 20.76 | 0.61 | 11.56 | 1.14 | 1.40 | 1.80 |
| X6MSH | 51.83 | 0.22 | 48.64 | 9.55 | 1.09 | 1.07 | |
| X7MSH | 261.68 | 0.18 | 277.93 | 88.23 | 1.19 | 0.94 | |
| X7MTH | 244.30 | 0.36 | 224.81 | 36.56 | 1.04 | 1.09 | |
| X8MSO | 2013.33 | 0.31 | 1528.42 | 169.92 | 0.87 | 1.32 | |
| X8MTO | 1278.38 | 0.17 | 1053.50 | 176.17 | 0.85 | 1.21 | |
| Other secondary metabolites | Apigenin rutinoside | 1140.31 | 0.31 | 848.50 | 73.33 | 0.63 | 1.34 |
| | Caffeic Acid | 30.01 | 0.32 | 0.96 | -0.20 | 0.74 | 31.31 |
| | Chlorogenic Acid | 29.55 | 0.66 | 16.29 | 1.38 | 1.09 | 1.81 |
| | Citrat | 2647.54 | 0.44 | 1894.98 | 169.09 | 1.08 | 1.40 |
| | Cyanidin rhamnoside | 1431.34 | 0.53 | 842.46 | -56.16 | 0.81 | 1.70 |
| | Cyanidin sophorosid glucoside | 674.85 | 0.31 | 387.08 | 88.61 | 1.04 | 1.74 |
| | Dihydro caffeoyl glucuronide | 27.05 | 0.85 | 8.96 | 0.01 | 1.12 | 3.02 |
| | Fumarat | 294.76 | 0.10 | 174.41 | 18.17 | 0.68 | 1.69 |
| | Kaempferol glucosyl rhamnosyl glucoside | 989.20 | 0.14 | 518.91 | 97.70 | 0.69 | 1.91 |

(Continued)

TABLE 3 | Continued

| Variable | Caibration validation | | | | | |
|---|-----------------------|----------------|---------|--------|-------|------|
| | SD | R ² | RMSE | Bias | Slope | RPD |
| Kaempherol rutinoside | 2788.98 | 0.59 | 1613.31 | 127.58 | 0.88 | 1.73 |
| Kaempherol xylosyl rhamnoside | 1362.13 | 0.56 | 774.66 | 7.04 | 0.88 | 1.76 |
| Malat | 1078.18 | 0.16 | 786.53 | 133.47 | 0.61 | 1.37 |
| m-Coumaric Acid | 144.26 | 0.00 | 143.67 | 18.09 | 0.84 | 1.00 |
| p-Coumaric Acid | 4.00 | 0.46 | 1.35 | -0.08 | 1.02 | 2.95 |
| Pelargonidin cumaroyl diglucoside glucoside | 69.47 | 0.65 | 34.69 | -0.28 | 0.94 | 2.00 |
| Pelargonidin sambubioside | 291.72 | 0.47 | 223.17 | 13.31 | 0.81 | 1.31 |
| Prenyl naringenin | 36.74 | 0.63 | 14.89 | -2.09 | 0.93 | 2.47 |
| Quercetin glucoside | 56.73 | 0.23 | 54.09 | 11.77 | 1.41 | 1.05 |
| Succinat | 60.74 | 0.16 | 45.15 | 0.70 | 0.93 | 1.35 |

Metabolites have been measured with GC-MS or LC-MS depending on the metabolite ($n = 124$ per metabolite) on leaves harvested on 4-week old plants grown in the greenhouse. Sugars are given in $\mu\text{mol/gFW}$; hormones in ng/gFW . For glucosinolates and other secondary metabolites, foliar relative concentrations were estimated by dividing the peak area corresponding to the metabolite by the fresh weight of the sample. SD, standard deviation; RMSE, root mean square deviation; and RPD, relative percent difference. All predictions have been obtained from CNN models (see **Supplementary Material** for details).

CONCLUSION

In this paper, we argue that NIRS coupled with recent advances in deep learning approaches is a promising method to broadly capture various information about plant functioning, ecological strategies, response to environment, and metabolism. In particular, NIRS affords considerable time and cost savings (spectrum acquisition lasts only a few seconds), and without using hazardous chemicals. In addition, samples can be analyzed in neither their natural form without destruction nor any special sample preparation. Thus, NIRS makes it possible to create extensive databases of traits at different temporal, spatial, and taxonomic scales and facilitate the adoption of phenomics into ecology. It might provide a reliable tool for the characterization of plant populations across geographical ranges, specifically if combined with other omics approaches and deep learning computation. Of course, developing calibration equations takes time, but selecting a suitable subset of samples to use in the calibration equation and validating the calibration equation take only a matter of hours in addition to standard laboratory work to chemically analyze the subset. Clearly, NIRS is more suited for larger data sets than those containing only a few samples. As calibration equations keep available for future studies, the time and financial cost of calibrations will decrease. Thus, adopting NIRS in trait-based ecology would literally multiply the number of species, genotypes, and environments potentially measurable, a key point to link functional trait variation to plant physiology and adaptation.

REFERENCES

1001 Genomes Consortium (2016). 1135 sequenced natural inbred lines reveal the global pattern of polymorphism in *Arabidopsis thaliana*. *Cell* 166, 481–491. doi: 10.1016/j.cell.2016.05.063

DATA AVAILABILITY STATEMENT

The raw data supporting the conclusions of this article will be made available by the authors, without undue reservation.

AUTHOR CONTRIBUTIONS

FV led the writing of the manuscript. All authors contributed critically to the drafts and gave final approval for publication.

FUNDING

This work was supported by INRAE, CNRS, the French Agency for Research (ANR grant ANR-17-CE02-0018-01; “AraBreed” to FV, DV, and CV), Région Occitanie (FEDER FSE IEJ 2014-2020; Project PHENOPSIS 2.0), and the European Research Council (ERC; “CONSTRAINTS”: grant ERC-StG-2014-639706-CONSTRAINTS to CV). Metabolite analytics were funded by the Deutsche Forschungsgemeinschaft (DFG, German Research foundation) – Projektnummer INST 37/696-1 FUGG.

SUPPLEMENTARY MATERIAL

The Supplementary Material for this article can be found online at: <https://www.frontiersin.org/articles/10.3389/fpls.2022.836488/full#supplementary-material>

Albert, C. H., Grassein, F., Schurr, F. M., Vieilledent, G., and Violle, C. (2011). When and how should intraspecific variability be considered in trait-based plant ecology? *Perspect. Plant Ecol. Evol. Syst.* 13, 217–225. doi: 10.1016/j.ppees.2011.04.003

Albert, C. H., Thuiller, W., Yoccoz, N. G., Soudant, A., Boucher, F., Saccone, P., et al. (2010). Intraspecific functional variability: extent, structure and

- sources of variation. *J. Ecol.* 98, 604–613. doi: 10.1111/j.1365-2745.2010.01651.x
- Anderegg, L. D. L., Berner, L. T., Badgley, G., Sethi, M. L., Law, B. E., and HilleRisLambers, J. (2018). Within-species patterns challenge our understanding of the leaf economics spectrum. *Ecol. Lett.* 21, 734–744. doi: 10.1111/ele.12945
- Andrés, S., Giráldez, F. J., López, S., Mantecón, Á. R., and Calleja, A. (2005). Nutritive evaluation of herbage from permanent meadows by near-infrared reflectance spectroscopy: 1. Prediction of chemical composition and in vitro digestibility. *J. Sci. Food Agric.* 85, 1564–1571. doi: 10.1002/jsfa.2138
- Arslan, M., Xiaobo, Z., Xuetao, H., Tahir, H. E., Shi, J., Khan, M. R., et al. (2018). Near infrared spectroscopy coupled with chemometric algorithms for predicting chemical components in black goji berries (*Lycium ruthenicum* Murr.). *J. Near Infrared Spectrosc.* 26, 275–286. doi: 10.1177/0967033518795597
- Asner, G. P., and Martin, R. E. (2008). Spectral and chemical analysis of tropical forests: scaling from leaf to canopy levels. *Remote Sens. Environ.* 112, 3958–3970. doi: 10.1016/j.rse.2008.07.003
- Asner, G. P., Martin, R. E., Ford, A. J., Metcalfe, D. J., and Liddell, M. J. (2009). Leaf chemical and spectral diversity in Australian tropical forests. *Ecol. Appl.* 19, 236–253. doi: 10.1890/08-0023.1
- Barradas, A., Correia, P. M. P., Silva, S., Mariano, P., Pires, M. C., Matos, A. R., et al. (2021). Comparing machine learning methods for classifying plant drought stress from leaf reflectance spectra in *Arabidopsis thaliana*. *NATO Adv. Sci. Inst. Ser. E Appl. Sci.* 11:6392. doi: 10.3390/app11146392
- Beale, D. J., Karpe, A. V., and Ahmed, W. (2016). “Beyond metabolomics: a review of multi-omics-based approaches” in *Microbial Metabolomics: Applications in Clinical, Environmental, and Industrial Microbiology*. eds. D. J. Beale, K. A. Kouremenos and E. A. Palombo (Cham: Springer International Publishing), 289–312.
- Biancolillo, A., and Marini, F. (2018). Chemometric methods for spectroscopy-based pharmaceutical analysis. *Front. Chem.* 6:576. doi: 10.3389/fchem.2018.00576
- Birth, G. S., and Hecht, H. G. (1987). “The physics of near-infrared reflectance” in *Near-Infrared Technology in the Agricultural and Food Industries*. ed. M. N. Saint-Paul (American Association of Cereal Chemists, Inc.), 1–15.
- Burnett, A. C., Serbin, S. P., Davidson, K. J., Ely, K. S., and Rogers, A. (2021). Detection of the metabolic response to drought stress using hyperspectral reflectance. *J. Exp. Bot.* 72, 6474–6489. doi: 10.1093/jxb/erab255
- Cabrera-Bosquet, L., Sánchez, C., Rosales, A., Palacios-Rojas, N., and Araus, J. L. (2011). Near-infrared reflectance spectroscopy (NIRS) assessment of $\delta^{18}O$ and nitrogen and ash contents for improved yield potential and drought adaptation in maize. *J. Agric. Food Chem.* 59, 467–474. doi: 10.1021/jf103395z
- Chan, E. K. F., Rowe, H. C., Hansen, B. G., and Kliebenstein, D. J. (2010). The complex genetic architecture of the metabolome. *PLoS Genet.* 6:e1001198. doi: 10.1371/journal.pgen.1001198
- Chavana-Bryant, C., Malhi, Y., Wu, J., Asner, G. P., Anastasiou, A., Enquist, B. J., et al. (2017). Leaf aging of Amazonian canopy trees as revealed by spectral and physiological measurements. *New Phytol.* 214, 1049–1063. doi: 10.1111/nph.13853
- Cozzolino, D., Fassio, A., and Gimenez, A. (2001). The use of near-infrared reflectance spectroscopy (NIRS) to predict the composition of whole maize plants. *J. Sci. Food Agric.* 81, 142–146. doi: 10.1002/1097-0010(20010101)81:1<142::AID-JSFA790>3.0.CO;2-I
- Cui, C., and Fearn, T. (2018). Modern practical convolutional neural networks for multivariate regression: applications to NIR calibration. *Chemom. Intell. Lab. Syst.* 182, 9–20. doi: 10.1016/j.chemolab.2018.07.008
- Curran, P. J. (1989). Remote sensing of foliar chemistry. *Remote Sens. Environ.* 30, 271–278. doi: 10.1016/0034-4257(89)90069-2
- Darvishzadeh, R., Skidmore, A., Schlerf, M., Atzberger, C., Corsi, F., and Cho, M. (2008). LAI and chlorophyll estimation for a heterogeneous grassland using hyperspectral measurements. *ISPRS J. Photogramm. Remote Sens.* 63, 409–426. doi: 10.1016/j.isprsjprs.2008.01.001
- de Oliveira, D. M., Fontes, L. M., and Pasquini, C. (2019). Comparing laser induced breakdown spectroscopy, near infrared spectroscopy, and their integration for simultaneous multi-elemental determination of micro- and macronutrients in vegetable samples. *Anal. Chim. Acta* 1062, 28–36. doi: 10.1016/j.aca.2019.02.043
- Díaz, S., Kattge, J., Cornelissen, J. H. C., Wright, I. J., Lavorel, S., Dray, S., et al. (2016). The global spectrum of plant form and function. *Nature* 529, 167–171. doi: 10.1038/nature16489
- Doughty, C. E., Field, C. B., and McMillan, A. M. S. (2011). Can crop albedo be increased through the modification of leaf trichomes, and could this cool regional climate? *Clim. Chang.* 104, 379–387. doi: 10.1007/s10584-010-9936-0
- Dreccer, M. F., Barnes, L. R., and Meder, R. (2014). Quantitative dynamics of stem water soluble carbohydrates in wheat can be monitored in the field using hyperspectral reflectance. *Field Crop Res.* 159, 70–80. doi: 10.1016/j.fcr.2014.01.001
- Earnot, M., Compan, F., and Roumet, P. (2013). Assessing leaf nitrogen content and leaf mass per unit area of wheat in the field throughout plant cycle with a portable spectrometer. *Field Crop Res.* 140, 44–50. doi: 10.1016/j.fcr.2012.10.013
- Ercioglu, E., Velioglu, H. M., and Boyaci, I. H. (2018). Chemometric evaluation of discrimination of aromatic plants by using NIRS. *LIBS. Food Anal. Methods* 11, 1656–1667. doi: 10.1007/s12161-018-1145-x
- Estarague, A., Vasseur, F., Sartori, K., Bastias, C., and Cornet, D. (2021). Into the range: a latitudinal gradient or a center-margins differentiation of ecological strategies in *Arabidopsis thaliana*? bioRxiv [Preprint]. doi: 10.1101/2021.10.15.461205
- Exposito-Alonso, M. (2020). Seasonal timing adaptation across the geographic range of *Arabidopsis thaliana*. *Proc. Natl. Acad. Sci. U. S. A.* 117, 9665–9667. doi: 10.1073/pnas.1921798117
- Exposito-Alonso, M., Exposito-Alonso, M., Gómez Rodríguez, R., Barragán, C., Capovilla, G., Chae, E., et al. (2019). Natural selection on the *Arabidopsis thaliana* genome in present and future climates. *Nature* 573, 126–129. doi: 10.1038/s41586-019-1520-9
- Farquhar, G. D., Ehleringer, J. R., and Hubick, K. T. (1989). Carbon isotope discrimination and photosynthesis. *Annu. Rev. Plant Biol.* 40, 503–537. doi: 10.1146/annurev.pp.40.060189.002443
- Foley, W. J., McIlwee, A., Lawler, I., Aragones, L., Woolnough, A. P., and Berding, N. (1998). Ecological applications of near infrared reflectance spectroscopy—a tool for rapid, cost-effective prediction of the composition of plant and animal tissues and aspects of animal performance. *Oecologia* 116, 293–305. doi: 10.1007/s004420050591
- Fu, P., Meacham-Hensold, K., Guan, K., Wu, J., and Bernacchi, C. (2020). Estimating photosynthetic traits from reflectance spectra: a synthesis of spectral indices, numerical inversion, and partial least square regression. *Plant Cell Environ.* 43, 1241–1258. doi: 10.1111/pce.13718
- Galvez-Sola, L., García-Sánchez, F., Pérez-Pérez, J. G., Gimeno, V., Navarro, J. M., Moral, R., et al. (2015). Rapid estimation of nutritional elements on citrus leaves by near infrared reflectance spectroscopy. *Front. Plant Sci.* 6:571. doi: 10.3389/fpls.2015.00571
- Garnier, E., Navas, M.-L., and Grigulis, K. (2016). *Plant Functional Diversity: Organism Traits, Community Structure, and Ecosystem Properties*. United Kingdom: Oxford University Press.
- Gitelson, A. A., and Merzlyak, M. N. (1997). Remote estimation of chlorophyll content in higher plant leaves. *Int. J. Remote Sens.* 18, 2691–2697. doi: 10.1080/014311697217558
- Grime, J. P. (1974). Vegetation classification by reference to strategies. *Nature* 250, 26–31. doi: 10.1038/250026a0
- Grime, J. P. (1977). Evidence for the existence of three primary strategies in plants and its relevance to ecological and evolutionary theory. *Am. Nat.* 111, 1169–1194. doi: 10.1086/283244
- Grime, J. P. (1988). Plant strategies and the dynamics and structure of plant-communities. *Nature* 336:630. doi: 10.1038/336630a0
- Jacquemoud, S., Verhoef, W., Baret, F., Bacour, C., Zarco-Tejada, P. J., Asner, G. P., et al. (2009). PROSPECT+SAIL models: A review of use for vegetation characterization. *Remote Sens. Environ.* 113, S56–S66. doi: 10.1016/j.rse.2008.01.026
- Kattenborn, T., Fassnacht, F. E., Pierce, S., Lopatin, J., Grime, J. P., and Schmidlein, S. (2017). Linking plant strategies and plant traits derived by radiative transfer modelling. *J. Veg. Sci.* 28, 717–727. doi: 10.1111/jvs.12525
- Kattenborn, T., Fassnacht, F. E., and Schmidlein, S. (2019). Differentiating plant functional types using reflectance: which traits make the difference? *Remote Sens. Ecol. Conserv.* 5, 5–19. doi: 10.1002/rse2.86
- Kattge, J., Bönsch, G., Díaz, S., and Lavorel, S. (2020). TRY plant trait database—enhanced coverage and open access. *Glob. Chang. Biol.* 26, 119–188. doi: 10.1111/gcb.14904
- Keddy, P. A. (1992). A pragmatic approach to functional ecology. *Funct. Ecol.* 6, 621–626. doi: 10.2307/2389954
- Kleinebecker, T., Schmidt, S. R., Fritz, C., Smolders, A. J. P., and Hölzel, N. (2009). Prediction of $\delta^{13}C$ and $\delta^{15}N$ in plant tissues with near-infrared

- reflectance spectroscopy. *New Phytol.* 184, 732–739. doi: 10.1111/j.1469-8137.2009.02995.x
- Knipling, E. B. (1970). Physical and physiological basis for the reflectance of visible and near-infrared radiation from vegetation. *Remote Sens. Environ.* 1, 155–159. doi: 10.1016/S0034-4257(70)80021-9
- Kokaly, R. F. (2001). Investigating a physical basis for spectroscopic estimates of leaf nitrogen concentration. *Remote Sens. Environ.* 75, 153–161. doi: 10.1016/S0034-4257(00)00163-2
- Kokaly, R. F., Asner, G. P., Ollinger, S. V., Martin, M. E., and Wessman, C. A. (2009). Characterizing canopy biochemistry from imaging spectroscopy and its application to ecosystem studies. *Remote Sens. Environ.* 113, S78–S91. doi: 10.1016/j.rse.2008.10.018
- Kothari, S., Beauchamp-Rioux, R., Laliberté, E., and Cavender-Bares, J. (2021). Reflectance spectroscopy allows rapid, accurate, and non-destructive estimates of functional traits from pressed leaves. bioRxiv [Preprint]. doi: 10.1101/2021.04.21.440856
- Lasky, J. R., Des Marais, D. L., McKay, J. K., Richards, J. H., Juenger, T. E., and Keitt, T. H. (2012). Characterizing genomic variation of *Arabidopsis thaliana*: the roles of geography and climate. *Mol. Ecol.* 21, 5512–5529. doi: 10.1111/j.1365-294X.2012.05709.x
- Le, B. T. (2020). Application of deep learning and near infrared spectroscopy in cereal analysis. *Vib. Spectrosc.* 106:103009. doi: 10.1016/j.vibspec.2019.103009
- Li, L., Zhang, Q., and Huang, D. (2014). A review of imaging techniques for plant phenotyping. *Sensors* 14, 20078–20111. doi: 10.3390/s141120078
- Lorts, C. M., and Lasky, J. R. (2020). Competition × drought interactions change phenotypic plasticity and the direction of selection on *Arabidopsis* traits. *New Phytol.* 227, 1060–1072. doi: 10.1111/nph.16593
- Lymburner, L., Beggs, P. J., and Jacobson, C. R. (2000). Estimation of canopy-average surface-specific leaf area using Landsat TM data. *Photogramm. Eng. Remote Sens.* 66, 183–192.
- May, R.-L., Warner, S., and Wingler, A. (2017). Classification of intra-specific variation in plant functional strategies reveals adaptation to climate. *Ann. Bot.* 119, 1343–1352. doi: 10.1093/aob/mcx031
- Messier, J., McGill, B. J., Enquist, B. J., and Lechowicz, M. J. (2016). Trait variation and integration across scales: is the leaf economic spectrum present at local scales? *Ecography* 40, 685–697. doi: 10.1111/ecog.02006
- Mishra, P., and Passos, D. (2021). A synergistic use of chemometrics and deep learning improved the predictive performance of near-infrared spectroscopy models for dry matter prediction in mango fruit. *Chemom. Intell. Lab. Syst.* 212:104287. doi: 10.1016/j.chemolab.2021.104287
- Ortiz-Monasterio, J. I., Palacios-Rojas, N., Meng, E., Pixley, K., Trethowan, R., and Peña, R. J. (2007). Enhancing the mineral and vitamin content of wheat and maize through plant breeding. *J. Cereal Sci.* 46, 293–307. doi: 10.1016/j.jcs.2007.06.005
- Pasquini, C. (2018). Near infrared spectroscopy: a mature analytical technique with new perspectives—A review. *Anal. Chim. Acta* 1026, 8–36. doi: 10.1016/j.aca.2018.04.004
- Petit Bon, M., Böhner, H., Kaino, S., Moe, T., and Bräthen, K. A. (2020). One leaf for all: chemical traits of single leaves measured at the leaf surface using near infrared-reflectance spectroscopy (NIRS). *Methods Ecol. Evol.* 11, 1061–1071. doi: 10.1111/2041-210x.13432
- Pierce, S., Negreiros, D., Cerabolini, B. E. L., Kattge, J., Díaz, S., Kleyer, M., et al. (2017). A global method for calculating plant CSR ecological strategies applied across biomes world-wide. *Funct. Ecol.* 31, 444–457. doi: 10.1111/1365-2435.12722
- Poggio, T., Mhaskar, H., Rosasco, L., Miranda, B., and Liao, Q. (2017). Why and when can deep-but not shallow-networks avoid the curse of dimensionality: a review. *Int. J. Autom. Comput.* 14, 503–519. doi: 10.1007/s11633-017-1054-2
- Prananto, J. A., Minasy, B., and Weaver, T. (2021). Rapid and cost-effective nutrient content analysis of cotton leaves using near-infrared spectroscopy (NIRS). *PeerJ* 9:e11042. doi: 10.7717/peerj.11042
- Price, N., Moyers, B. T., Lopez, L., Lasky, J. R., Monroe, J. G., Mullen, J. L., et al. (2018). Combining population genomics and fitness QTLs to identify the genetics of local adaptation in *Arabidopsis thaliana*. *Proc. Natl. Acad. Sci. U. S. A.* 115, 5028–5033. doi: 10.1073/pnas.1719998115
- Rascher, U., Damm, A., van der Linden, S., Okujeni, A., Pieruschka, R., Schickling, A., et al. (2010). “Sensing of photosynthetic activity of crops” in *Precision Crop Protection—the Challenge and Use of Heterogeneity*. eds.
- E.-C. Oerke, R. Gerhards, G. Menz and R. A. Sikora (Dordrecht: Springer Netherlands), 87–99.
- Ratzka, A., Vogel, H., Kliebenstein, D. J., Mitchell-Olds, T., and Kroymann, J. (2002). Disarming the mustard oil bomb. *Proc. Natl. Acad. Sci. U. S. A.* 99, 11223–11228. doi: 10.1073/pnas.172112899
- Rincent, R., Charpentier, J.-P., Faivre-Rampant, P., Paux, E., Le Gouis, J., Bastien, C., et al. (2018). Phenomic selection is a low-cost and high-throughput method based on indirect predictions: proof of concept on wheat and poplar. *G3* 8, 3961–3972. doi: 10.1534/g3.118.200760
- Roelofsen, H. D., van Bodegom, P. M., Kooistra, L., and Witte, J.-P. M. (2014). Predicting leaf traits of herbaceous species from their spectral characteristics. *Ecol. Evol.* 4, 706–719. doi: 10.1002/ece3.932
- Sabatier, D. R., and Rutherford, R. S. (2013). Can near Infrared Reflectance (NIR) Spectroscopy Be Used to Predict the Resistance of Sugarcane to Pests and Diseases in a Generic Analysis of the Plant Surface. Picking up good vibrations. Available at: <http://citeseerx.ist.psu.edu/viewdoc/download?doi=10.1.1.655.4723&rep=rep1&type=pdf#page=39> (Accessed March 22, 2022).
- Salguero-Gómez, R., Violle, C., Gimenez, O., and Childs, D. (2018). Delivering the promises of trait-based approaches to the needs of demographic approaches, and vice versa. *Funct. Ecol.* 32, 1424–1435. doi: 10.1111/1365-2435.13148
- Sartori, K., Vasseur, F., Violle, C., Baron, E., Gerard, M., Rowe, N., et al. (2019). Leaf economics and slow-fast adaptation across the geographic range of *Arabidopsis thaliana*. *Sci. Rep.* 9:10758. doi: 10.1038/s41598-019-46878-2
- Serbin, S. P., Singh, A., McNeil, B. E., Kingdon, C. C., and Townsend, P. A. (2016). Spectroscopic determination of leaf morphological and biochemical traits for northern temperate and boreal tree species. *Ecol. Appl.* 24, 1651–1669. doi: 10.1890/13-2110.1
- Shepherd, K. D., and Walsh, M. G. (2007). Infrared spectroscopy—enabling an evidence-based diagnostic surveillance approach to agricultural and environmental management in developing countries. *J. Near Infrared Spectrosc.* 15, 1–19. doi: 10.1255/jnirs.716
- Silva-Perez, V., Molero, G., Serbin, S. P., Condon, A. G., Reynolds, M. P., Furbank, R. T., et al. (2018). Hyperspectral reflectance as a tool to measure biochemical and physiological traits in wheat. *J. Exp. Bot.* 69, 483–496. doi: 10.1093/jxb/erx421
- Sims, D. A., and Gamon, J. A. (2002). Relationships between leaf pigment content and spectral reflectance across a wide range of species, leaf structures and developmental stages. *Remote Sens. Environ.* 81, 337–354. doi: 10.1016/S0034-4257(02)00010-X
- Singh, A., Serbin, S. P., McNeil, B. E., Kingdon, C. C., and Townsend, P. A. (2015). Imaging spectroscopy algorithms for mapping canopy foliar chemical and morphological traits and their uncertainties. *Ecol. Appl.* 25, 2180–2197. doi: 10.1890/14-2098.1
- Spinelli, F., Noferini, M., and Costa, G. (2004). “Near infrared spectroscopy (NIRS): perspective of fire blight detection in asymptomatic plant material” in X International Workshop on Fire Blight 704, 87–90.
- Takou, M., Wieters, B., Kopriva, S., Coupland, G., Linstädter, A., and de Meaux, J. (2018). Linking genes with ecological strategies in *Arabidopsis thaliana*. *J. Exp. Bot.* 70, 1141–1151. doi: 10.1093/jxb/ery447
- Tohge, T., Borghi, M., and Fernie, A. R. (2018). The natural variance of the *Arabidopsis* floral secondary metabolites. *Sci. Data* 5:180051. doi: 10.1038/sdata.2018.51
- Ustin, S. L., Roberts, D. A., Gamon, J. A., Asner, G. P., and Green, R. O. (2004). Using imaging spectroscopy to study ecosystem processes and properties. *Bioscience* 54, 523–534. doi: 10.1641/0006-3568(2004)054[0523:UISTSE]2.0.CO;2
- Vasseur, F., Exposito-Alonso, M., Ayala-Garay, O. J., Wang, G., Enquist, B. J., Vile, D., et al. (2018a). Adaptive diversification of growth allometry in the plant *Arabidopsis thaliana*. *Proc. Natl. Acad. Sci. U. S. A.* 115, 3416–3421. doi: 10.1073/pnas.1709141115
- Vasseur, F., Sartori, K., Baron, E., Fort, F., Kazakou, E., Segrestin, J., et al. (2018b). Climate as a driver of adaptive variations in ecological strategies in *Arabidopsis thaliana*. *Ann. Bot.* 122, 935–945. doi: 10.1093/aob/mcy165
- Vasseur, F., Violle, C., Enquist, B. J., Granier, C., and Vile, D. (2012). A common genetic basis to the origin of the leaf economics spectrum and metabolic

- scaling allometry. *Ecol. Lett.* 15, 1149–1157. doi: 10.1111/j.1461-0248.2012.01839.x
- Violle, C., Navas, M. L., Vile, D., Kazakou, E., Fortunel, C., Hummel, I., et al. (2007). Let the concept of trait be functional! *Oikos* 116, 882–892. doi: 10.1111/j.0030-1299.2007.15559.x
- Violle, C., Reich, P. B., Pacala, S. W., Enquist, B. J., and Kattge, J. (2014). The emergence and promise of functional biogeography. *Proc. Natl. Acad. Sci.* 111, 13690–13696. doi: 10.1073/pnas.1415442111
- Wadoux, A. M. J.-C., Heuvelink, G. B. M., de Bruin, S., and Brus, D. J. (2021). Spatial cross-validation is not the right way to evaluate map accuracy. *Ecol. Model.* 457:109692. doi: 10.1016/j.ecolmodel.2021.109692
- Wójcicki, K. (2015). Application of NIR spectroscopy for whisky identification and determination the content of ethanol. *Curr. Trends Comm. Sci.* 123, 123–133.
- Wold, S., Martens, H., and Wold, H. (1983). “The multivariate calibration problem in chemistry solved by the PLS method,” in *Matrix Pencils* (Berlin Heidelberg: Springer), 286–293.
- Wright, I. J., Reich, P. B., Westoby, M., Ackerly, D. D., Baruch, Z., Bongers, F., et al. (2004). The worldwide leaf economics spectrum. *Nature* 428, 821–827. doi: 10.1038/nature02403
- Wu, J., Chavana-Bryant, C., Prohaska, N., Serbin, S. P., Guan, K., Albert, L. P., et al. (2016). Convergence in relationships between leaf traits, spectra and age across diverse canopy environments and two contrasting tropical forests. *New Phytol.* 214, 1033–1048. doi: 10.1111/nph.14051
- Wu, S., Tohge, T., Cuadros-Inostroza, Á., Tong, H., Tenenboim, H., Kooke, R., et al. (2018). Mapping the Arabidopsis metabolic landscape by untargeted metabolomics at different environmental conditions. *Mol. Plant* 11, 118–134. doi: 10.1016/j.molp.2017.08.012
- Yu, E., Zhao, R., Cai, Y., Huang, J., Li, C., Li, C., et al. (2019). Determination of manganese content in cottonseed meal using near-infrared spectrometry and multivariate calibration. *J. Cotton Res.* 2:12. doi: 10.1186/s42397-019-0030-5

Conflict of Interest: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Publisher’s Note: All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

Copyright © 2022 Vasseur, Cornet, Beurier, Messier, Rouan, Bresson, Ecarnot, Stahl, Heumos, Gérard, Reijnen, Tillard, Lacombe, Emanuel, Floret, Estarague, Przybylska, Sartori, Gillespie, Baron, Kazakou, Vile and Violle. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.