



HAL
open science

Fate of polychlorobiphenyls in the insect *Tenebrio molitor*: consequences for further use as food and feed

Christelle Planche, Frédéric Mercier, Johanna Rivas, Magaly Angénieux, Hanli Wang, Benjamin Calmont, Sébastien Crépieux, Jérémy Ratel, Erwan Engel

► To cite this version:

Christelle Planche, Frédéric Mercier, Johanna Rivas, Magaly Angénieux, Hanli Wang, et al.. Fate of polychlorobiphenyls in the insect *Tenebrio molitor*: consequences for further use as food and feed. Proceedings of the Nutrition Society, 2022, 81, 10.1017/s0029665122000945 . hal-03685700

HAL Id: hal-03685700

<https://hal.inrae.fr/hal-03685700v1>

Submitted on 2 Jun 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

3rd International Symposium on Nutrition (ISN 2022): Urban food policies for sustainable nutrition and health, 27–28 January 2022

Fate of polychlorobiphenyls in the insect *Tenebrio molitor*: consequences for further use as food and feed

Christelle Planche¹, Frédéric Mercier¹, Johanna Rivas¹, Magaly Angénieux¹, Hanli Wang¹, Benjamin Calmont², Sébastien Crépieux³, Jérémy Ratel¹ and Erwan Engel¹

¹MASS group, UR QuaPA, INRAE, F-63122 Saint Genès Champanelle,
²SHNAO, Société d'histoire naturelle Alcided'Orbigny, F-63170 Aubière, and
³INVERS, F-63720 Saint-Ignat, France

Background/Objectives: Given the world population increase, there is a growing need to find alternative and sustainable protein sources for food and feed. Edible insects are an interesting solution with low greenhouse gas emissions, high feed conversion efficiency, low land use, and their ability to transform low value organic co-products into high value protein products⁽¹⁾. In counterpart, one of the potential risks associated with their consumption is the chemical contamination. In urban areas, this chemical risk may be linked to emission sources of persistent organic pollutants (POPs)⁽²⁾ such as polychlorobiphenyls (PCBs), dioxins (PCDD/Fs), polycyclic aromatic hydrocarbons (PAHs)...inducing their ubiquitous presence at trace levels in food supply⁽³⁾. In this context, this study explored the ability of yellow mealworm (*Tenebrio molitor*), the most widely bred and traded insect species in Europe⁽⁴⁾, to bioaccumulate PCBs from their feeding substrate.

Methods: *Tenebrio molitor* larvae were reared during 20 days in a temperature and humidity-controlled incubator and fed with wheat bran artificially contaminated with PCBs at a concentration of 0.67 ppb, 4 ppb or 24.4 ppb (n = 3 for each concentration). The larvae PCB content was then measured based on an extraction method inspired by Planche et al.⁽⁵⁾ and an analysis by gaz chromatography coupled with a micro-Electron Capture Detector (μ ECD).

Results: Whatever the PCB contamination level, no significant impact was observed on the larval growth. The bioaccumulation factors (BAF (%)) = $100 \times$ concentration of PCBs in larvae / concentration of PCBs in wheat bran obtained with fresh larvae are between 77% and 82%. With dried larvae, the most common commercially available form, the bioaccumulation factors ranged between 216% and 233%.

Discussion / Conclusion: This study highlights the ability of *Tenebrio molitor* to accumulate PCBs from their diet during rearing and the significant impact of the drying process on the bioaccumulation factors. This demonstrates the importance of considering the quality of the substrates used for farming insects as food and feed in terms of content in chemical contaminants including POPs like PCBs.

References

1. van Huis A (2016) *Proc Nutr Soc* **75**, 294–305.
2. Saini A, Harner T et al. (2020) *Environ Pollut* **267**, 115416.
3. Guo W, Pan B et al. (2019) *Int J Environ Res Public Health* **16**, 4361.
4. Bordiean A, Krzyżaniak M et al. (2020) *Agriculture* **10**, 233.
5. Planche C, Ratel J et al. (2015) *J Chromatogr* **A1392**, 74–81.

Disclosure of Interest

None Declared