



**HAL**  
open science

# Specificity of resistance and tolerance to cucumber vein yellowing virus in melon accessions and evidence for resistance-breaking associated with a single mutation in VPg

Cecile Desbiez, Maria Luisa Domingo Calap, Michel Pitrat, Catherine Wipf-Scheibel, Gregory Girardot, Immaculada Ferriol, Juan López-Moya, Hervé Lecoq

## ► To cite this version:

Cecile Desbiez, Maria Luisa Domingo Calap, Michel Pitrat, Catherine Wipf-Scheibel, Gregory Girardot, et al.. Specificity of resistance and tolerance to cucumber vein yellowing virus in melon accessions and evidence for resistance-breaking associated with a single mutation in VPg. 15. International symposium of plant virus epidemiology, Consejo Superior de Investigaciones Científicas, Madrid, Spain, Jun 2022, Madrid, Spain. hal-03689874

**HAL Id: hal-03689874**

**<https://hal.inrae.fr/hal-03689874>**

Submitted on 7 Jun 2022

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



## Specificity of resistance and tolerance to cucumber vein yellowing virus in melon accessions and evidence for resistance-breaking associated with a single mutation in VPg

**C. Desbiez<sup>1</sup>, M-L. Domingo Calap<sup>2</sup>, M. Pitrat<sup>3</sup>, C. Wipf-Scheibel<sup>1</sup>, G. Girardot<sup>1</sup>, I. Ferriol<sup>2,4</sup>, J. J. López-Moya<sup>2</sup>, and H. Lecoq<sup>1</sup>**

<sup>1</sup>Pathologie Végétale, INRAE, 84140 Montfavet, France; <sup>2</sup>CRAG, CSIC-IRTA-UAB-UB, Barcelona, Spain; <sup>3</sup>GAFL, INRAE, 84140 Montfavet, France; <sup>4</sup>ICA-CSIC, Madrid, 28006, Spain.

(cecile.desbiez@inrae.fr)

S6-O3

Genetic resistance is a sustainable way of controlling plant viruses, but it requires the presence of resistance genes in the host germplasm. Besides, its durability can be reduced by resistance-breaking isolates, either pre-existing in the environment or emerging after the deployment of the resistance. In the case of cucumber vein yellowing virus (CVYV, a member of the genus *Ipomovirus*), an emerging virus on cucurbits in the Mediterranean Basin (1), few resistances are available in melon. The melon accession PI 164323 was found to display complete resistance to the strain CVYV-Esp, and melon accession HSD 2458 presented a tolerance, i.e. very mild symptoms in spite of virus accumulation in inoculated plants. The resistance is controlled by one dominant allele *Cvy-1<sup>1</sup>*, while the tolerance is controlled by a recessive allele *cvy-2* independent from *Cvy-1<sup>1</sup>*. Upon inoculation with eight geographically diverse CVYV isolates corresponding to the known molecular diversity of the virus (2), the resistance was found to be strain-specific since many CVYV isolates induced necrosis on PI 164323, whereas the tolerance presented a broader range. A resistance-breaking isolate inducing severe mosaics on PI 164323 was obtained. It differed from the parental strain by a single amino-acid change in the VPg-coding region. The effect on the mutation was confirmed by reverse genetics using a newly-obtained CVYV infectious clone (3). Competition experiments suggested a fitness cost of the resistance-breaking mutation in susceptible melon. Our results highlight the need to combine cultural practices and/or additional genes to develop a more durable control of CVYV than the use of the *Cvy-1<sup>1</sup>* allele alone.

### References

- (1) Desbiez C., & Lecoq H., (2012). *Adv. Virus Res.*, 84: 67-126.
- (2) Desbiez C., et al., (2019). *Plant Pathol.*, 68: 1268-1275.
- (3) Desbiez C., et al., (2022). *Phytopathology*, *in press*.