

Table 3. Associations between infant feeding practices and ages at AP or at AR for boys and girls from the EDEN mother–child cohort study.

	Age at AP (days)		Age at AR (days)	
	Girls β [95% CI]	Boys β [95% CI]	Girls β [95% CI]	Boys β [95% CI]
Model 1				
Unadjusted				
N	n=585	n=640	n=585	n=640
Breastfeeding duration (months)	3 [2 ; 5]	1 [0 ; 2]	17 [5 ; 29]	-5 [-16 ; 6]
Age at complementary food introduction (months)	1 [-2 ; 5]	1 [-2 ; 4]	19 [-10 ; 48]	36 [11 ; 61]
Adjusted *				
N	n=585	n=640	n=585	n=640
Breastfeeding duration (months)	2 [1 ; 4]	1 [0 ; 2]	18 [6 ; 30]	-10 [-21 ; 1]
Age at complementary food introduction (months)	0 [-3 ; 4]	0 [-3 ; 3]	15 [-15 ; 45]	29 [5 ; 54]
Model using multiple imputations of missing data*				
N	n=664	n=735	n=664	n=735
Breastfeeding duration (months)	2 [1 ; 4]	1 [0 ; 2]	14 [3 ; 25]	-10 [-20 ; 0]
Age at complementary food introduction (months)	1 [-3 ; 4]	0 [-3 ; 3]	14 [-14 ; 42]	36 [14 ; 59]
Model 2				
Unadjusted				
N	n=386	n=409	n=386	n=409
Late dairy introduction and use of ready-prepared baby food (per SD)	8 [1 ; 15]	1 [-4 ; 7]	-21 [-75 ; 33]	68 [20 ; 115]
Long breastfeeding, late food introduction and home-made food (per SD)	13 [7 ; 20]	0 [-5 ; 6]	76 [25 ; 126]	18 [-30 ; 66]
Use of ready-prepared adult food (per SD)	3 [-4 ; 10]	4 [-1 ; 10]	-41 [-96 ; 14]	21 [-25 ; 67]
Adjusted *				
N	n=386	n=409	n=386	n=409
Late dairy introduction and use of ready-prepared baby foods (per SD)	4 [-2 ; 11]	0 [-6 ; 6]	-28 [-84 ; 28]	50 [1 ; 98]
Long breastfeeding, late food introduction and home-made food (per SD)	11 [5 ; 17]	0 [-5 ; 6]	76 [23 ; 129]	-12 [-61 ; 37]
Use of ready-prepared adult food (per SD)	5 [-1 ; 11]	5 [0 ; 11]	-44 [-98 ; 10]	27 [-18 ; 72]
Model using multiple imputations of missing data*				
N	n=664	n=735	n=664	n=735
Late dairy introduction and use of ready-prepared baby food (per SD)	2 [-5 ; 9]	-1 [-6 ; 4]	-13 [-67 ; 40]	37 [-1 ; 75]
Long breastfeeding, late food introduction and home-made food (per SD)	7 [2 ; 12]	1 [-3 ; 6]	50 [8 ; 92]	-5 [-48 ; 39]
Use of ready-prepared adult food (per SD)	3 [-2 ; 8]	3 [-1 ; 7]	-20 [-70 ; 29]	14 [-24 ; 51]

*Adjusted for maternal education level, smoking status during pregnancy, gestational weight gain, monthly income, maternal and paternal BMI, maternal age, sex, preterm birth, birth weight category, weight at 2 months and recruitment center. AP: adiposity peak; AR: adiposity rebound