

## First Report of Fusarium avenaceum, Fusarium oxysporum, Fusarium redolens, and Fusarium solani Causing Root Rot in Pea in France

S. Gibert, Véronique Edel-Hermann, E. Gautheron, N. Gautheron, J.-M. Sol, G. Capelle, R. Galland, A. Bardon-Debats, C. Lambert, C. Steinberg

## ▶ To cite this version:

S. Gibert, Véronique Edel-Hermann, E. Gautheron, N. Gautheron, J.-M. Sol, et al.. First Report of Fusarium avenaceum , Fusarium oxysporum , Fusarium redolens , and Fusarium solani Causing Root Rot in Pea in France. Plant Disease, 2022, 106 (4), pp.art. 1297. 10.1094/PDIS-04-21-0833-PDN . hal-03697644

## HAL Id: hal-03697644 https://hal.inrae.fr/hal-03697644

Submitted on 18 Jul 2023

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Page 1 of 3

S. Gibert

Plant disease

1 First Report of Fusarium avenaceum, Fusarium oxysporum, Fusarium redolens and Fusarium

2 solani Causing Root Rot in Pea in France

4

3

- 5 S. Gibert<sup>1,2</sup>, V. Edel-Hermann<sup>1,†</sup>, E. Gautheron<sup>1</sup>, N. Gautheron<sup>1</sup>, J. M. Sol<sup>2</sup>, G. Capelle<sup>2</sup>, R.Galland<sup>2</sup>,
- 6 A. Bardon-Debats<sup>2</sup>, C.Lambert<sup>2</sup>, and C.Steinberg<sup>1</sup>

7

- <sup>1</sup>Agroécologie, AgroSup Dijon, INRAE, Univ. Bourgogne, Univ. Bourgogne Franche-Comté, F-21000
- 9 Dijon, France
- <sup>2</sup>Société BONDUELLE SAS, Rue Nicolas Appert BP 17, 59653 Villeneuve d'Ascq CEDEX, France
- †Corresponding Author: V. Edel-Hermann; email: <u>veronique.edel-hermann@inrae.fr</u>

12

**Keywords:** Pea, France, root rot, Fusarium avenaceum, F. oxysporum, F. redolens, F. solani

14

- 15 **Funding:** Simon Gibert was financially supported by a PhD funding from Bonduelle SAS (59653
- Villeneuve d'Ascq, France) and the National Association of Technical Research (ANRT) (CIFRE
- 17  $n^{\circ}2017/0852$ ).

18

- 19 Pea (*Pisum sativum* L.) is a commonly cultivated legume in France. It can be affected by root rot diseases
- 20 caused by a pathogenic complex of soilborne fungi (Harveson et al. 2021). Symptomatic peas with black
- 21 rot lesions on their root system, sometimes associated with stunting, yellowing or wilting of the aerial parts,
- were collected in 2017 and 2018 in the Hauts-de-France, a pea-producing region in France. Fungal colonies
- 23 were obtained from diseased roots, purified by single-spore culturing, and cultivated on potato dextrose
- agar and carnation leaf agar to characterize them morphologically. Sixteen isolates, with morphological
- characteristics similar to those of *Fusarium* (Nelson et al. 1983), were identified using the translation
- elongation factor 1-α gene (*TEF1*) (O'Donnell et al. 2015). *TEF1* was amplified with primers EF1/EF2

27 (O'Donnell et al. 1998) or primers Fa/Ra (Edel-Hermann et al. 2015) and sequenced. Their GenBank acc. nos. are MW694287 through MW694302. A BLAST search showed that the sequences of strains 28 MIAE07688, MIAE07689, MIAE07690, and MIAE07691 had 100% similarity with sequences of F. 29 30 avenaceum, and 98.6 to 100% similarity with the sequence KU587615 of F. avenaceum collected from symptomatic pea in Canada. The sequences of strains MIAE07692, MIAE07693, MIAE07694, and 31 MIAE07695 had 100% similarity with sequences of F. oxysporum collected from symptomatic peas in UK 32 (acc. nos. MT630325, MT630331, MT630325, and MT630322, respectively). The sequences of strains 33 MIAE07696, MIAE07697, MIAE07698, and MIAE07699 showed 100% similarity with sequences of F. 34 35 redolens, and 99.3 to 99.5% similarity with the sequence KU587620 of F. redolens collected from symptomatic pea in Canada. The sequences of strains MIAE07700, MIAE07701, MIAE07702, and 36 MIAE07703 showed 99 to 100% similarity with sequences of F. solani, and 99.7 to 100% similarity with 37 38 the sequence KY556458 of F. pisi comb. nov. lineage (formerly F. solani f. sp. pisi) (Šišić et al. 2018). To assess pathogenicity, surface-sterilized pea seeds (cv. Firenza) were placed on malt agar and after 48 h, 39 germinated seeds were transferred into sterile glass tubes containing 20 ml of Hoagland's No. 2 basal salt 40 41 mixture at 1.6 g/liter added with 8 g/liter of agar. They were inoculated 5 days post germination with 1 ml of a conidial suspension at 10<sup>5</sup> conidia/ml or sterile deionized water for the control plantlets. Eight tubes 42 per strain and eight control tubes were prepared. The tubes were incubated in a growth chamber at 25°C 43 day / 23°C night and 16 h light / 8 h dark for 30 days. The control plantlets were asymptomatic. The four 44 F. avenaceum strains caused brown rot symptoms on the roots and wilting of aerial parts, leading to 62 to 45 46 100% of dead plantlets. The four *F. oxysporum* strains caused discolored to brown symptoms on the roots and wilting of the aerial parts, causing 100% death. The four F. redolens strains caused honey-colored to 47 brown symptoms on the roots and wilting of the aerial parts, leading to 75 to 100% of dead plantlets. Finally, 48 49 the four F. solani strains caused brown to black rot symptoms on the entire root systems up to the base of the stem, causing 100% death. Strains reisolated from infected plants had the same morphological 50 51 characteristics as the inoculated strains. To our knowledge, this is the first report of F. avenaceum, F.

Page **3** of **3**S. Gibert

Plant disease

52 oxysporum, F. redolens and F. solani pathogenic in pea in France. A large sampling and isolation campaign

is currently being performed to determine the contribution of each pathogen.

54

55

- 56 *References*:
- 57 Edel-Hermann, V., et al. 2015. J. Microbiol. Methods 111:64. https://doi.org/10.1016/j.mimet.2015.01.026
- Harveson, R. M., et al. 2021. Compendium of Pea Diseases and Pests, 3rd Edition. APS Press, St. Paul,
- 59 MN, USA. https://doi.org/10.1094/9780890546567.fm
- Nelson, P. E., et al. 1983. Fusarium species. An Illustrated Manual for Identification. The Pennsylvania
- 61 State University Press. University Park, PA, USA. ISBN: 978-0-271-00349-8
- 62 O'Donnell, K., et al. 1998. PNAS 95:2044. https://doi.org/10.1073/pnas.95.5.2044
- 63 O'Donnell, K., et al. 2015. Phytoparasitica 43:583. https://doi.org/10.1007/s12600-015-0484-z
- 64 Šišić, A., et al. 2018. Sci. Rep. 8:1. https://doi.org/10.1038/s41598-018-19779-z