



**HAL**  
open science

## Circular RNAs detection and cross-species conservation

Chloé Cerutti, Annie Robic, Thomas Faraut

► **To cite this version:**

Chloé Cerutti, Annie Robic, Thomas Faraut. Circular RNAs detection and cross-species conservation. JOBIM 2020 - Journées Ouvertes en Biologie, Informatique et Mathématiques, Jun 2020, virtual meeting, France. hal-03703612

**HAL Id: hal-03703612**

**<https://hal.inrae.fr/hal-03703612v1>**

Submitted on 24 Jun 2022

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Circular RNAs detection and cross-species conservation

Chloé CERUTTI<sup>1</sup>, Annie ROBIC<sup>1</sup> and Thomas FARAUT<sup>1</sup>  
GenPhySE, Université de Toulouse, INRAE, ENVT, 31326, Castanet Tolosan, France

Corresponding author: `Thomas.Faraut@inrae.fr`

For many years, circular RNAs (circRNAs) were considered as splicing byproducts because they were associated with low levels of expression. However, the development of high-throughput RNA sequencing and circRNAs-specific computational tools highlighted the abundance of these circRNAs in eukaryotic cells. These single-stranded RNAs are made of closed continuous loops lacking free ends and are generated during the splicing of pre-mRNA [1]. According to recent studies, circRNAs are mainly produced by exonic sequences of coding genes (exonic circRNAs) through an alternative splicing mechanism called “backsplicing”. More precisely, the end of a “donor” exon is linked to the beginning of an upstream “acceptor” exon. More rarely, circRNAs can also be produced from intronic sequences (intronic circRNAs) from intronic lariat [1]. The specific conformation of circRNAs makes their detection, quantification and functional characterization difficult [2]. Several circRNA mechanisms of action were already identified. According to existing studies, circRNAs would mainly act as microRNAs sponges or by protein interactions [2]. Recent studies have also shown that some circRNAs are evolutionarily conserved [3].

To screen the exhaustive circRNAs genomic content, we analyzed Total-RNAseq data obtained from pig (*sus scrofa*) and bovine (*bos taurus*) testicular and liver tissues. In an attempt to obtain a comprehensive overview of circRNAs in those tissues we developed an approach with a detection step agnostic to genome annotation [4]. This approach enables us to quantify the relative proportion of exonic and intronic circRNAs from coding and non-coding genes, the variability between individuals, tissues and species. Differential recruitment of splice junctions for circular transcripts versus linear transcript is also addressed.

## Acknowledgements

We thank the genotoul bioinformatics facility for providing computing resources and the institute of genome biology of FBN (Dummersdorf, Germany) for providing bovine datasets.

## References

- [1] A. Robic, T. Faraut, S. Djebali, R. Weikard, K. Feve, S. Maman, and C. Kuehn. Analysis of pig transcriptomes suggests a global regulation mechanism enabling temporary bursts of circular RNAs. *RNA Biol*, 16(9):1190–1204, 09 2019.
- [2] L. S. Kristensen, M. S. Andersen, L. V. W. Stagsted, K. K. Ebbesen, T. B. Hansen, and J. Kjems. The biogenesis, biology and characterization of circular RNAs. *Nat. Rev. Genet.*, 20(11):675–691, 11 2019.
- [3] S. Xia, J. Feng, L. Lei, J. Hu, L. Xia, J. Wang, Y. Xiang, L. Liu, S. Zhong, L. Han, and C. He. Comprehensive characterization of tissue-specific circular RNAs in the human and mouse genomes. *Brief. Bioinformatics*, 18(6):984–992, Nov 2017.
- [4] Y. Gao and F. Zhao. Computational Strategies for Exploring Circular RNAs. *Trends Genet.*, 34(5):389–400, 05 2018.