

1 Fast improvement of macrofauna communities and soil quality in plantain crops 2 converted to agroecological practices

3
4 Gladys Loranger-Merciris^{a,b,*}, Harry Ozier-Lafontaine^b, Jean-Louis Diman^c, Jorge Sierra^b, Patrick
5 Lavelle^d
6

7 ^a Université des Antilles, UMR ISYEB-MNHN-CNRS-Sorbonne Université-EPHE, UFR Sciences Exactes
8 et Naturelles, Campus de Fouillole, Pointe-à-Pitre Cedex, 97157 Guadeloupe, France

9 ^b ASTRO Agrosystèmes tropicaux, INRAE, 97170 Petit-Bourg, Guadeloupe, France

10 ^c UE PEYI, INRAE, 97170 Petit-Bourg, Guadeloupe, France

11 ^d Institut de Recherche pour le Développement/ Université Paris Sorbonne IEES, 32 rue Henri
12 Varagnat, 93143 BONDY Cedex, France

13 * Corresponding author. E-mail address: Gladys.Loranger@univ-antilles.fr (G. Loranger-Merciris).
14
15

16 ABSTRACT

17 Plantain (*Musa paradisiaca* AAB) is one of the main staple food crops in tropical areas, particularly in
18 the Caribbean region. Therefore, supporting a sustainable production of plantain in this zone is
19 crucial to secure food and to provide income to millions of people. We hypothesized that
20 agroecological practices based on multi- functionality and biodiversity and that preserve soil should
21 i) increase macrofauna abundance, ii) enhance soil chemical fertility, iii) improve soil physical
22 structure, iv) reduce plant infestation by parasitic nematodes, and v) immediately achieve similar
23 yield as in conventional systems. For this purpose, we compared plantain conventional and
24 agroecological systems at the farm scale. The agroecological system consisted in a combination of
25 three innovations: i) inputs of vermicompost to replace mineral fertilizers, ii) use of healthy planting
26 material and iii) use of a mulching of living crop to control weeds. Ten months after the plantation,
27 soil macrofauna abundance had doubled and taxonomic richness had increased by 45% in the
28 agroecological system. In addition, soil structure was improved in the agroecological system (52% of
29 biogenic aggregates in the agroecological system vs. 21 % in conventional systems), as well as the
30 abundance of free-living nematodes associated to plantain roots increased by 73 % in the
31 agroecological system. The abundance of *Radopholus similis*, one of main plant parasitic nematodes,
32 decreased in the agroecological system. Moreover, the agroecological system achieved similar crop
33 yield level than conventional systems. Agroecological practices are thus an efficient option to allow a
34 fast transition from conventional to environmentally friendly cropping systems.
35

36 Keywords: Legume cover crop, Plants Issued from Fragments (PIF), Soil aggregates, Soil fauna,
37 Tropical agroecosystems, Vermicompost
38
39

40 1. Introduction

41
42 One of the most relevant challenges of the 21st century is to accelerate the transition of
43 conventional intensive agricultural systems to agroecological, while maintaining high levels of yields,
44 quality of products and enhance the provision of soil-based ecosystem services (Dendoncker et al.,
45 2018). However, the green revolution paradigms supporting crop production are still in use and, at a
46 global scale, generate major environmental concerns (Tilman et al., 2002; Den doncker et al., 2018).

47 The intensive agricultural practices are at the origin of massive pollution of ecosystems and soil
48 degradation. These intensive agricultural practices also greatly decrease soil biological diversity and
49 activity (Clermont-Dauphin et al., 2014; Tsiafouli et al., 2015; Lavelle et al., 2022) and ecosystem
50 regulation services (e.g., decrease in water quality, increase of greenhouse gases in the atmosphere,
51 pollution) (Millennium Ecosystem Assessment (MEA), 2005; Lavelle et al., 2014). Development of
52 new, more environmentally friendly cropping systems, preventing the depletion of natural resources,
53 is currently a priority for stakeholders, legislators and consumers, according to the growing interest
54 in developing alternative sustainable farming strategies driven in the 1990s (Nkonya et al., 2016). All
55 of these strategies share the same objective of minimizing or suppressing the use of synthetic inputs,
56 enhancing organic matter inputs, and improving agroecosystems health, while maintaining high
57 production level (Altieri, 1999; Andres and Bhullar, 2016; Meynard, 2017). These agroecological
58 strategies promote the development of practices based on the use of organic resources produced on
59 farm and the enhancement of natural ecological processes. According to Pretty (2008), sustainable
60 agriculture jointly produces food and goods for farmers and the environment. In these agroecological
61 systems, pests and diseases are supposed to be controlled by natural predators and parasites, and
62 favourable soil structure to be maintained by the roots and macrofauna activities (Kibblewhite et al.,
63 2008). However, these agroecological systems still need to be evaluated in terms of agronomic and
64 environmental impact and economic feasibility (El Mujtar et al., 2019). A critical issue is the decrease
65 in production generally observed in the early stages of transition (Schrama et al., 2018). The
66 restoration of soils degraded by aggressive practices may last several years (Mosier et al., 2021). This
67 delay may be a severe obstacle to the adoption of environmentally friendly practices in the absence
68 of supporting public policies. Practices able to accelerate this transition are needed (Peters, 1991;
69 Zinati, 2002).

70 Plantain (*Musa paradisiaca* AAB), a staple food grown throughout the tropics, constitutes a
71 major source of carbohydrates for millions of people in Africa, the Caribbean, Latin America, Asia and
72 the Pacific (e.g., Robinson, 1996; Tchango et al., 1999; Lescot, 2004; Ekunwe and Ajayi, 2010). In
73 2020, plantain was grown in 54 countries and more than 43 million tons were produced worldwide
74 (FAOSTAT, 2021). This crop provided food security and income for small growers who represent the
75 majority of producers in the tropics (Kahane et al., 2013). Latin America and the Caribbean produced
76 18 % of the global production of plantain (FAOSTAT, 2021). Therefore, sustainable production of
77 plantain in this region is crucial to provide food security and income to millions of people (Picq et al.,
78 1998). Pests are a major constraint for plantain production (Godefroid et al., 2017). Thus, there is a
79 potential for yield gains through pest control improving. Plant-parasitic nematodes are major
80 underground pests of plantain that impair the transport of nutrients and water to the main stem,
81 causing plant toppling and yield reduction (Gowen et al., 2005). Nematodes are mainly controlled by
82 synthetic pesticides that have negative effects on other soil living organisms (Lavelle et al., 2004;
83 Carrascosa et al., 2014). Weeds can also affect plantain production by competing for water and
84 nutrient resources, and by hosting parasites and diseases that affect plantain (Obiefuna, 1989). These
85 weeds are mainly chemically controlled. Herbicide treatments are usually carried out at the
86 beginning of plantation, and can be repeated every month. This chemical control has environmental
87 and economic costs. The availability and quality of plantain planting material is also one of the
88 limitations for the development of this crop. Farmers often used suckers already infested with pests
89 and diseases as their only source of planting material. Due to the high disease and pest pressure, the
90 lifetime of the majority of French West Indies plantain plantations is very short, i.e., 2–3 years for
91 plantains vs. 7 years or more for intensively managed dessert bananas (Qu'én'éherv'é, 2009).
92 However, some plantain plantations easily last about twenty years in South America or in Africa

93 (Gold, 1991). It is particularly important to develop methods which increase crop longevity and
94 reduce soil biological degradation due to the excessive use of chemical inputs.

95 We tested the hypotheses that agroecological practices based on multi-functionality and
96 biodiversity and that preserve soil should be able to i) increase macrofauna abundance and benefit
97 from their favourable activities as ecosystem engineers (Lavelle et al., 2016), ii) enhance soil fertility
98 associated to organic fertilization, iii) improve soil physical structure as a result of enhanced biogenic
99 aggregation, iv) reduce plant infestation by plant-parasitic nematodes associated to direct control by
100 earthworms (Blouin et al., 2005; Lafont et al., 2007), and v) achieve similar yield as in conventional
101 systems. In order to test these hypotheses, the conventional intensive practices in three farms and
102 an innovative cropping system were compared on the basis of agronomic and ecological criteria.
103 Reduction of agrochemical inputs is the intended goal, especially in French West Indies where
104 ecosystems have attained high levels of contamination (Ozier-Lafontaine and Lesueur-Jannoyer,
105 2014). To achieve this goal, we used: i) healthy plants issued from the Plants Issued from Fragments
106 (PIF) method (Lefranc et al., 2010; Sodom et al., 2010), a method of vegetative propagation instead
107 of suckers which are most of the time already infected by pests, ii) nutrient supply with organic
108 fertilization with vermicompost instead of mineral fertilizers, and iii) use of a cover crop instead of
109 herbicides to control weeds.

110

111 **2. Material and methods**

112

113 2.1. Site information and experimental design

114

115 The study was carried out in the Western part of Guadeloupe where most plantain
116 plantations are found. In order to carry out our experimentation, three farmers' plots were chosen.
117 These farms were located from 0 to 150 m altitude, with annual rainfall ranging from 2000 and 3000
118 mm year⁻¹ and mean annual temperature around 23°C (France Meteorological Service,
119 <http://www.meteo.gp>). Each plot had uniform management, soil type and topography. Soils are
120 Haplic Nitisols dominated by halloysite clay minerals (FAO-UNESCO classification, Driessen et al.,
121 2001; Clermont-Dauphin et al., 2004). The first two plots farms (Farm 1 and Farm 2) were located at
122 "La Sarde" in the city of Capesterre-Belle-Eau (respectively, 16° 6' 88" N, 61° 34' 12" W, and 16° 6'
123 15" N, 61° 34' 69" W). The third one was located at "Valombreuse", in the city of Petit-Bourg (16° 10'
124 44" N, 61° 38' 97" W). The experimentations were set up during the dry season between February
125 and March 2013. Before the plantation, the three plots have been left in fallow since at least two
126 years (Table 1).

127 Two cropping systems were set up in each of the three farms. The first one was the
128 conventional cropping system of each farmer which is described in Table 1. Planting material
129 (suckers) was collected from an old unproductive field, where suckers were separated from their
130 mother plant with a machete. The suckers' corms have been carefully peeled with a machete and the
131 pseudostems were cut off a few centimeters above the corm. Peeling of the corm was supposed to
132 delay the development of nematode infestation, while cutting the pseudostem reduces bulkiness and
133 improves early growth of the newly planted sucker.

134 The second cropping system was agroecological and integrated 3 innovations (Table 1): i)
135 healthy plants, obtained through the PIF technique (Lefranc et al., 2010; Sodom et al., 2010), were
136 used in order to reduce pest infestations at the beginning of the plantation; ii) vermicompost
137 (mixture of cattle manure and green wastes transformed by local epigeic earthworms) was applied in
138 order to fertilize plants and stimulate soil food web; and iii) a living vegetal covering using *Arachis*

139 pintoï Krapov. and W.C. Greg (Fabaceae) was used to control weeds through mulching. Ultimately,
140 we added 18.75 t ha⁻¹ of fresh vermicompost (corresponding to 11 t ha⁻¹ of dry matter) during the
141 experiment, in the planting hole mixed with soil at the beginning, then at the soil surface, 2 and 6
142 months after the start of the experiment (Table 1). Characteristics of the vermicompost were: pH 7.5,
143 Total C 215 g.kg⁻¹, Total N 20.3 g kg⁻¹, C/N 12.8, Total P 5.4 g kg⁻¹, Total K 13.3 g kg⁻¹, Total Mg 6.2 g kg⁻¹,
144 Total Ca 86.3 g kg⁻¹, NO₃-N 2449 mg kg⁻¹, NH₄-N 9.7 mg kg⁻¹. *Arachis pintoï* forms a very dense mat
145 that remarkably controls weeds. Seven weeks after its implantation, the plant already covered partly
146 the soil and reached 100 % of coverage in six months. *Arachis pintoï* was allowed to senesce naturally
147 within the plantain cropping field.

148 In each farm, an experimental plot involved 150 plantain plants divided between the two
149 treatments with 3 repetitions by treatment (Fig. 1A). Each elementary experimental unit contained 9
150 central plantains and 16 border plantains (Fig. 1B). Spacing between the plantain rows was 2.5 m and
151 2 m within the row (i.e., 2000 plants per ha). Before plantation, the soil was ploughed in order to
152 burry weeds. Vegetal material (suckers or PIF) was planted immediately after field preparation. At
153 the flowering stage, 10 months after the plantation, soil chemical, morphological and biological
154 characteristics were measured. Plant health (root necrosis index and plant parasitic nematodes com-
155 munities) and yield were also measured.

156

157 2.2. Soil chemical characteristics

158

159 Soil chemical properties were measured in each of the 18 experimental units (6 experimental
160 units × 3 farms). In each experimental unit, soil samples were taken with an 8 cm diameter × 15 cm
161 height cylinder, near the 9 central plants (Fig. 1). These 9 samples were mixed in a composite sample
162 in which chemical analyses were done. Total soil C and N were determined using an auto-analyser
163 (NF ISO 13878, 1998). Available P was measured using the Olsen-Dabin method (Dabin, 1967). Soil
164 mineral N (NO₃ and NH₄) was measured by colorimetry after extraction with a 0.5 M KCl solution.
165 Exchangeable Ca, Mg, K and Na were determined after extraction with ammonium acetate (NF X
166 31108). Cation-exchange capacity (CEC) was measured with IF07–10D (NFX 31–130) method and pH-
167 H₂O and pH-KCl with NF ISO 1770, 3696 and 11464.

168

169 2.3. Morphology of soil aggregates

170

171 The morphological assessment of soil macroaggregates provides an integrative index of soil
172 biological activity. An undisturbed soil sample using an 8 cm diameter × 8.5 cm height cylinder was
173 taken at 50 cm from the central plantain (plant n° 5, Fig. 1B) in each of the 18 experimental units. In
174 the laboratory, blocks were gently separated and the samples were air-dried then sieved at 2 mm.
175 The soil retained by the sieve was placed on a filter paper and the various elements were separated
176 according to their origin and shape. The elements were sorted according to categories: i) biogenic
177 aggregates of rounded forms, created by macroinvertebrates, ii) physical aggregates of angular
178 forms, produced by the physical processes of the environment (especially alternating dry and wet
179 periods), iii) non-macroaggregated soil (soil passing through the 2 mm sieve), and iv) other
180 components (roots, stones, litter, wood). The separated samples were put in an oven at 60°C for 15
181 days and weighed (Velasquez et al., 2007).

182

183 2.4. Soil biological characteristics

184

185 Ten months after the plantation, at the end of the wet season, soil macro-invertebrates were
186 hand sorted from soil monoliths of 25 cm x 25 cm x 20 cm, taken at 50 cm from the central plantain
187 in each of the 18 experimental units, following the ISO 23611-S methodology (ISO, 2011). Soil macro-
188 invertebrates were counted and classified into eighteen taxonomic groups: Diplopoda, Formicidae
189 (ants), Isoptera (termites), Lumbricina (earthworms), Coleoptera, Coleoptera larvae, Chilopoda,
190 Isopoda, Dermaptera, Blattaria, Araneae, Gastropoda, terrestrial Turbellaria, Hemiptera, Orthoptera,
191 Diptera larvae, Lepidoptera larvae and Thysanoptera (Table 2). Taxonomic richness was calculated
192 based on these groups. Earthworms were identified at the morphospecies level.

193

194 2.5. Plant health and yield

195

196 Plant parasitic and root-associated free-living nematodes were extracted from primary roots
197 removed from a 25 cm x 25 cm x 30 cm soil block taken at 50 cm from the central plantain in each of
198 the 18 experimental units. Nematodes were extracted in a mist chamber (Viglierchio and Schmitt,
199 1983). Individuals were counted in aliquots and expressed as number of nematodes per 100 g of root
200 fresh biomass. The main plant parasitic nematodes species were identified under a light microscope.

201 To assess the severity of root damage, a 100 cm³ volume of soil was taken at the base of the
202 central plant for each of the 18 experimental units. Roots were removed, carefully washed and cut
203 longitudinally. Necrosis rates on the internal surfaces of these roots were evaluated. The color of
204 root necrosis due to soil-borne pathogens varied from reddish to black while healthy roots were
205 white. Notations were rated with a root necrosis index (RNI) based on different classes: 0, no lesion;
206 1, 1–25% of root area is necrotic; 2, 26–50%; 3, 51–75%; and 4, more than 75% (Bridge and Gowen,
207 1993). Average sums of necrosis notations obtained for each plant were calculated for each
208 treatment. In each experimental unit, the bunches of the 9 central plantains (Fig. 1B) were collected
209 and weighted at the end of the experiment (11–12 months after plantation). Plant yield was
210 estimated by bunch weight.

211

212 2.6. Statistical analysis

213

214 Generalized linear mixed models (GLMM, Bolker et al., 2009) were used to analyse the
215 relationships between cultural practices and soil physico-chemical and biological features. In GLMM,
216 the linear predictor contains random effects in addition to fixed effects. The inclusion of random
217 effects allows reporting the effect of variables that create variance and contain unobserved
218 heterogeneity but that are not important to test (Duyck et al., 2012). In our study, we treated “Farm”
219 as a random effect and “Cropping system” as a fixed effect. These analyses were made using XLSTAT®
220 software. Analysis of Similarity (ANOSIM) was performed in order to compare earthworm
221 communities in both systems, using XLSTAT® software (XLSTAT-R, vegan package).

222 In addition, we performed principal component analyses (PCA) on biological and chemical
223 soil parameters, respectively. Biological parameters comprised main groups of soil macrofauna,
224 morphological characteristics, root necrosis index, nematode communities and plant yield. Soil
225 parameters comprised a set of 12 standard variables describing soil chemistry (Table 3). Monte Carlo
226 tests on coordinates of sampling points on factorial axes allowed to test for significant differences
227 among farms and experimental situations. These analyses were done with the Ade4 library in the R
228 environment (Thioulouse et al., 2018).

229

230 3. Results

231
 232
 233
 234
 235
 236
 237
 238
 239
 240
 241
 242
 243
 244
 245
 246
 247
 248
 249
 250
 251
 252
 253
 254
 255
 256
 257
 258
 259
 260
 261
 262
 263
 264
 265
 266
 267
 268
 269
 270
 271
 272
 273
 274
 275
 276

3.1. Soil chemical characteristics

Soils were acidic to slightly alkaline, with average values of pH-H₂O between 4.5 and 7.0. Soil pH, N-NH₄, N-NO₃, total N, total C, CEC, K, Na, Ca, Mg and P content did not significantly differ among the two cropping systems (Table 3).

3.2. Soil morphological characteristics

Non-macro aggregated soil represented on average 50.0 % of the total soil mass in conventional systems and 28.0 % in agroecological systems. In contrast, biogenic aggregates represented on average 20.7 % of the total soil mass in conventional systems and 52.1 % of the total soil mass in agroecological systems. Agroecological systems had significantly higher proportions of biogenic aggregates ($F_{1,14} = 127.80$, $p < 0.0001$), and lower proportions of non-macro aggregated soil ($F_{1,14} = 36.99$, $p < 0.0001$) than conventional systems. Proportion of physical aggregates was not significantly different in both systems (respectively 27.9% in conventional systems and 19.0% in agroecological systems, $F_{1,14} = 3.24$, $p = 0.09$), Fig. 2.

3.3. Soil biological characteristics

Earthworms were the most abundant group of macroinvertebrates (40.7%) followed by “Diplopoda” (24.6%), and “Ants” (9.7%). We found 206.8 ± 34.0 (mean \pm SE) individual m⁻² in conventional systems compared to 416.6 ± 72.0 individual m⁻² in agroecological systems. The density of total soil macrofauna was significantly higher in agroecological systems than in conventional systems ($F_{1,14} = 7.45$, $p = 0.016$). The abundance of “Earthworms” was significantly higher in agroecological systems compared to conventional systems ($F_{1,14} = 7.91$, $p = 0.014$). However, there was no significant difference between agricultural systems for other macroinvertebrates taxonomic groups (Fig. 3, Table 2). Yet, the taxonomical richness (number of taxonomic groups) was significantly higher in agroecological systems than in conventional systems ($F_{1,14} = 7.61$, $p = 0.015$); 14 taxonomic groups were found in conventional systems compared to 17 in agroecological systems (Table 2). Two species of earthworms were found in the conventional system: *Pontoscolex corethrurus* (Müller) and an undetermined species. Four species were found in the agroecological system: *P. corethrurus*, *P. spiralis* (Borges and Moreno) and two undetermined species. The endogeic earthworm species *P. corethrurus* prevailed in the earthworm community in both treatments (91 % of earthworm community in conventional systems and 81 % in agroecological system). The community of earthworms was significantly different in both treatments (ANOSIM, $R = 0.218$, $p = 0.007$).

3.4. Plant health and yield

Population of *Radopholus similis* (Cobb) was significantly higher in conventional systems compared to agroecological systems ($F_{1,14} = 4.98$, $p = 0.042$). Moreover, the abundance of free-living nematodes associated to plantain roots was significantly higher in agroecological systems than in conventional systems ($F_{1,14} = 17.91$, $p = 0.001$). We found 2399 ± 349 ind 0.100 g roots⁻¹ in conventional systems and 4150 ± 275 ind 0.100 g roots⁻¹ in the agroecological system (Table 4). The root necrosis index (65 ± 9 % in conventional systems and 57 ± 11 % in agroecological systems) was not significantly different among systems ($F_{1,14} = 0.79$, $p = 0.389$).

277 Bunch weight and estimated plantain yield were not significantly different among cropping
278 systems. The average bunch weight was 24 ± 3 Kg by plant (corresponding to 48 ± 6 t ha⁻¹) in
279 agroecological systems and 28 ± 2 Kg by plant (corresponding to 57 ± 5 t ha⁻¹) in conventional
280 systems ($F_{1,14} = 1.67$, $p = 0.218$).

281

282 3.5. Agroecological systems versus conventional systems

283

284 Principal Component Analyses of the data matrix confirmed the trend expressed above. PCA
285 of biological parameters (Fig. 4) significantly separated conventional from agroecological plots (18 %
286 variance explained, $p < 0.01$). Axis 1 opposed sites with high abundance and diversity of
287 macroinvertebrates and free-living nematodes with predominant biogenic aggregates or non-
288 macroaggregated soil, on the left end side, to sites with dominance of physical macroaggregates,
289 high plant yield, and plant parasitic nematodes of genera *Meloidogyne*, *Heli-* *cotylenchus* and
290 *Radopholus*, on the right end. Axis 2 opposed sites with high proportions of biogenic aggregates and
291 high general abundance of Myriapoda and macroinvertebrates in general to sites with dominant non-
292 macroaggregated soil and high root necrosis index. Points with agroecological farming practices were
293 associated with highest macro- invertebrate abundance, biogenic macroaggregation and high
294 densities of free-living nematodes while points with conventional systems, with lower abundance of
295 macroinvertebrates, projected closer to plant parasitic nematodes and physical macroaggregation.

296 Farms were also significantly separated (26 % explained variance, $p < 0.01$) independent of
297 the systems, with Farm 1 on the left end side with higher abundance of *Pratylenchus* sp., non-
298 macroaggregated soil and root necrosis and the other two farms on the opposite side, with higher
299 yield and physical macroaggregation. PCA of chemical parameters did not significantly separate
300 conventional from agroecological plots.

301

302

303 4. Discussion

304

305 The major finding of this study was that changes in practices in plantain agrosystems
306 significantly affected macrofauna taxonomical richness, soil morphological characteristics and
307 nematodes communities, in only 10 months. This shows a fast transition from conventional to
308 agroecological cropping systems, with no significant loss of production. As expected, the
309 agroecological plantain system was characterized by a higher soil macrofauna richness, a better soil
310 structure (higher proportions of biogenic aggregates), a decrease in plant parasitic nematodes and an
311 increase of the abundance of free-living nematodes associated to plantain roots. However, the
312 agroecological plantain system did not affect significantly soil chemical parameters. Similar
313 experiments in several places have shown that, while soil macro-invertebrates communities respond
314 immediately to any changes in soil conditions, physical parameters change with some time lag and
315 chemical only in the long term (Velasquez and Lavelle, 2019).

316

317 4.1. Agroecological practices and soil macrofauna

318

319 The significant increase in macrofauna abundance (due to increase of earthworm abundance)
320 and taxonomic richness within the agroecological system may be firstly due to non-application of
321 herbicides, pesticides and inorganic fertilizers in this agrosystem. In fact, herbicides used in
322 conventional systems, especially Basta® (active ingredient glufosinate) and Reglone® (active

ingredient diquat) are known to negatively impact terrestrial soil invertebrates (Griffiths et al., 2008; Druart et al., 2010; Niedobova et al., 2019). Moreover, Sánchez-Moreno et al. (2009) showed that the fosthiazate pesticide (active ingredient of the nematicide Nemathorin®), usually used in conventional plantain systems, had a high toxicity to non-target fauna (including arthropods), and a high persistence in the environment. In our study, the earthworm community is very low in conventional systems. This result is supported by the study of Pelosi et al. (2014) which showed that earthworms are highly impacted by pesticides at all organization levels. The positive impact of the agroecological plantain system on soil macrofauna may be also due to vermicompost input. Few studies deal with the impact of vermicompost on earthworm populations. In the same way, Chaudhuri et al. (2016) also showed that vermicompost can increase the abundance and biomass of earthworms. It is known that inputs of organic matter such as plant-derived material usually increase the biological activity and number of trophic groups (Brussaard, 1998). At last, the presence of the legume cover crop *A. pintoï* may also impact soil macrofauna in the studied agroecological system.

As in our study, it has been showed in previous studies that *A. pintoï* significantly enhance earthworm density and fauna diversity in Amazonian pastures (Laossi et al., 2008; Velasquez et al., 2012). Earthworms may respond to litter quality or biomass, *A. pintoï* producing high amounts of organic matter (Laossi et al., 2008). *A. pintoï* produced a very dense leaf cover just above the ground and thus might also provide a better habitat for soil organisms with its complete soil cover (Perin et al., 2003).

342

343 4.2. Agroecological practices and soil structure

344

Soil physical characteristics (linked to soil aggregation) were improved in the agroecological plantain system with a great increase in biogenic aggregates produced by macro-invertebrate ecosystem engineers. Besides, the abundance of earthworms was significantly higher in this cropping system. Several studies have already reported such effects (Velasquez et al., 2007, 2012). The endogeic earthworm species, *P. corethrus*, known to produce large amounts of biogenic aggregates, prevailed in the earthworm community in the present study. Their casts, which were stabilized by ageing, gave the soil high structural stability (Blanchart et al., 1999). This increase of biogenic aggregates may be enhanced by the legumes groundcover which has been showed to control soil physical characteristics via their impact on soil engineers. As an example, biogenic macroaggregates increased by 87 % in plots with the herbaceous legume *A. pintoï* compared to plots without *A. pintoï* (Velasquez et al., 2012). Moreover, in their study, Velasquez et al. (2012) showed that there was a parallel increase in earthworm abundance and decrease in soil C:N ratio under *A. pintoï*. This result suggests that earthworms benefited from improved nutrition in the presence of *A. pintoï*, thus enhancing their activity and their influence on soil structure. Soil structure enhancement associated with earthworms and growing plants could have important benefits for water capture, erosion control, gas exchange, root penetration and C stabilization in soils (Velasquez et al., 2012).

361

362 4.3. Agroecological practices and regulation of pests

363

Our study also revealed that the populations of *R. similis* were lower and of root-associated free-living nematodes were significantly greater in the agroecological system compared with the conventional system. Non-use of chemical nematicides and insecticides and organic fertilization may explain this result. Other studies also showed that organic amendments increased the diversity and functional composition of soil nematode communities (Pattison et al., 2006, 2021). In particular, the

368

369 supply of organic matter boosted the populations of microbivorous nematodes, particularly
370 bacterivores (Ferris and Bongers, 2006) and of carnivorous nematode populations (Tabarant et al.,
371 2011) and not of plant parasitic nematodes. Thus, the input of organic matter would be an
372 interesting alternative method to reduce plant parasitic nematodes, maintain plant health and
373 lengthen the duration of plantain plots. Additionally, in the agroecological system, the reduction of *R.*
374 *similis* population may be linked to earthworm abundance since they are known to control their
375 impact in various ways (Blouin et al., 2005; Lafont et al., 2007; Wurst, 2010; Loranger-Merciris et al.,
376 2012; Rowen et al., 2019).

377

378 4.4. Agroecological practices, soil fertility and plantain yield

379

380 Our study did not show any differences between the two plantain cropping systems
381 regarding soil fertility (chemical properties). It is important to note that, in our study, the soil
382 chemical properties were determined 10 months after the plantation. It has been showed that *A.*
383 *pintoï* mulching significantly improved soil fertility in long-term experiments in Australia (Johns,
384 1994) and China (Zhong et al., 2018). Vermicomposting converted organic waste into an organic
385 product with high content of nutrients, microbial matter, and stabilized humic substances (Sierra et
386 al., 2013). In a recent study, Sihi et al. (2017) found that long-term application of organic residues in
387 cultivations of basmati rice favored biologically available nutrients in organic systems. Moreover, a
388 recent meta-analysis by Chen et al. (2018) showed that the continuous application of organic
389 amendments caused greater gains in soil organic carbon, significantly increase of Olsen P over time
390 and increased the resilience of agronomic systems. Organic substrates stimulated soil
391 microorganisms that produce enzymes responsible for the conversion of unavailable nutrients to
392 plant available forms and should benefit to chemical soil fertility in the long term or if massive
393 amendments were practiced (Lavelle et al., 2014; Velasquez and Lavelle, 2019).

394 Our study also showed that plantain yield (approximated by bunch weight) in the
395 agroecological system was similar to the conventional systems. It had been showed in a recent study,
396 in plantain cropping systems of Cameroon, that the highest bunch weights were measured in fields
397 belonging to farmers who applied herbicide and nitrogen fertilizers more frequently and at higher
398 rates (D'épigny et al., 2019). In our study, the agroecological practices (use of vermicompost, legume
399 crop cover, healthy plant material) allow to maintain a level of plantain productivity similar to that of
400 conventional systems. The similarity between both systems regarding plantain yield may be also
401 linked to the fact that the soil fertility was similar in the 2 systems.

402

403

404 5. Conclusion

405

406 We verified the hypothesis that there was a fast significant increase in soil macrofauna
407 abundance and diversity and associated ecosystem services (soil aggregation, regulation of pests,
408 production) in agroecological versus conventional plantain systems. The agroecological system
409 improved soil quality and achieved similar plant yield levels than the conventional system, only 10
410 months after the plantation. Agroecological practices, which promote biotic and abiotic interactions
411 in the soil, may contribute to the goal of achieving sustainable tropical cropping systems. The regular
412 communication of the experiment results to the producers and their own analysis developed during
413 several seminars led them to reevaluate their production systems and consider adopting some of the
414 proposed agroecological options. Some of plantain producers of Guadeloupe have adopted PIF

415 plants, which would ideally extend the life of their banana plantations by two years, and organic
 416 fertilization with vermicompost (innovations produced on the farm). However, further research is
 417 necessary to evaluate the long-term impact of agroecological practices in plantain agricultural
 418 systems in particular on chemical soil fertility.

419
 420

421 **Declaration of Competing Interest**

422

423 The authors declare that they have no known competing financial interests or personal
 424 relationships that could have appeared to influence the work reported in this paper

425
 426

427 **Acknowledgments**

428

429 This work was supported financially by the French Ministry of Ecology and Sustainable
 430 Development, France (ECOPHYTO program, project ALTERBIO). The authors thank the three farmers
 431 involved in this study. We also thank Fred Burner, Chantal Fléreau, Danielle Ramaël and Frédérique
 432 Razan (INRAE Antilles Guyane) for their technical help. The authors would also like to thank Lucienne
 433 Desfontaines (INRAE Antilles Guyane) for soil chemical analyses

434
 435

436 **References**

437

- 438 Altieri, M.A., 1999. The ecological role of biodiversity in agroecosystems. *Agric. Ecosyst. Environ.* 74,
 439 19–31. [https://doi.org/10.1016/S0167-8809\(99\)00028-6](https://doi.org/10.1016/S0167-8809(99)00028-6).
- 440 Andres, C., Bhullar, G., 2016. Sustainable intensification of tropical agro-ecosystems: need and
 441 potentials. *Front. Environ. Sci.* 4, 1–10. <https://doi.org/10.3389/fenvs.2016.00005>.
- 442 Blanchart, E., Albrecht, A., Alegre, J., Duboisset, A., Villenave, C., Pashanasi, B., Lavelle, P., Brussaard,
 443 L., 1999. Effects of earthworms on soil structure and physical properties. In: Lavelle, P.,
 444 Brussaard, L., Hendrix, J. (Eds.), *Earthworm Management in Tropical Agroecosystems*. CAB
 445 International, London, pp. 149–172.
- 446 Blouin, M., Zuily-Fodil, Y., Pham-Thi, A.-T., Laffray, D., Reversat, G., Pando, A., Tondoh, G., Lavelle, P.,
 447 2005. Belowground organism activities affect plant aboveground phenotype, inducing plant
 448 tolerance to parasites. *Ecol. Lett.* 8, 202–208. [https://doi.org/10.1111/j.1461-](https://doi.org/10.1111/j.1461-0248.2004.00711.x)
 449 [0248.2004.00711.x](https://doi.org/10.1111/j.1461-0248.2004.00711.x).
- 450 Bolker, B.M., Brooks, M.E., Clark, C.J., Geange, S.W., Poulsen, J.R., Stevens, M.H.H., White, J.S.S.,
 451 2009. Generalized linear mixed models: a practical guide for ecology and evolution. *Trends*
 452 *Ecol. Evol.* 24, 127–135. <https://doi.org/10.1016/j.tree.2008.10.008>.
- 453 Bridge, J., Gowen, S.R., 1993. Visual assessment of plant parasitic nematode and weevil damage on
 454 bananas and plantain, in: Gold C.S., Gemmel, B. (Eds.), *Biological and Integrated control of*
 455 *highland banana and Plantain pests and diseases in Africa*. Cotonou, Benin, Davis, CA, UA, pp.
 456 147–154.
- 457 Brussaard, L., 1998. Soil fauna, guilds, functional groups and ecosystem processes. In: *Appl. Soil Ecol*,
 458 9, pp. 123–135. [https://doi.org/10.1016/S0929-1393\(98\)00066-3](https://doi.org/10.1016/S0929-1393(98)00066-3).
- 459 Carrascosa, M., Sánchez-Moreno, S., Alonso-Prados, J.L., 2014. Relationships between nematode
 460 diversity, plant biomass, nutrient cycling and soil suppressiveness in fumigated soils. *Eur. J.*

- 461 Soil Biol. 62, 49–59. <https://doi.org/10.1016/j.ejsobi.2014.02.009>.
- 462 Chaudhuri, P.S., Paul, T.K., Dey, A., Datta, M., Dey, S.K., 2016. Effects of rubber leaf litter
463 vermicompost on earthworm population and yield of pineapple (*Ananas comosus*) in West
464 Tripura, India. *Int. J. Recycl. Org. Waste Agric.* 5, 93–103. [https://doi.org/10.1007/s40093-](https://doi.org/10.1007/s40093-016-0120-z)
465 016-0120-z.
- 466 Chen, Y., Camps-Arbestain, M., Shen, Q., Singh, B., Cayuela, M.L., 2018. The long-term role of organic
467 amendments in building soil nutrient fertility: a meta-analysis and review. *Nutr. Cycl.*
468 *Agroecosyst.* 111, 103–125. <https://doi.org/10.1007/s10705-017-9903-5>.
- 469 Clermont-Dauphin, C., Cabidoche, Y.M., Meynard, J.M., 2004. Effects of intensive monocropping of
470 bananas on properties of volcanic soils in the uplands of the French west indies. *Soil Use*
471 *Manag.* 20, 105–113. <https://doi.org/10.1111/j.1475-2743.2004.tb00345.x>.
- 472 Clermont-Dauphin C.M., Blanchart E., Loranger-Merciris G., Meynard J.M., 2014. Cropping systems to
473 improve soil biodiversity and ecosystem services: the outlook and lines of research, in: Ozier-
474 Lafontaine H., Lesueur-Jannoyer M. (Eds.), *Sustain. Agric. Rev.* 14, 117–158.
475 https://doi.org/10.1007/978-3-319-06016-3_5.
- 476 Dabin, B., 1967. Sur une méthode d'analyse du phosphore dans les sols tropicaux. Colloque sur la
477 fertilité des sols tropicaux. Tananarive 99–115, 19-25 Nov., I.
- 478 Dendoncker, N., Boeraeve, F., Crouzat, E., Dufrêne, M., König, A., Barnaud, C., 2018. How can
479 integrated valuation of ecosystem services help understanding and steering agroecological
480 transitions ? *Ecol. Soc.* 23, 1–13. <https://doi.org/10.5751/ES-09843-230112>.
- 481 D'épigny, S., Delrieu Wils, E., Tixier, P., Ndoumbé Keng, M., Cilas, C., Lescot, T., Jagoret, P., 2019.
482 Plantain productivity: insights from Cameroonian cropping systems. *Agr. Syst.* 168, 1–10.
483 <https://doi.org/10.1016/j.agsy.2018.10.001>.
- 484 Driessen, P., Deckers, J., Spaargaren, O., Nachtergaele, F., 2001. *Lecture Notes on the Major Soils of*
485 *the World*. FAO, Rome.
- 486 Druart, C., Scheifler, R., de Vaufleury, A., 2010. Towards the development of an embryotoxicity
487 bioassay with terrestrial snails: Screening approach for cadmium and pesticides. *J. Hazard.*
488 *Mater.* 184, 26–33. <https://doi.org/10.1016/j.jhazmat.2010.07.099>.
- 489 Duyck, P.F., Dortel, E., Vinatier, F., Gaujoux, E., Carval, D., Tixier, P., 2012. Effect of environment and
490 fallow period on *Cosmopolites sordidus* population dynamics at the landscape scale. *Bull.*
491 *Entomol. Res.* 102, 583–588. <https://doi.org/10.1017/S0007485312000089>.
- 492 Ekunwe, P.A., Ajayi, H.I., 2010. Economics of plantain production in Edo State Nigeria. *Res. J. Agric.*
493 *Biol. Sci.* 6, 902–905.
- 494 El Mujtar, V., Muñoz, N., Mc Cormick, B.P., Pulleman, M., Tittonell, P., 2019. Role and management
495 of soil biodiversity for food security and nutrition; where do we stand. *Glob. Food Secur.* 20,
496 132–144. <https://doi.org/10.1016/j.gfs.2019.01.007>.
- 497 FAOSTAT, 2021. Food and Agriculture Organization of the United Nations. FAO,, Rome.
- 498 Ferris, H., Bongers, T., 2006. Nematode indicators of organic enrichment. *J. Nematol.* 38, 3–12.
- 499 Godefroid, M., Tixier, P., Chabrier, C., Djigal, D., Qu'énéhervé, P., 2017. Associations of soil type and
500 previous crop with plant-feeding nematode communities in plantain agrosystems. *Appl. Soil*
501 *Ecol.* 113, 63–70. <https://doi.org/10.1016/j.apsoil.2017.01.012>.
- 502 Gold, C.S., 1991. Introduction: biological and integrated control of highland banana and plantain
503 Pests and Diseases. Cotonou, Bénin.
- 504 Gowen, S., Qu'énéhervé, P., Fogain, R., Luc, M., Sikora, R., Bridge, J., 2005. Nematode Parasites of
505 Bananas and Plantains. *Plant Parasitic Nematodes in Subtropical and Tropical Agriculture*.
506 CABI, pp. 611–643. <https://doi.org/10.1079/9780851997278.0000>.

- 507 Griffiths, B.S., Caul, S., Thompson, J., Hackett, C.A., Cortet, J., Pernin, C., Krogh, P.H., 2008. Soil
 508 microbial and faunal responses to herbicide tolerant maize and herbicide in two soils. *Plant*
 509 *Soil* 308, 93–103. <https://doi.org/10.1007/s11104-008-9609-1>.
- 510 ISO, 2011. *Qualité du sol - Prélèvement des invertébrés du sol - Partie 5: Prélèvement et extraction*
 511 *des macro-invertébrés du sol. ISO 23611–5, 2011.*
- 512 Johns, G.G., 1994. Effect of *Arachis pintoi* groundcover on performance of bananas in northern New
 513 South Wales. *Aust. J. Exp. Agr.* 34, 1197–1204. <https://doi.org/10.1071/EA9941197>.
- 514 Kahane, R., Hodgkin, T., Jaenicke, H., Hoogendoorn, C., Hermann, M., Keatinge, D.J.D. H., D'arros,
 515 H.J., Padulosi, S., Looney, N.E., 2013. Agrobiodiversity for food security, health and income.
 516 *Agron. Sustain. Dev.* 33, 671–693.
- 517 Kibblewhite, M.G., Ritz, K., Swift, M.J., 2008. Soil health in agricultural systems. *Philos. T. Roy. Soc. B.*
 518 363, 685–701. <https://doi.org/10.1098/rstb.2007.2178>.
- 519 Laossi, K.M., Barot, S., Carvalho, D., Desjardins, T., Lavelle, P., Martins, M., Mitja, D., Rendeiro, A.,
 520 Rousseau, G., Sarrazin, M., Velasquez, E., Grimaldi, M., 2008. Effects of plant diversity on
 521 plant biomass production and soil macrofauna in Amazonian pastures. *Pedobiologia* 51, 397–
 522 407. <https://doi.org/10.1016/j.pedobi.2007.11.001>.
- 523 Lafont, A., Risède, J.M., Loranger-Merciris, G., Clermont-Dauphin, C., Dorel, M., Rhino, B., Lavelle, P.,
 524 2007. Effects of the earthworm *Pontoscolex corethrurus* on banana plants infected with the
 525 plant-parasitic nematode *Radopholus similis*. *Pedobiologia* 51, 311–318.
 526 <https://doi.org/10.1016/j.pedobi.2007.05.004>.
- 527 Lavelle, P., Blouin, M., Boyer, J., Cadet, P., Laffray, D., Pham-Thi, A.T., Reversat, G., Settle, W., Zuily,
 528 Y., 2004. Plant parasite control and soil fauna diversity. *C. R. Biol.* 327, 629–638.
 529 <https://doi.org/10.1016/j.crv.2004.05.004>.
- 530 Lavelle, P., Rodríguez, N., Arguello, O., Bernal, J., Botero, C., Chaparro, P., Gómez, Y., Gutiérrez, A.,
 531 Hurtado, M., Loaiza, S., Xiomara Pullido, S., Rodríguez, E., Sanabria, C., Velásquez, E., Fonte,
 532 S.J., 2014. Soil ecosystem services and land use in the rapidly changing Orinoco River Basin of
 533 Colombia. *Agric. Ecosyst. Environ.* 185, 106–117.
 534 <https://doi.org/10.1016/j.agee.2013.12.020>.
- 535 Lavelle, P., Spain, A., Blouin, M., Brown, G., Decaëns, T., Grimaldi, M., Jiménez, J.J., McKey, D.,
 536 Mathieu, J., Velasquez, E., Zangerlé, A., 2016. Ecosystem engineers in a self-organized soil: a
 537 review of concepts and future research questions. *Soil Sci.* 181, 91–109.
 538 <https://doi.org/10.1097/SS.0000000000000155>.
- 539 Lavelle, P., Mathieu, J., Spain, A., Brown, G., Fragoso, C., Lapied, E., De Aquino, A., Barois, I., Barrios,
 540 E., Barros, M.E., Bedano, J.C., Blanchart, E., Caulfield, M., Chagueza, Y., Dai, J., Decaens, T.,
 541 Dominguez, A., Dominguez, Y., Feijoo, A., Folgarait, P., Fonte, S.J., Gorosito, N., Huerta, E.,
 542 Jimenez, J.J., Kelly, C., Loranger, G., Marchao, R., Marichal, R., Praxedes, C., Rodriguez, L.,
 543 Rousseau, G., Rousseau, L., Sanabria, C., Suarez, J.C., Tondoh, J.E., De Valenca, A., Vanek, S.J.,
 544 Vasquez, J., Velasquez, E., Webster, E., Zhang, C., 2022. Soil macroinvertebrate communities:
 545 a worldwide assessment. *Glob. Ecol. Biogeogr.* <https://doi.org/10.1111/geb.13492>.
- 546 Lefranc, L.M., Lescot, T., Staver, C., Nkapnang, I., Temple, L., 2010. Macropropagation as an
 547 innovative technology: lessons and observations from projects in Cameroon. *Acta Hort.*
 548 879, 727–734. <https://doi.org/10.17660/ActaHortic.2010.879.78>.
- 549 Lescot, T., 2004. Banane: production, commerce et variétés. *Fruitrop* 118, 5–9.
- 550 Loranger-Merciris, G., Cabidoche, Y.M., Deloné, B., Quénéhervé, P., Ozier-Lafontaine, H., 2012. How
 551 earthworm activities affect banana plant response to nematode parasitism. *Appl. Soil Ecol.*
 552 52, 1–8. <https://doi.org/10.1016/j.apsoil.2011.10.003>.

- 553 Millennium Ecosystem Assessment (MEA), 2005. Ecosystems and human well-being. Island Press,,
554 Washington, D.C.
- 555 Meynard, J.M., 2017. L'agroécologie, un nouveau rapport aux savoirs et à l'innovation. Tropical Issue.
556 24. <https://doi.org/10.1051/ocl/2017021>.
- 557 Mosier, S., Córdova, S.C., Robertson, G.P., 2021. Restoring soil fertility on degraded lands to meet
558 food, fuel, and climate security needs via perennialization. *Front. Sustain. Food Syst.* 5,
559 706142 <https://doi.org/10.3389/fsufs.2021.706142>.
- 560 Niedobová, J., Kristofová, L., Michalko, R., Hula, V., Kýnický, J., Brtnický, M., 2019. Effects of
561 glufosinate-ammonium herbicide and pod sealant on spider *Pardosa agrestis*. *J. Appl.*
562 *Entomol.* 143, 196–203. <https://doi.org/10.1111/jen.12574>.
- 563 Nkonya, E., Mirzabaev, A., von Braun, J., 2016. Economics of land degradation and improvement - a
564 global assessment for sustainable development. Springer International Publishing,, Cham.
- 565 Obiefuna, J.C., 1989. Biological weed control in plantains (*Musa AAB*) with egusi melon (*Colocynthis*
566 *citrullus L.*). *Biol. Agric. Hortic.* 6, 221–227. <https://doi.org/10.1080/01448765.1989.9754519>.
- 567 Ozier-Lafontaine, H., Lesueur-Jannoyer, M., 2014 (Eds). *Sustain. Agric. Rev. 14, Agroecology and*
568 *Global Change.* <https://doi.org/10.1007/978-3-319-06016-3>.
- 569 Pattison, T., Cobon, J., Sikora, R., 2006. Soil quality improvement and nematode management on
570 banana farms in Australia, 15-20 October 2006. In: XVII Reuniao Internacional ACORBAT, Vol.
571 1. Joinville, Santa Catarina,, Brasil, pp. 268–283, 15-20 October 2006.
- 572 Pelosi, C., Barot, S., Capowiez, Y., Hedde, M., Vandenbulcke, F., 2014. Pesticides and earthworms: a
573 review. *Agron. Sustain. Dev.* 34, 199–228. <https://doi.org/10.1007/s13593-013-0151-z>.
- 574 Perin, A., Guerra, J.G.M., Texeira, M.G., 2003. Soil coverage and nutrient accumulation by pinto
575 peanut. *Pesqui. Agropecu. Bras.* 38, 791–796. <https://doi.org/10.1590/S0100-204X2003000700002>.
- 576
- 577 Peters, S., 1991. Organic and conventional beyond transition. *Org. Farmer* 11, 1–5.
- 578 Picq, C. Fouré, E., Frison, E.A., 1998. Bananas and Food Security. Les productions bananières: un
579 enjeu économique majeur pour la sécurité alimentaire. International symposium, Douala,
580 Cameroon, 10–14 November 1998.
- 581 Pretty, J.N., 2008. Agricultural sustainability: concepts, principles and evidence. *Philos. Trans. R. Soc.*
582 *B.* 363, 447–465. <https://doi.org/10.1098/rstb.2007.2163>.
- 583 Puissant, J., Villenave, C., Chauvin, C., Plassard, C., Blanchart, E., Trap, J., 2021. Quantification of the
584 global impact of agricultural practices on soil nematodes: A meta-analysis. *Soil Biol. Bioch.*
585 161, 108383 <https://doi.org/10.1016/j.soilbio.2021.108383>.
- 586 Quénéhervé, P., 2009. Integrated Management Of Banana Nematodes. In: Ciancio, A., Mukerji, K.
587 (Eds.), *Integrated Management of Fruit Crops Nematodes. Integrated Management of Plant*
588 *Pests and Diseases*, vol 4. Springer, Dordrecht. https://doi.org/10.1007/978-1-4020-9858-1_1.
- 589
- 590 Robinson, J.C., 1996. *Banana and Plantains.* Cab. Int., Wallingford, UK.
591 <https://doi.org/10.1079/9781845936587.0000>.
- 592 Rowen, E., Tooker, J.F., Blubaugh, C.K., 2019. Managing fertility with animal waste to promote
593 arthropod pest suppression (Review). *Biol. Control* 134, 130–140.
594 <https://doi.org/10.1016/j.biocontrol.2019.04.012>.
- 595 Sodom, L., Tomekpé, K., Folliot, M., Côte, F.X., 2010. Comparaison de l'efficacité de deux méthodes
596 de multiplication rapide de plants de bananier à partir de l'étude des caractéristiques
597 agronomiques d'un hybride de bananier plantain (*Musa spp.*). *Fruits* 65, 3–9.
598 <https://doi.org/10.1051/fruits/2009036>.

- 599 Sánchez-Moreno, S., Alonso-Prados, E., Alonso-Prados, J.L., García-Baudín, J.M., 2009. Multivariate
600 analysis of toxicological and environmental properties of soil nematicides. *Pest Manag. Sci.*
601 65, 82–92. <https://doi.org/10.1002/ps.1650>.
- 602 Schrama, M., de Haan, J.J., Kroonen, M., Versteegen, H., Van der Putten, W.H., 2018. Crop yield gap
603 and stability in organic and conventional farming systems. *Agric. Ecosyst. Environ.* 256, 123–
604 130. <https://doi.org/10.1016/j.agee.2017.12.023>.
- 605 Sierra, J., Desfontaines, L., Faverial, J., Loranger-Merciris, G., Boval, M., 2013. Composting and
606 vermicomposting of cattle manure and green wastes under tropical conditions: carbon and
607 nutrient balances and end-product quality. *Soil Res* 51, 142–151.
608 <https://doi.org/10.1071/SR13031>.
- 609 Sihi, D., Dari, B., Sharma, D.K., Pathak, H., Nain, L., Sharma, O.P., 2017. Evaluation of soil health in
610 organic vs. conventional farming of basmati rice in North India. *J. Plant Nutr. Soil Sc.* 180,
611 389–406. <https://doi.org/10.1002/jpln.201700128>.
- 612 Tabarant, P., Villenave, C., Risède, J.M., Estrade, J.R., Thuriès, L., Dorel, M., 2011. Effects of four
613 organic amendments on banana parasitic nematodes and soil nematode communities. *Appl.*
614 *Soil Ecol.* 49, 59–67. <https://doi.org/10.1016/j.apsoil.2011.07.001>.
- 615 Tchango, J., Bikoï, A., Achard, R., Escalan, J.V., Ngalani, J.A., 1999. Plantain: postharvest
616 operations. AGSI/FAO, 60 p.
- 617 Tilman, D., Cassman, K.G., Matson, P.A., Naylor, R., Polasky, S., 2002. Agricultural sustainability and
618 intensive production practices. *Nature* 418, 671–677. <https://doi.org/10.1038/nature01014>.
- 619 Thioulouse, J., Dray, S., Dufour, A.B., Siberchicot, A., Jombart, T., 2018. Multivariate analysis of
620 ecological data with ade4. Springer, New York, p. 2018. [https://doi.org/10.1007/978-1-4939-](https://doi.org/10.1007/978-1-4939-8850-1)
621 [8850-1](https://doi.org/10.1007/978-1-4939-8850-1).
- 622 Tsiafouli, M., Thebault, E., Sgardelis, S.P., de Ruiter, P.C., van der Putten, W.H., Birkhofer, K.,
623 Hemerik, L., de Vries, F.T., Bardgett, R.D., Brady, M.V., Bjornlund, L., Jorgensen, H.B.,
624 Christensen, S., D' Hertefeldt, T., Hotes, S., Hol, W.H.G., Frouz, J., Liiri, M., Mortimer, S.R.,
625 Setälä, H., Tzanopoulos, J., Uteseny, K., Pizl, V., Stary, J., Wolters, V., Hedlund, K., 2015.
626 Intensive agriculture reduces soil biodiversity across Europe. *Glob. Change Biol.* 21, 973–985.
627 <https://doi.org/10.1111/gcb.12752>.
- 628 Velasquez, E., Pelosi, C., Brunet, D., Grimaldi, M., Martins, M., Rendeiro, A.C., Barrios, E., Lavelle, P.,
629 2007. This ped is my ped: Visual separation and near infrared spectra allow determination of
630 the origins of soil macroaggregates. *Pedobiologia* 51, 75–87.
631 <https://doi.org/10.1016/j.pedobi.2007.01.002>.
- 632 Velasquez, E., Fonte, S.J., Barot, S., Grimaldi, M., Desjardins, T., Lavelle, P., 2012. Soil macrofauna-
633 mediated impacts of plant species composition on soil functioning in Amazonian pastures.
634 *Appl. Soil Ecol.* 56, 43–50. <https://doi.org/10.1016/j.apsoil.2012.01.008>.
- 635 Velasquez, E., Lavelle, P., 2019. Soil macrofauna as an indicator for evaluating soil based ecosystem
636 services in agricultural landscapes. *Acta Oecol* 100, 103446.
637 <https://doi.org/10.1016/j.actao.2019.103446>.
- 638 Viglierchio, D.R., Schmitt, R.V., 1983. On the methodology of nematode extraction from field
639 samples: Baermann funnel modifications. *J. Nematol.* 15, 438–444.
- 640 Wurst, S., 2010. Effects of earthworms on above- and belowground herbivores. *Appl. Soil Ecol.* 45,
641 123–130. <https://doi.org/10.1016/j.apsoil.2010.04.005>.
- 642 Zhong, Z., Huang, X., Feng, D., Xing, S., Weng, B., 2018. Long-term effects of legume mulching on soil
643 chemical properties and bacterial community composition and structure. *Agric. Ecosyst.*
644 *Environ.* 268, 24–33. <https://doi.org/10.1016/j.agee.2018.09.001>.

645 Zinati, G., 2002. Transition from conventional to organic farming systems: I. Challenges,
646 recommendations, and guidelines for pest management. HortTechnology 12, 606–610.
647 <https://doi.org/10.21273/HORTTECH.12.4.606>.
648
649
650

651

652

653

Table 1

Description of the plantain cropping systems in three studied farms in Guadeloupe (conventional and agroecological practices). Number into brackets following pesticide, herbicide and fertilizer levels are the number of applications during the plantation cycle observed in the plots.

	Farm 1		Farm 2		Farm 3	
	Conventional practices	Agroecological practices	Conventional practices	Agroecological practices	Conventional practices	Agroecological practices
Planting date	03.14.2013		03.08.2013		02.19.2013	
Fallow duration (before plantation)	2 years		4 years		5 years	
Origin of plant material	Sucker	PIF	Sucker	PIF	Sucker	PIF
Plantain cultivar	French		French		French	
Plot longevity	Annual (1 cycle)		Annual (1 cycle)		Annual (1 cycle)	
Pesticides application	Nemathorin® (1)	None	Nemathorin® (1)	None	Nemathorin® (1)	None
⊕ Herbicides application	Basta® F1 (3)	None§	Basta® F1 (3)	None	Reglone® (3)	None§
Mulching	None	<i>Arachis pintoï</i> †	None	<i>Arachis pintoï</i> †	None	<i>Arachis pintoï</i> †
Fertilization	NPK fertilizer (5)	Vermicompost*	NPK fertilizer (5)	Vermicompost*	NPK fertilizer (5)	Vermicompost*

⊕ Herbicides were applied 1 month, 5 months and 8 months after the plantation. Two months occurred between the last herbicide application and soil sampling.

§ manual weeding during the two first months, then brushcutting every 3 weeks.

† density of *Arachis pintoï*: 4 cuttings per m² at the beginning of the experiment.

* 1.5 kg Vermicompost per plant at the beginning of the experiment; 3 kg Vermicompost per plant 2 months after the beginning of the experiment; and 3 kg Vermicompost per plant 6 months after the beginning of the experiment.

673

674

675

676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713

Table 2

Soil macro-invertebrates (individuals.m⁻²) identified at crop flowering stage in two plantain cropping systems in Guadeloupe (conventional and agroecological) and compared using GLMM. Values are means of 9 experiment units. Standard errors are given in parentheses. For each taxonomic group, means with the same superscript letter are not significantly different.

Soil macro- Invertebrates taxonomic group	Conventional cropping systems	Agroecological cropping systems	F _{1,14}	P
Cl. Diplopoda	46 (19) ^a	114 (39) ^a	3.026	0.104
F. Formicidae (ants)	20 (8) ^a	41 (19) ^a	1.067	0.317
infra U. Isoptera (termites) S/O	15 (15) ^a	19 (17) ^a	0.024	0.879
Lumbricina (earthworms)	88 (17) ^a	164 (24) ^b	7.91	0.014
O. Coleoptera	2 (2) ^a	3 (1) ^a	0.085	0.774
O. Coleoptera (larvae)	14 (2) ^a	15 (3) ^a	0.115	0.739
Cl. Chilopoda	14 (7) ^a	19 (11) ^a	0.137	0.716
O. Isopoda	1 (1) ^a	4 (2) ^a	2.041	0.172
O. Dermaptera	0 (0) ^a	3 (2) ^a	2.703	0.120
S/O. Blattaria	1 (1) ^a	2 (1) ^a	0.160	0.694
O. Araneae	2 (1) ^a	5 (2) ^a	4.000	0.063
Cl. Gastropoda	0 (0) ^a	4 (3) ^a	1.455	0.245
Cl. Turbellaria	0 (0) ^a	2 (1) ^a	3.368	0.085
O. Hemiptera	1 (1) ^a	4 (2) ^a	2.041	0.172
O. Orthoptera	2 (1) ^a	1 (1) ^a	0.160	0.694
O. Diptera (larvae)	1 (1) ^a	16 (16) ^a	0.853	0.370
O. Lepidoptera (larvae)	0 (0) ^a	1 (1) ^a	2.286	0.150
O. Thysanoptera	1 (1) ^a	0 (0) ^a	1.000	0.332

714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746

Table 3

Soil chemical characteristics measured at crop flowering stage in two plantain cropping systems in Guadeloupe (conventional and agroecological) and compared using GLMM. Values are means of 9 experiment units. Standard errors are given in parentheses. For each soil chemical characteristic, means with the same superscript letter are not significantly different.

Soil characteristics	Conventional cropping systems	Agroecological cropping systems	F _{1,14}	P
pH-H ₂ O	5.14 (0.20) ^a	5.15 (0.25) ^a	0.002	0.967
pH-KCl	4.45 (0.18) ^a	4.49 (0.28) ^a	0.035	0.853
N-NH ₄ ⁺ (mg.kg ⁻¹) ₁	3.07 (0.75) ^a	4.01 (0.78) ^a	1.416	0.254
N-NO ₃ ⁻ (mg.kg ⁻¹) ₁	2.33 (0.75) ^a	1.42 (0.24) ^a	1.577	0.230
N (%)	0.26 (0.03) ^a	0.25 (0.02) ^a	0.429	0.523
C (%)	2.59 (0.24) ^a	2.71 (0.21) ^a	0.464	0.507
CEC (cmol.kg ⁻¹)	9.74 (0.74) ^a	9.54 (0.79) ^a	0.124	0.730
K (cmol.kg ⁻¹)	0.66 (0.10) ^a	0.59 (0.13) ^a	0.168	0.688
Na (cmol.kg ⁻¹)	0.08 (0.01) ^a	0.08 (0.01) ^a	0.630	0.441
Ca (cmol.kg ⁻¹)	3.76 (0.70) ^a	3.88 (0.89) ^a	0.027	0.871
Mg (cmol.kg ⁻¹)	1.28 (0.14) ^a	1.34 (0.10) ^a	0.390	0.542
Available P (%)	0.07 (0.004) ^a	0.06 (0.002) ^a	0.816	0.380

747
 748
 749
 750
 751
 752
 753
 754
 755
 756
 757
 758
 759
 760
 761
 762
 763
 764
 765
 766
 767
 768
 769
 770
 771
 772
 773
 774
 775
 776
 777
 778
 779




Fig. 1. A) Design of the plantain field trials carried out in each of the three farms located in the western part of Guadeloupe. B) Design of each pair of experimental units. The number within the boxes indicate the plants used for measurements.

780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807




Fig. 2. Morphology of soil aggregates in two plantain cropping systems in Guadeloupe (conventional and agroecological).

808
 809
 810
 811
 812
 813
 814
 815
 816
 817
 818
 819
 820
 821
 822
 823
 824
 825
 826
 827
 828
 829
 830
 831
 832
 833
 834




Fig. 3. Soil macrofauna abundance in two plantain cropping systems in Guadeloupe [conventional and agroecological]. Only the 6 main groups were represented. The abundance of the 9 remaining groups has been gathered in the




Fig. 4. Soil biological properties in cropping systems and farms. Projection in the factorial plane F1F2 of a PCA of biological variables (left) and sampling points separated among cropping systems (upper right, $p < 0.01$) and farms (lower right, $p < 0.01$). LEFT: *Macroinvertebrate community* (DIP: *Diplopoda*, CHI: *Chilopoda*, COL: *Coleoptera*, FOR: *Formicidae*, LUM: *Lumbricina*, MTOT: Total Macrofauna, OTH: Other invertebrates, TAX: Taxonomic richness); Soil macro aggregation status (PMAG: Physical macroaggregates, BMAG: Biogenic macroaggregates, NMAG: non macroaggregated soil); Nematodes (PRAT: *Pratylenchus* sp., MELO: *Meloidogyne* sp., HELI: *Helicotylenchus* sp., RAD: *Radopholus similis*, FLN: Free-living nematodes); RNI: Root necrosis index; YIELD: Plant yield. RIGHT: AGRO (Agroecological system), CONV (Conventional system); 1: Farm 1, 2: Farm 2, 3: Farm 3.