



# **Tuning the pH during the fermentation has a strong effect on the wine protein composition and the stability of the resulting white wines**

Eric Meistermann, Aude Vernhet, Philippe Cottureau, Frédéric Charrier,  
Patrick Chemardin, Céline Poncet-Legrand

## **► To cite this version:**

Eric Meistermann, Aude Vernhet, Philippe Cottureau, Frédéric Charrier, Patrick Chemardin, et al..  
Tuning the pH during the fermentation has a strong effect on the wine protein composition and the  
stability of the resulting white wines. Macrowine 2021, Jun 2021, Virtual, Italy. hal-03712957

**HAL Id: hal-03712957**

**<https://hal.inrae.fr/hal-03712957>**

Submitted on 4 Jul 2022

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



1 – Institut Français de la Vigne et du Vin,  
30240 Le Grau du Roi, France  
2 – SPO, INRAE, Institut Agro, Univ  
Montpellier  
34060 Montpellier, France  
celine.poncet-legrand@inrae.fr

## Tuning the pH during the fermentation has a strong effect on the wine protein composition and the stability of the resulting white wines.

E. Meistermann<sup>1</sup>, A. Vernhet<sup>2</sup>, P. Cottureau<sup>1</sup>, F. Charrier<sup>1</sup>, P. Chemardin<sup>2</sup>,  
C. Poncet-Legrand<sup>2</sup>


**AIM:** The context of global warming implies increases of ethanol content and pH in wines. Elsewhere, previous results have shown the impact of pH on the stability of proteins in white wines. Our aim was to assess for different varieties the effect of pH on the protein content of wines at the end of the alcoholic fermentation. Fermentations were performed using the same starting must, to avoid any other difference in terms of matrix composition (polyphenols, polysaccharides, ions...).

**METHODS:** Fermentations were carried out using musts from different grape varieties (Sauvignon, Sylvaner, Riesling, Gewurztraminer and Pinot Gris), adjusting the pH of the initial musts to 3.0, 3.3, 3.6 and 3.9 either with hydrochloric acid or sodium hydroxide. For each wine thus obtained, the proteins were analyzed and quantified by gel electrophoresis. Heat tests (heating at 40°C during 4 hours) were carried out and the differences between the initial and final turbidities ( $\Delta$ NTU) were calculated.

### RESULTS:

#### SDS – PAGE analysis of proteins in musts and wines

**Invertase 70 kDa**    **$\beta$ -glucanase ~35 kDa**   **Chiti27 : chitinase 27 kDa**   **Chiti 24: chitinase 24 kDa**   **TL22: Thaumatin like protein 22 kDa**  
**TL21: Thaumatin like protein 21 kDa**   **TL19-20: Thaumatin like protein 19-20 kDa**   **LTP: Lipid transfer Proteins ~13-15 kDa**



On the whole and in accordance with previous works, protein concentrations in wines decreased during fermentation.


However, this decrease was more marked for the lowest pHs (3.0 and 3.3).

Furthermore, two families of proteins (chitinases and  $\beta$ -glucanases) were particularly fragile, but their behavior also depended on the initial pH of the must and the matrix composition : for instance chitinases were eliminated in Gewurztraminer and Sauvignon, but not in Pinot Gris, nor Riesling and Sylvaner.

On the contrary, invertases and lipid transfer proteins remained stable when they were initially present.

All these behaviors are in agreement with previous results [1, 2, 3].

#### Evolution of $\Delta$ NTU as a function of the pH



Even though the concentration of proteins either decreased or remained stable when the pH decreased, the turbidity measured after the heat tests evolved differently: a maximum was observed at pH 3.6.

This is probably due to a balance between different phenomena:

- at higher pHs (>3.6) the temperature of denaturation of most proteins is increased above 50°C and are thus not unfolded at 40°C;
- at lower pH (3.0 and 3.3), proteins unfold at lower temperature, but once unfolded, there are more charged and their aggregation is limited due to electrostatic repulsions, limiting the visual haze formation.

### CONCLUSIONS:

This study confirms that pH has a decisive impact on protein composition in white wines, but the latter is not directly related to the risk of haze formation. Other parameters such as the composition in ions, polyphenols, and polysaccharides play a part. They are themselves strongly dependent on the grape variety and cultural practices.

### References

- [1] Dufrechou, M.; Poncet-Legrand, C.; Sauvage, F.-X.; Vernhet, A. Stability of White Wine Proteins: Combined Effect of pH, Ionic Strength, and Temperature on Their Aggregation. *J. Agric. Food Chem.* **2012**, *60*, 1308– 1319, DOI: 10.1021/jf204048j
- [2] Lambri, M.; Dordoni, R.; Giribaldi, M.; Riva Violetta, M.; Giuffrida, M. G. Effect of pH on the Protein Profile and Heat Stability of an Italian White Wine. *Food Res. Int.* **2013**, *54*, 1781– 1786, DOI: 10.1016/j.foodres.2013.09.038
- [3] Dufrechou, M.; Vernhet, A.; Roblin, P.; Sauvage, F.-X.; Poncet-Legrand, C. White Wine Proteins: How Does the pH Affect Their Conformation at Room Temperature?. *Langmuir* **2013**, *29*, 10475– 10482, DOI: 10.1021/la401524w