



HAL
open science

Traditional agroforestry vineyards, sources of inspiration for the agroecological transition of viticulture

Pablo Oliva Oller, Martin Notaro, Erick Langer, Christian Gary

► To cite this version:

Pablo Oliva Oller, Martin Notaro, Erick Langer, Christian Gary. Traditional agroforestry vineyards, sources of inspiration for the agroecological transition of viticulture. terclim2022 XIVth International Terroir Congress 2nd ClimWine Symposium, Jul 2022, Bordeaux, France. , 2022, Proceedings of the XIVth International Terroir Congress 2nd ClimWine Symposium. hal-03716265

HAL Id: hal-03716265

<https://hal.inrae.fr/hal-03716265>

Submitted on 7 Jul 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial - NoDerivatives 4.0
International License



2nd ClimWine Symposium



terclim

July 3 – 8, 2022 | Bordeaux, France

XIVth International Terroir Congress



Book of abstracts

Traditional agroforestry vineyards, sources of inspiration for the agroecological transition of viticulture

Pablo Oliva Oller^{1,2}, Martin Notaro^{3,5}, Erick Langer⁴ and Christian Gary⁵

¹*IDEPRO Desarrollo Empresarial, Tarija, Bolivia*

²*Fundación FAUTAPO, Camargo, Bolivia*

³*CIRAD, UMR ABSys, Montpellier, France*

⁴*School of Foreign Service, Georgetown University, Washington. D.C., USA*

⁵*ABSys, Univ Montpellier, CIHEAM-IAMM, CIRAD, INRAE, Institut Agro, Montpellier, France*

Presenting author: christian.gary@inrae.fr

A unique “terroir” can be found in southern Bolivia, which combines the specific features of climate, topography and altitude of high valleys, with the management of grapevines staked on trees. It is one of the rare remnants of agroforestry viticulture. A survey was carried out among 29 grapegrowers in three valleys, to characterize the structure and management of these vineyards, and identify the services they expect from trees. Farms were small (2.2 ha on average) and 85% of vineyards were less than 1 ha. Viticulture was associated with vegetable, fruit and fodder production, sometimes in the same fields. Molle trees were found in all plots, together with one or two other native tree species. Traditional grapevine varieties such as Negra Criolla, Moscatel de Alejandria and Vicchoqueña were grown with a large range of densities from 1550 to 9500 vines ha⁻¹. From 18 to 30% of them were staked on trees, with 1.2 to 4.9 vines per tree. The management of these vineyards (irrigation, fertilization and grapevine protection) was described, the most particular technical operation being the coordinated pruning of trees and grapevines. Three types of management could be identified in the three valleys. Grapegrowers had a clear idea of the ecosystem services they expected from trees in their vineyards. The main one was protection against climate hazards (hail, frost, flood). Then they expected benefits in terms of pest and disease control, improvement of soil fertility and resulting yield. At last, some producers claimed that tree-staking was quicker and cheaper than conventional trellising. It can be hypothesized then that agroforestry is a promising technique for the agroecological transition of viticulture. Its contribution to the “terroir” of the high valleys of southern Bolivia and its link with the specificities of the wines and spirits produced there remain to be explored.

Keywords: agroforestry, ecosystem services, native trees, criollo grapes, agroecology

Traditional agroforestry vineyards, sources of inspiration for the agroecological transition of viticulture

Pablo Oliva Oller^{1*}, Martin Notaro², Erick Langer³ and Christian Gary⁴

¹IDEPRO and Fundación FAUTAPO, Bolivia - ²Cirad, France - ³Georgetown University, USA - ⁴Inrae, France

*pablorodrigooliva@gmail.com

A unique *terroir* can be found in southern Bolivia, which combines the specific features of climate, topography and altitude of high valleys, with the management of grapevines staked on trees. It is one of the rare remnants of agroforestry viticulture in the world. Winegrowers attribute several ecosystem services to trees in these vineyards. Surveys were carried out in the valleys of Cinti, Cotagaita and Paicho to characterize 29 of these winegrowing systems.



Field and aerial view of a vineyard with grapes staked on trees in the Valley of Cinti (credit: P. Oliva Oller and Google Maps)



Structure and management of agroforestry vineyards in southern Bolivia

Farms were small (2.2 ha on average) and 85% of vineyards were less than 1 ha. Viticulture was associated with vegetable, fruit and fodder production, sometimes in the same fields. Molle trees were found in all plots, together with one or two other native tree species. Traditional grapevine varieties such as Negra Criolla, Moscatel de Alejandria and Vicchoqueña were grown with a large range of densities from 1550 to 9500 vines ha⁻¹. From 18 to 30% of them were staked on trees, with 1.2 to 4.9 vines per tree.

Tree and grapevine densities in vineyards located in 3 high-valleys of southern Bolivia

	Cinti	Cotagaita	Paicho
tree density (no./ha)	240-440	330-950	640-1020
vine density (no./ha)	3050-9500	5100-7200	1550-6650
tree-staked vines (%)	18	30	28
no. vines/tree	2.3	4.9	1.2

Frequency of tree species, grape cultivars (local names) and farming practices over 29 agroforestry vineyards. Several tree species and grape cultivars could be grown in the same plot.

Tree species	No.	Grape cultivars	No.	Farming practices	No.
Molle (<i>Schinus molle</i>)	29	Negra criolla	26	Irrigation	29
Chañar (<i>Geoffroea decorticans</i>)	10	Moscatel de Alejandria	17	Winter pruning	29
Taco (<i>Prosopis alba</i>)	6	Vicchoqueña	12	Green pruning	29
Algarrobo (<i>Ceratonia siliqua</i>)	4	Pedro Ximenez	1	Tree pruning	29
Churqui (<i>Acacia caven</i>)	2	Cabernet	1	Manure supply	29
Peach tree (<i>Prunus persica</i>)	1			Pesticide appl.	27
Nogal (<i>Juglans australis</i>)	1			Tillage	7
				Mineral fertil.	2

The management of these vineyards (irrigation, fertilization and grapevine protection) was described, the most particular technical operation being the coordinated pruning of trees and grapevines. Three types of management could be identified in the three valleys.

Ecosystem services expected by winegrowers from trees in their vineyards

The main ecosystem service expected from trees by grape growers was the protection against climate hazards (hail, frost, flood). Then they expected benefits in terms of pest and disease control, improvement of soil fertility and resulting grape yield. At last, some producers claimed that tree-staking was quicker and cheaper than conventional trellising.

It can be hypothesized then that agroforestry is a promising technique for the agroecological transition of viticulture. Its contribution to the "terroir" of the high valleys of southern Bolivia and its link with the specificities of the wines and spirits produced there remain to be explored.

Frequency of services expected from trees by winegrowers over 29 agroforestry vineyards. Several services could be expected by the same winegrower.

