



**HAL**  
open science

## Apiaceae Family as a Valuable Source of Biocidal Components and their Potential Uses in Agriculture

Punniamoorthy Thiviya, Niroshan Gunawardena, Ashoka Gamage, Terrence Madhujith, Othmane Merah

► **To cite this version:**

Punniamoorthy Thiviya, Niroshan Gunawardena, Ashoka Gamage, Terrence Madhujith, Othmane Merah. Apiaceae Family as a Valuable Source of Biocidal Components and their Potential Uses in Agriculture. *Horticulturae*, 2022, 8 (7), pp.614. 10.3390/horticulturae8070614 . hal-03726418

**HAL Id: hal-03726418**

**<https://hal.inrae.fr/hal-03726418>**

Submitted on 18 Jul 2022

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution| 4.0 International License



Review

# Apiaceae Family as a Valuable Source of Biocidal Components and their Potential Uses in Agriculture

Punniamoorthy Thiviya <sup>1</sup>, Niroshan Gunawardena <sup>2</sup>, Ashoka Gamage <sup>3,\*</sup>, Terrence Madhujith <sup>4,\*</sup> and Othmane Merah <sup>5,6,\*</sup>

<sup>1</sup> Postgraduate Institute of Agriculture, University of Peradeniya, Peradeniya 20400, Sri Lanka; thiviya904@gmail.com

<sup>2</sup> Department of Food, Agriculture and Bioresources, School of Environment, Resources and Development, Asian Institute of Technology, Khlong Nueng 12120, Thailand; nirosan170@gmail.com

<sup>3</sup> Chemical and Process Engineering, Faculty of Engineering, University of Peradeniya, Peradeniya 20400, Sri Lanka

<sup>4</sup> Department of Food Science and Technology, Faculty of Agriculture, University of Peradeniya, Peradeniya 20400, Sri Lanka

<sup>5</sup> Laboratoire de Chimie Agro-industrielle (LCA), Institut National de la Recherche Agronomique, Université de Toulouse, CEDEX 4, 31030 Toulouse, France

<sup>6</sup> Département Génie Biologique, IUT Paul Sabatier, Université Paul Sabatier, 32000 Auch, France

\* Correspondence: ashogamage@gmail.com (A.G.); tmadhujith@agri.pdn.ac.lk (T.M.); othmane.merah@ensiacet.fr (O.M.); Tel.: +94-714430714 (A.G.); +94-713412171 (T.M.); +33-5-3-432-3523 (O.M.)

**Abstract:** Synthetic chemicals are used to protect crops and agricultural products, thereby producing high yields. However, intensive use of these synthetic chemicals significantly affects the environment and sustainable agriculture production. Moreover, direct or indirect exposure to these synthetic chemicals may cause acute or chronic toxicity in humans and animals. Due to their biodegradability, low toxicity, and being environmentally friendly, secondary metabolites derived from plant sources are being studied as a sustainable approach. Apiaceae family crops are a good source of bioactive phytochemicals. Many studies have found that Apiaceae extracts and essential oils possess various biocidal activities: antibacterial, antifungal, herbicidal, insecticidal or repellent, and larvicidal activities, among others. These various potent bioactivities make the Apiaceae an excellent alternative source for synthetic chemicals. In this context, the present review highlights the biocidal activities of some Apiaceae species and their potential applications in agriculture to protect the plant and agricultural products against pests, weeds, phytopathogens, and foodborne and food spoilage microorganisms.

**Keywords:** Antibacterial; antifungal; biocides; essential oil; herbicides; insecticides; pesticides; phytopathogens; Umbelliferae



**Citation:** Thiviya, P.; Gunawardena, N.; Gamage, A.; Madhujith, T.; Merah, O. Apiaceae Family as a Valuable Source of Biocidal Components and their Potential Uses in Agriculture. *Horticulturae* **2022**, *8*, 614. <https://doi.org/10.3390/horticulturae8070614>

Academic Editor: Stefania Galletti

Received: 1 June 2022

Accepted: 5 July 2022

Published: 7 July 2022

**Publisher's Note:** MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



**Copyright:** © 2022 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

## 1. Introduction

Apiaceae (formerly Umbelliferae) is one of the largest plant families in the order Apiales. Commonly known as parsley or carrot family, it consists of nearly 3780 species belonging to 434 genera distributed throughout many regions in the world [1]. The Apiaceae family includes some of the widely used vegetables and aromatic herbs such as carrot (*Daucus carota*), parsnip (*Pastinaca sativa*), celery (*Apium graveolens*), Gotu kola (*Centella asiatica*), parsley (*Petroselinum crispum*), coriander (*Coriandrum sativum*), dill (*Anethum graveolens*), fennel (*Foeniculum vulgare*), cumin (*Cuminum cyminum*), anise (*Pimpinella anisum*), caraway (*Carum carvi*), as well as poisonous species like water hemlock (*Cicuta maculata*), northern water hemlock (*Cicuta virosa*), poison hemlock (*Conium maculatum*), *Aethusa cynapium* (fool's parsley), and hemlock water-dropwort (*Oenanthe crocata*). Plants of the Apiaceae family are grown, not only as a nutritional source, but also as a source of flavour, fragrance, and medicine. Different parts, such as the root, leaf, leaf stalk, pseudo-bulb, and seed, serve

these purposes. All parts of Apiaceae plants contain secretory glands that are important in storing essential oils (EOs), giving rise to the distinct flavour of each species [2–4]. Moreover, EO plays a vital role in plants: defence against bacteria, viruses, fungi, insects, and herbivores, and as an attractant to pollinators [5].

Agrochemicals, such as pesticides, fertilizers, and chemical growth agents are widely used to increase crop yield and manage weeds, pests, and diseases. Extensive use of these synthetic agrochemicals poses potential risks to human health and the environment, including agrochemical poisoning, negative impact on biocontrol agents, reduced biodiversity and ecosystem functions, residues in the environment and water and soil pollution, food, and drinking water, accumulation in the food chain, high and acute toxicity in humans, occupational and non-occupational health impacts, and pest-resistance development. Therefore, the development of non-toxic, biodegradable, and eco-friendly natural agrochemicals as an alternative to synthetic biocides gained much attention [6–8].

In recent years, the extraction of active compounds (secondary metabolites) from plant sources gained interest among scientists. Plants synthesize secondary metabolites in response to defence mechanisms against biotic or abiotic stress. These secondary metabolites protect plants from phytopathogens (fungi, bacteria, viruses, nematodes, protozoa), predators, UV-B radiation, and drought, and play a role in growth and development. Plants produce several secondary metabolites, such as phenolics, terpenes, carotenoids, alkaloids, and other nitrogen/sulphur-containing compounds [9,10]. Various plant extracts and secondary metabolites are known to have various bioactivities, such as herbicidal, insecticidal, and antimicrobial activities [11]. Medicinal aromatic plants contain high concentrations of these secondary metabolites or bioactive compounds and can be used in pharmaceuticals, agrochemicals, food additives, and as ingredients in cosmetics [11].

Apiaceae species are a good source of secondary metabolite compounds with diverse biological activities, such as apoptosis inducers, antibacterial, antifungal, phytotoxic activities, and cyclooxygenase inhibitory activity [12,13]. Thus, Apiaceae can be used in agriculture as a biopesticide, sprouting inhibitor, antifungal agent, and insect repellent [14]. In this respect, this current review focuses on the Apiaceae as a source of biocidal components and their potential uses in agriculture. Relevant articles published in English were searched in different databases, specifically, Google Scholar, Scopus, Wiley Online Library, Semantic Scholar, and Pub Med. This review cites data from 150 articles, published between 2005 and 2021, and compiles the biocidal potential of 54 Apiaceae species.

## 2. Phytochemicals in Apiaceae Extracts

Several Apiaceae species are an excellent source of EOs, which contain more than 760 different chemical components [1,2]. The chemical composition varies with the plant part used for extraction, various extraction methods, phenological stages of the plant, harvesting season, plant age, soil nature, and environmental conditions. These variations in chemical composition have a direct effect on their biological activities [15,16]. Knowledge of these chemical constituents is vital to make use of this economically important plant family, not only for medicinal benefits, but also for environmental applications. Essential oils and their components, mainly monoterpenoids and sesquiterpenoids, show antimicrobial, repellent, chemosterilant, antifeeding, and other biocidal activities [17,18]. Table 1 shows some Apiaceae species and their major components, while Figure 1 shows the chemical structure of several main components.

**Table 1.** Phytochemicals in Apiaceae plant extracts.

Botanical Name	Common Name	Part	Extract	Main Compounds with Biocidal Activity	Reference
<i>Aegopodium podagraria</i>	Goutweed Ground Elder	Leaf	Ethanollic extracts	Quercetin-3-O-glucoside (399 mg/kg), apigenin (1.3 mg/kg), rutin (0.7 mg/kg)	[19]
<i>Ammi visnaga</i>	Toothpick weed, Khella	Seeds	Ethanollic extracts	Quercetin-3-O-glucoside (49.5 mg/kg), rutin (24 mg/kg), kaempferol (11 mg/kg), biochanin A (40 mg/kg), genistein (23 mg/kg)	[19]
<i>Ammodaucus leucotrichus</i>	Woolly Cumin	Aerial parts	Essential oil	Perillaldehyde (58.3%), D-limonene (23.33%), $\alpha$ -pinene (5.74%), $\beta$ -pinene (2.26%)	[20]
<i>Anethum graveolens</i>	Dill	Seeds	Essential oil	Carvone (42%), limonene (32%), dill ether, $\alpha$ -phellandrene (14.2), cymene (2%)	[21]
				Carvone (55.2%), limonene (16.6%), dill apiole (14.4%), linalool (3.7%)	[22]
<i>Angelica archangelica</i>	Garden Angelica	Root	Essential oil	$\alpha$ -Pinene (29.7%), $\delta$ -3-carene (14.2%), $\beta$ -phellandrene + limonene (13.2%)	[23]
<i>Angelica sylvestris</i>	Wild angelica	Leaf	Ethanollic extract	Quercetin 3-O-glucoside (20.5 mg/kg), rutin (136 mg/kg), formononetin (12.4 mg/kg)	[19]
<i>Anthriscus cerefolium</i>	Chervil	Leaf	Ethanollic extract	Naringin (6.5 mg/kg), quercetin 3-O-glucoside (5.3 mg/kg)	[19]
<i>Anthriscus sylvestris</i>	Cow parsley, Wild chervil	Leaf	Ethanollic extract	Rutin (8.0 mg/kg), quercetin 3-O-glucoside (3.8 mg/kg)	[19]
<i>Apium graveolens</i>	Celery	Leaf, Root	CO <sub>2</sub> -extract	Leaf: Limonene (32.16%) Root: Limonene (21.87%), carvone (17.71%)	[24]
<i>Apium graveolens</i>	Celery	Leaves	Essential oil	$\beta$ -Pinene (39.63%), limonene (15.11%), cis-ocimene (16.18%), $\gamma$ -terpinene (7.73%), $\beta$ -selinene (3.81%)	[25]
<i>Apium nodiflorum</i>	Water celery	Aerial parts	Essential oil	Myristicin (47.0%), limonene (16.7%), terpinolene (9.9%), (Z)- $\beta$ -ocimene (6.1%)	[26]
<i>Azorella cryptantha</i>		Aerial parts	Essential oil	$\alpha$ -Pinene (9.6–21.9%), $\alpha$ -thujene (5.7–12.5%), $\beta$ -pinene (1.5–5.9%)	[27]
				Carvone (70.1%), $\gamma$ -terpinene (12.6%), limonene (5.1%)	[28]
<i>Carum carvi</i>	Caraway	Seeds	Essential oil	Carvone (66.4%), limonene (32.5%)	[29]
				$\gamma$ -Terpinene (31.03%), $\beta$ -pinene (18.77%), p-cymene (17.16%), carvone (12.20%)	[30]

Table 1. Cont.

Botanical Name	Common Name	Part	Extract	Main Compounds with Biocidal Activity	Reference
<i>Carum carvi</i>	Caraway	Fruits	Essential oil	2-Caren-10-al (34.03%), anethole (28.46%), d-terpinene (16.06%), p-cymene (5.59%), limonene (3.55%)	[31]
				Carvone (50.6%), limonene (46.48%)	[32]
				(R)-Carvone (37.98%), D-limonene (26.55%), $\alpha$ -pinene (5.21), cis-carveol (5.01%), $\beta$ -myrcene (4.67%)	[33]
				Carvone (46.62%), limonene (45.49%)	[34]
<i>Carum nigrum</i>		Seeds	Essential oil	Dillapiole (29.9%), germacrene B (21.4%), $\beta$ -caryophyllene (7.8%)	[35]
<i>Carum copticum</i>	Ajowan	Fruits	Essential oil	Thymol (63.17%), p-cymene (21.4%), d-terpinene (13.76%)	[31]
<i>Centella asiatica</i>	Gotu Kola, Centella, Indian penny	Leaves	Essential oil	$\alpha$ -Humulene (21.06%), $\beta$ -caryophyllene (19.08%), bicyclogermacrene (11.22%), germacrene B (6.29%), myrcene (6.55%), $\gamma$ -terpinene (5.77%)	[36]
<i>Chaerophyllum aromaticum</i>		Root	Essential oil and methanolic extract	Viridiflorol (22.2%), germacrene d (5.1%), m-cresol (1.6%), limonene (2.1%), $\alpha$ -pinene (0.8%)	[37]
<i>Chaerophyllum aureum</i>	-	Fruits and aerial parts	Essential oil	Sabinene (18.5–31.6%), p-cymene (7.9–25.4%) and limonene (1.9–10.9%)	[38]
<i>Chaerophyllum crinitum</i>	-	Aerial parts	Essential oil	$\alpha$ -Terpinolene (20.3%), $\alpha$ -terpineol (7.2%), limonene (5.8%)	[39]
<i>Chaerophyllum macropodium</i>	-	Aerial parts	Essential oils and hexane extracts	Trans- $\beta$ -ocimene (21.2–22.8%), cis- $\beta$ -ocimene (8.5–10.0%), p-cymene (1.4–10.3%), $\beta$ -phellandrene (5.1–7.5%), $\gamma$ -terpinene (1.1–9.2%), $\beta$ -pinene (2.3–7.1%), (+) spathulenol (4.4–6.0%)	[40]
<i>Coriandrum sativum</i>	Coriander	Seeds	Essential oil	Linalool (40.9%), geranyl acetate (12.8%), $\gamma$ -terpinene (10.6%)	[28]
				Linalool (66.8%), $\alpha$ -pinene (7.79%), camphor (6.46%), terpinene (3.97%), limonene (3.79%)	[41]
				Linalool (57.57%), geranyl acetate (15.09%), Camphor (3.02%), p-cymene (2.52%)	[42]

Table 1. Cont.

Botanical Name	Common Name	Part	Extract	Main Compounds with Biocidal Activity	Reference
				Linalool (58.80%), menthol (12.89%), $\alpha$ -pinene (5.29%), $\gamma$ -terpinene (4.76%)	[43]
				Linalool (76.41%), $\gamma$ -terpinene (5.35%), $\alpha$ -pinene (4.44%), Camphor (2.20%)	[30]
<i>Coriandrum sativum</i>	Coriander	Fruits	Essential oil	Linalool (70.9%), $\alpha$ -pinene (4.17%), p-cymene (3.63%)	[32]
				$\gamma$ -Terpinene (33.6 %), sabinene (32.0 %), thymol methyl ether (15.7 %)	[44]
<i>Crithmum maritimum</i>	Sea Fennel	Aerial parts	Essential oil	$\beta$ -myrcene (13.66%), p-cymene (11.67%), $\beta$ -phellandrene (6.57%), $\alpha$ -pinene (5.51%), camphene (5.16%)	[45]
<i>Crithmum maritimum</i>	Sea Fennel	-	Essential oil	Sabinene (49.45%), $\gamma$ -terpinene (31.37%), pinenes (9.57%), limonene (2.73%)	[46]
<i>Crithmum maritimum</i>	Sea fennel	Aerial parts and seeds	Essential oil	Dillapiole (55.7 and 39.9%), myristicin (4.4 and 12.8%), $\gamma$ -terpinene (14.0 and 21.2%), thymol methyl ether (11.8 and 11.1%), $\alpha$ -pinene (2.3 and 4.7%), sabinene (4.7 and 1.6%), p-cymene (3.5 and 4.6%), respectively.	[47]
<i>Crithmum maritimum</i>	Sea fennel	Seeds	Essential oil	Dillapiol (39.9%), $\gamma$ -terpinene (21.2%), myristicin (12.8%), thymol methyl ether (11.1%), $\alpha$ -pinene (4.7%), p-cymene (4.6%)	[48]
<i>Crithmum maritimum</i>	Sea fennel	Leaves	Essential oil	$\alpha$ -Phellandrene (71.05%), dill ether (10.83%), limonene (10.74%), p-cymene (2.12%)	[25]
				$\beta$ -Pinene, p-cymene, $\gamma$ -terpinene, cuminal, $\alpha$ -terpinen-7-al and $\gamma$ -terpinen-7-a	[16]
<i>Cuminum cyminum</i>	Cumin	Seeds	Essential oil	$\gamma$ -Terpinen-7-al (35.3%), cuminaldehyde (21.8%), $\alpha$ -terpinen-7-al (15.4%), $\gamma$ -terpinene (12.5%), $\beta$ -pinene (6.4%)	[49].
				Cuminaldehyde (30.42–33.24 %), $\gamma$ -terpinen-7-al (20.54–28.36 %), $\alpha$ -terpinen-7-al (about 13 %), $\gamma$ -terpinene (6.15–12.60 %), $\beta$ -cymene (4.19–5.38 %), $\beta$ -pinene (3.10–5.36 %)	[50]

Table 1. Cont.

Botanical Name	Common Name	Part	Extract	Main Compounds with Biocidal Activity	Reference
<i>Cuminum cyminum</i>	Cumin	Fruits	Essential oil	Cuminaldehyde (25.17%), p-cymene (17.5%), $\beta$ -pinene (13.56%)	[32]
<i>Daucus carota</i>	Wild carrot	Leaf	Ethanollic extract	Quercetin-3-O-glucoside (68.4 mg/kg), rutin (9.5 mg/kg)	[19]
<i>Deverra scoparia</i>		Aerial parts	Essential oil	$\alpha$ -Pinene (31.95%), sabinene (17.24%), $\Delta$ 3-carene (16.85%), ocimene (9.75%), myrcene (3.46%), terpinene-4-ol (2.84%), eugenol (1.8%)	[51]
<i>Echinophora spinosa</i>	Prickly Parsnip	Air dried plants	Essential oil	$\delta^3$ -carene (60.86%), $\alpha$ -phellandrene (7.12%), p-cymene (6.22%), myrcene (4.82%), $\beta$ -phellandrene (2.73%)	[52]
<i>Eryngium alpinum</i>	Alpine Sea Holly	Aerial parts	Essential oil	Caryophyllene oxide (27.9%), bicyclogermacrene (13.2%), germacrene D (8.2%)	[53]
<i>Eryngium amethystinum</i>	Amethyst Sea Holly	Aerial parts	Essential oil	$\beta$ -Caryophyllene (15.2%), $\alpha$ -pinene (10.2%), 2,3,6-trimethylbenzaldehyde (9.3%)	[53]
<i>Eryngium triquetrum</i>	Choukzerk	Flowers	Essential oil	Pulegone (50.6%), piperitenone (30.5%), menthone (7.0%), limonene (1.3%)	[54]
<i>Eryngium triquetrum</i>	Choukzerk	Aerial parts	Essential oil	Falcarinol (74.8%), octane (5.6%)	[55]
<i>Ferula orientalis</i>		Aerial parts	Essential oil	$\alpha$ -Pinene (75.9%), camphene (3.4%), p-cymene (2.2%)	[56]
<i>Ferulago sandrasica</i>		Roots	Essential oil	Limonene (28.9%), $\alpha$ -pinene (15.6%), terpinolene (13.9%)	[56]
<i>Ferulago angulata</i>		Seeds	Essential oil	(Z)- $\beta$ -Ocimene (19.93%), $\alpha$ -pinene (15.50%), p-cymene (7.67%), sabinene (7.49%), $\beta$ -phellandrene (5.5%), $\alpha$ -phellandrene (4.95%), $\gamma$ -terpinene (3.3%)	[57]
<i>Ferulago cassia</i>		Root	Dichloromethane (CH <sub>2</sub> Cl <sub>2</sub> ) extract	Coumarins; peucedanol, suberosin, grandivitol, umbelliferone	[58]
<i>Ferulago trachycarpa</i>		Rhizomes of plant	Dichloromethane, n-hexane, and methanolic extract	Coumarins; crenulatin, suberosin, marmesin senecioate	[59]
<i>Ferulago trifida</i>		Flowers	Essential oil	(E)- $\beta$ -Ocimene (37.3%), $\alpha$ -pinene (16.3%), bornyl acetate (9.4%), cis-verbenol (8.6%), $\gamma$ -terpinene (7.5%), $\alpha$ -limonene (4.3%), $\beta$ -myrcene (2.8%)	[60]


Table 1. Cont.

Botanical Name	Common Name	Part	Extract	Main Compounds with Biocidal Activity	Reference	
<i>Foeniculum vulgare</i>	Fennel	Stems		$\alpha$ -Pinene (22.6%), (E)- $\beta$ -ocimene (20.7%), trans-verbenol (22.1%), bornyl acetate (8.5%), $\alpha$ -limonene (3.3%)		
		Leaves		(E)- $\beta$ -Ocimene (22.6%), $\alpha$ -pinene (19.6%), bornyl acetate (16.7%), cis-verbenol (10.4%), $\beta$ -myrcene (3.2%)		
		Fruits		(E)- $\beta$ -Ocimene (30.5%), $\alpha$ -pinene (18.0%), bornyl acetate (11.0%), $\alpha$ -terpinolene (9.4%)		
			Essential oil		Estragole (50.1%), limonene (20.2%)	[28]
			Essential oil		Trans-anethole (65.1%), fenchone (25.6%), methyl chavicol (3.4%), $\alpha$ -pinene (1.5%), limonene (1.3%)	[21]
			Essential oil		Trans-anethole, fenchone, anisketone, p-anisaldehyde, d-limonene, estragol, estragol, carvone	[61]
			Essential oil		Estragole (71.64%)	[62]
		Essential oil		E-Anethol (76.22%), estragole (9.54%), fenchone (10.91%)	[43]	
		Essential oil		Estragole (43.38%), anethole (29.34%), fenchone (15.16%), Limonene (5.02%)	[63]	
			Ethanollic extract	Rutin (62 mg/kg), quercetin 3-O-glucoside (60.0 mg/kg)	[19]	
<i>Foeniculum vulgare</i>	Fennel	Fruits	Essential oil	Trans-Anethole (64.42%), fenchone (14.59%), methyl cavicol (6.62%), limonene (3.37%)	[31]	
<i>Foeniculum vulgare</i>	Fennel	Leaves	Essential oil	E-Anethol (41.18%), z-anethol (26.49%), $\beta$ -phellandrene (5.38%), fenchone (2.46%)	[25]	
<i>Heracleum anisactis</i>		Aerial parts	Essential oil	Myristicin (95.15%)	[64]	
<i>Heracleum transcaucasicum</i>		Aerial parts	Essential oil	Myristicin (96.87%)	[64]	
<i>Hippomarathrum microcarpum</i>		Aerial parts	Essential oil	$\beta$ -Caryophyllene (31.4%), caryophyllene oxide (23.1%), $\beta$ -phellandrene (4.6%), Germacrene D (4.2%), $\alpha$ -pinene (3.0%)	[56]	
<i>Levisticum officinale</i>	Lovage	leaf and seed	Essential oil	$\gamma$ -Terpinene (14.52% and 12.37%), $\beta$ -phellandrene (13.85% and 15.54%), (Z)- $\beta$ -ocimene (12.91% and 23.70%)	[65]	



Table 1. Cont.

Botanical Name	Common Name	Part	Extract	Main Compounds with Biocidal Activity	Reference
<i>Ostericum sieboldii</i>	Water Dropwort			Myristicin (30.31%), $\alpha$ -terpineol (9.92%), $\alpha$ -cadinol (7.29%), $\beta$ -farnesene (6.26%), linalool (5.94%)	[66]
<i>Pastinaca sativa</i>	Wild parsnip	Leaf	Ethanollic extract	Rutin (652 mg/kg), quercetin 3-O-glucoside (517 mg/kg)	[19]
<i>Petroselinum hortense</i>	Parsley	Seeds	Essential oil	$\alpha$ -Pinene (42.15%), $\beta$ -pinene (30.21%), $\beta$ -phellandrene (6.03%), myristicine (4.37%), limonene (2.02%), sabinene (2.26%), myrtenal (2.14%)	[43]
<i>Pimpinella anisum</i>	Anise	Seeds	Essential oil	E-Anethol (76.56%), estragole (13.01%), linalool (7.42%)	[43]
			Essential oil	Trans-anethole (80.8%), 1,8-cineol (4.7%), linalool (1.5%)	[21]
			Essential oil	(E)-Anethole (93%)	[49].
			Ethanollic extract	Rutin (13.0 mg/kg), quercetin 3-O-glucoside (6.5 mg/kg)	[19]
<i>Pimpinella saxifraga</i>	Burnet saxifrage	Leaf	Ethanollic extract	Quercetin 3-O-glucoside (421.2 mg/kg), rutin (205.2 mg/kg), apigenin (62.0 mg/kg), quercetin (20.4 mg/kg)	[19]
<i>Prangos peucedanifolia</i>		Flower	Essential oil	$\beta$ -pinene (35.58%), $\alpha$ -pinene (22.13%), $\beta$ -phellandrene (12.54%), myrcene (8.27%), $\gamma$ -terpinene (5.97%), $\alpha$ -phellandrene (3.23%)	[67]
		Leaves	Essential oil	m-Cresol (50.38%), trans-p-menth-2-en-1-ol (6.63%), $\gamma$ -terpineol (3.9%), $\alpha$ -terpinen-7-al (3.83%), and $\beta$ -pinene (3.63%)	[67]
<i>Seseli gummiferum</i>	Moon Carrot	Aerial parts	Methanollic and water extracts	Chlorogenic acid (211 and 199 $\mu$ g/g), Gallic acid (37.81 and 22.27 $\mu$ g/g), p-coumaric acid (18.69 and 27.16 $\mu$ g/g), rutin (15.22 and 14.42 $\mu$ g/g)	[68]
<i>Sison amomum</i>	Stone Parsley	Flowering aerial parts	Essential oil	Sabinene (54.4%), $\beta$ -phellandrene (16.6%), germacrene D (6.7%), terpinen-4-ol (3.8%), $\gamma$ -terpinene (2.4%), myrcene (2.0%)	[69]
<i>Smyrniolum olusatrum</i>	Alexander, Horse parsley	Roots	Essential oil	Furanoeremophilone (31.5%), furanodiene (19.1%), (E)- $\beta$ -caryophyllene (11%), $\beta$ -pinene (3%)	[55]
<i>Trachyspermum ammi</i>	Ajwain, Ajowan caraway	Seeds	Essential oil	Thymol (63.4%), p-cymene (19%), $\gamma$ -terpinene (16.9%)	[70]



**Figure 1.** Chemical structure of several major components with potential biocidal activity;

### 3. Biocidal Effects of Apiaceae Extracts

The biocidal activities of phytochemicals, such as monoterpene, phthalides, terpenoids, phenylpropanoids (coumarins and phenylpropenes), and polyacetylenes, are commonly found in Apiaceae plants [4,5]. Compared with single components, oil in its complete composition showed high antimicrobial activity due to the synergistic effect of minor and major constituents of oil [21]. Table 2 highlights the biocidal potential of several Apiaceae species.

**Table 2.** Phytochemical constituents of Apiaceae plant extracts.

Botanical Name	Common Name	Biocidal Effect	Results/Mechanisms	References
<i>Actinolema macrolema</i>		Antibacterial activity ( <i>Staphylococcus epidermidis</i> ) Antifungal activity ( <i>Candida albicans</i> )	MIC using microdilution broth assay 62.5 ( <i>S. epidermidis</i> ) 125 ( <i>C. albicans</i> ) $\mu\text{g/mL}$ .	[71]
<i>Aegopodium podagraria</i>	Goutweed Ground Elder	Antibacterial activity ( <i>Enterobacter cloacae</i> , <i>Klebsiella pneumonia</i> , <i>Pseudomonas fluorescens</i> , <i>Bacillus subtilis</i> , <i>Bacillus mycoides</i> , <i>Staphylococcus aureus</i> )	MIC using tube dilution method 1.25–5.00 mg/mL (in ethanol extract).	[72]
		Larvicidal activity ( <i>Culex quinquefasciatus</i> )	LD <sub>50</sub> = 9 ppm.	[73]
		Insecticidal activity ( <i>Toxoptera aurantii</i> )	LD <sub>50</sub> of seed extract 0.054 ng/aphids	[74]
<i>Ammi visnaga</i>	Toothpick Weed, Khella	Larvicidal activity ( <i>Aedes aegypti</i> )	Khellin, a major compound showed 100% and 93% mortality at 1 $\mu\text{g}/\mu\text{L}$ and at 0.5 $\mu\text{g}/\mu\text{L}$ .	[75]
		Insecticidal activity ( <i>Schistocerca gregaria</i> )	LC <sub>50</sub> of ethanol, petroleum ether and n-butanol extracts (21.0, 12.0 and 22.5%, respectively).	[76]

Table 2. Cont.

Botanical Name	Common Name	Biocidal Effect	Results/Mechanisms	References
<i>Ammodaucus leucotrichus</i>	Woolly Cumin	Antibacterial ( <i>Klebsilla pneumonia</i> ) and Antifungal ( <i>Trichophyton rubrum</i> , <i>Candida albicans</i> ) activities	Inhibition zones in disc diffusion method 6–20 mm.	[77]
		Insecticidal activity ( <i>Tribolium castaneum</i> )	Fumigant toxicity (LC <sub>50</sub> 14.78 µL), repellency, and reduces oviposition potential.	[78]
		Insecticidal activity ( <i>Lymantria dispar</i> )	Antifeedant activity at concentration of 0.5 and 1% of EO (Tot = 59.18 and 56.12, respectively).	[79]
<i>Anethum graveolens</i>	Dill	Antifungal activity ( <i>Candida</i> spp.)	MIC and MFC were 0.63–2.5 mg/mL and 1.25–5 mg/mL, respectively.	[21]
		Antifungal activity ( <i>Penicillium citrinum</i> , <i>Penicillium veridicatum</i> , <i>Aspergillus flavus</i> , <i>Aspergillus niger</i> , <i>Fusarium graminearum</i> ), antibacterial activity ( <i>Staphylococcus aureus</i> , <i>Bacillus subtilis</i> , <i>Pseudomonas aeruginosa</i> , <i>Salmonella typhi</i> , <i>Escherichia coli</i> , <i>Bacillus cereus</i> )	Antifungal activity by inverted petri dish method 27.5–100% at 6 µL. Inhibition zones determined by agar well diffusion method 13.2–25.3 mm (bacteria).	[22]
<i>Angelica archangelica</i>	Garden Angelica	Antibacterial activity against <i>Staphylococcus aureus</i> and <i>Escherichia coli</i>	MIC using broth microdilution method = 14.2 µL/mL ( <i>S. aureus</i> ) and 28.4 µL/mL ( <i>E. coli</i> ).	[23]
<i>Angelica sylvestris</i>	Wild Angelica	Larvicidal activity ( <i>Culex pipiens</i> )	LC <sub>50</sub> > 150 mg/L.	[80]
<i>Anthriscus cerefolium</i>	Chervil	Insecticidal activity (wheat granary weevil <i>Sitophilus granaries</i> )	Repellent (62%) and lethal activity (54%).	[81]
		Antibacterial activity ( <i>Staphylococcus aureus</i> , <i>Salmonella typhi</i> )	MIC 0.12 µg/mL ( <i>S. aureus</i> ) and 0.5 µg/mL ( <i>S. typhi</i> ).	[82]
<i>Apium graveolens</i>	Celery	Insecticidal activity against <i>Pseudaletia unipuncta</i> (Armyworm)	Antifeedant/feeding deterrence index (84%), growth inhibitory (91%), fumigant and contact toxicant action (100% after 24 h).	[83]
		Antibacterial activity against <i>Bacillus cereus</i> , <i>Staphylococcus aureus</i> , <i>Listeria monocytogenes</i> , <i>Enterococcus faecalis</i>	11.5–40 mm inhibition zone in agar diffusion method.	[24]
<i>Azilia eryngioides</i>		Insecticidal activity ( <i>Sitophilus granaries</i> , <i>Tribolium castaneum</i> )	Fumigant toxicity 24-LC <sub>50</sub> were 20.05 µL/L ( <i>S. granaries</i> ) and 46.48 µL/L ( <i>T. castaneum</i> ).	[84]

Table 2. Cont.

Botanical Name	Common Name	Biocidal Effect	Results/Mechanisms	References
<i>Azorella cryptantha</i>		Repellent activity on <i>Triatoma infestans</i> , insecticidal activity against <i>Ceratitis capitata</i> , antifungal (dermatophytes <i>Microsporium gypseum</i> , <i>Trichophyton rubrum</i> , <i>Trichophyton mentagrophytes</i> ), antibacterial ( <i>Escherichia coli</i> and <i>Yersinia enterocolitica</i> )	Repellent activity (92–100%), insecticidal contact toxicity (LD <sub>50</sub> at 72 h <11 mg/fly), MIC (microbroth dilution method) of dermatophytes and bacteria 125–1000 µg/mL.	[27]
		Antibacterial activity ( <i>Escherichia coli</i> , <i>Salmonella enteritidis</i> )	MIC (microbroth dilution method) values of <i>S. enteritidis</i> and <i>E. coli</i> 125–250 µg/mL and 500 µg/mL, respectively.	[85]
<i>Carum carvi</i>	Caraway	Antifungal activity ( <i>Aspergillus flavus</i> ) and inhibition of aflatoxin production	Complete inhibition of <i>A. flavus</i> growth and aflatoxin B <sub>1</sub> production at 1000 ppm caraway EO.	[28]
		Insecticidal activity (African cotton leafworm <i>Spodoptera littoralis</i> )	Fumigant toxicity LD <sub>50</sub> 41.45 µL/L air.	[86]
		Insecticidal activity against <i>Sitophilus zeamais</i> and <i>Tribolium castaneum</i>	Contact (LD <sub>50</sub> 3.07 and 3.39 µg/adult) and fumigant (LC <sub>50</sub> 3.37 and 2.53 mg/L air) toxicity against <i>S. zeamais</i> and <i>T. castaneum</i> , respectively, and may be attributed to strong contact and fumigant toxicity of (R)-carvone and D-limonene.	[33]
		Herbicidal activity against barnyard grass <i>Echinochloa crus-galli</i>	Causing leaf injuries and reduction in biomass, impaired photosynthetic activity and plant metabolism.	[29]
		Antibacterial ( <i>S. epidermidi</i> , <i>S. aureus</i> , <i>M. luteus</i> , <i>E. faecalis</i> , <i>B. cereus</i> , <i>E. coli</i> , <i>L. monocytogenes</i> , <i>S. typhimurium</i> ) Antifungal ( <i>C. albicans</i> , <i>C. glabrata</i> , <i>C. parapsilosis</i> , <i>C. krusei</i> , <i>S. cerevisiae</i> )	Inhibition zone 11.00 to 25.00 mm diameter in disc-diffusion assay. MIC = 0.059 to 1.875 mg/mL (bacteria), 0.029–0.059 mg/mL (fungi). AChE inhibition (IC <sub>50</sub> = 0.82 mg/mL) α-Glucosidase inhibition (IC <sub>50</sub> = 6.83 mg/mL).	[30]
<i>Carum copticum</i>	Ajowan	Antifungal activities against <i>Penicillium digitatum</i> and <i>Alternaria alternata</i>	Mycelial growth inhibition 67.78% ( <i>A. alternata</i> ) and 44.88% ( <i>P. digitatum</i> ). Thymol may cause severe damage to cell walls, cell membranes and cellular organelles, such as mitochondria in fungi.	[31]

Table 2. Cont.

Botanical Name	Common Name	Biocidal Effect	Results/Mechanisms	References
<i>Carum nigrum</i>		Antifungal activity ( <i>Penicillium citrinum</i> , <i>Penicillium purpurogenum</i> , <i>Aspergillus flavus</i> , <i>Aspergillus terreus</i> , <i>Aspergillus niger</i> , <i>Fusarium graminearum</i> ), antibacterial activity ( <i>Staphylococcus aureus</i> , <i>Bacillus subtilis</i> , <i>Pseudomonas aeruginosa</i> , <i>Salmonella typhimurium</i> , <i>Escherchia coli</i> , <i>Bacillus cereus</i> )	Antifungal activity by inverted petri dish method 34–100% at 3000 ppm. Inhibition zones determined by agar well diffusion method 17.9 mm to complete inhibition at 3000 ppm (bacteria).	[35]
<i>Centella asiatica</i>	Gotu Kola, Centella, Indian penny	Larvicidal activity ( <i>Culex quinquefasciatus</i> )	Larvicidal activity (24 h-LC <sub>50</sub> was 1.12–6.84 ppm at the temperatures of 31–19 °C) and acts as adult emergence inhibitor.	[87]
		Antibacterial activity ( <i>Bacillus subtilis</i> , <i>Staphylococcus aureus</i> , <i>Escherichia coli</i> , <i>Pseudomonas aeruginosa</i> , <i>Shigella sonnei</i> )	MIC using microplate dilution method ranging from 1.25 to 0.039 mg/mL.	[36]
		Antifungal ( <i>Aspergillus niger</i> ) and Antibacterial ( <i>Bacillus subtilis</i> ) activity	6.3–15.4 mm ( <i>A. niger</i> ) and 8.4–16.4 mm ( <i>B. subtilis</i> ) inhibition zone in disc diffusion method.	[88]
<i>Chaerophyllum aromaticum</i>		Antibacterial activity	Bactericidal activity of Root EO against <i>Bacillus spizizenii</i> and <i>Staphylococcus aureus</i> (diameter of zone of inhibition 17 mm and 35 mm). BChE inhibitory activity of EO from root (47.65%) and aerial (50.88%).	[37]
<i>Chaerophyllum aureum</i>		Antibacterial activity ( <i>Staphylococcus aureus</i> , <i>Staphylococcus epidermidis</i> , <i>Micrococcus luteus</i> , and <i>Escherichia coli</i> )	10.0–22.3 mm inhibition zone in agar diffusion method. Mechanisms of inhibition may be attributed to cell membrane damage due to hydrophobicity and impairment of bacterial enzyme.	[38]
<i>Chaerophyllum crinitum</i>		Antifungal ( <i>Candida tropicalis</i> ) and antibacterial ( <i>Acinetobacter baumannii</i> , <i>Staphylococcus aureus</i> ) activities	Inhibition zone 16–10 mm diameter in disc diffusion method.	[39]
<i>Chaerophyllum macropodum</i>		Antibacterial activity ( <i>Escherichia coli</i> , <i>Salmonella typhi</i> )	Inhibition zone 6–12 mm diameter in disc diffusion method.	[89]

Table 2. Cont.

Botanical Name	Common Name	Biocidal Effect	Results/Mechanisms	References
		Antibacterial activity against <i>Staphylococcus epidermidis</i> , <i>Bacillus subtilis</i> , <i>Staphylococcus aureus</i> , <i>Escherichia coli</i>	Inhibition zone 10.7–18.1 mm diameter in agar disc diffusion method.	[40]
		Antifungal activity ( <i>Aspergillus flavus</i> ) and inhibition of aflatoxin production	Complete inhibition of <i>A. flavus</i> growth and aflatoxin B <sub>1</sub> production at 1000 ppm coriander EO.	[28]
		Acaricidal and insecticidal activities ( <i>Plodia interpunctella</i> , <i>Sitotroga cerealella</i> and <i>Tyrophagus putrescentiae</i> )	Fumigant toxicity (LD <sub>50</sub> 4.19–18.76 µg/cm <sup>3</sup> ), contact toxicity (LD <sub>50</sub> 19.29 µg/cm <sup>2</sup> for <i>T. putrescentiae</i> ).	[41]
		Insecticidal activity ( <i>Callosobruchus maculatus</i> , <i>Tribolium confusum</i> )	Insect mortality LC <sub>50</sub> of <i>C. maculatus</i> and <i>T. confusum</i> were 1.34 and 318.02 µL/L air, respectively.	[42]
<i>Coriandrum sativum</i>	Coriander	Antibacterial activity ( <i>Staphylococcus aureus</i> , <i>Bacillus spp.</i> , <i>Escherichia coli</i> , <i>Salmonella typhi</i> , <i>Klebsiella pneumonia</i> , <i>Proteus mirabilis</i> )	MIC 108–217 mg/mL.	[90]
		Antibacterial ( <i>S. epidermidis</i> , <i>S. aureus</i> , <i>M. luteus</i> , <i>E. faecalis</i> , <i>B. cereus</i> , <i>E. coli</i> , <i>L. monocytogenes</i> , <i>S. typhimurium</i> ) Antifungal ( <i>C. albicans</i> , <i>C. glabrata</i> , <i>C. parapsilosis</i> , <i>C. krusei</i> , <i>S. cerevisiae</i> )	Inhibition zone 8.33 to 21.66 mm diameter in disc diffusion assay. MIC = 0.234 to 1.875 mg/mL (bacteria), 0.234–0.469 mg/mL (fungi). AChE inhibition (IC <sub>50</sub> = 0.68 mg/mL) α-Glucosidase inhibition (IC <sub>50</sub> = 6.24 mg/mL).	[30]
		Antiparasitic activity ( <i>Trypanosoma cruzi</i> )	EC <sub>50</sub> = 17.7 µg/mL.	[91]
<i>Crithmum maritimum</i>	Sea Fennel	Antifungal properties against <i>Candida albicans</i> , <i>Cryptococcus neoformans</i> and several dermatophytes and <i>Aspergillus spp.</i>	<i>C. neoformans</i> 260 µg/mL (MIC), <i>C. albicans</i> (Inhibition of biofilm formation and germ tube formation).	[44]
		Larvicidal ( <i>Spodoptera littoralis</i> ; tobacco cut worm), AChE inhibitory activities	Toxicity 62.3–71.7 µg/larva (LD <sub>50</sub> ). AChE inhibitory activity 7.4 mg/mL (IC <sub>50</sub> ).	[47]
		Insecticidal activity ( <i>Sitophilus zeamais</i> )	Contact (LC <sub>50</sub> 0.246 µL/cm <sup>2</sup> area) and fumigant (LC <sub>50</sub> 0.346 µL/cm <sup>3</sup> air) toxicity, inhibition of AChE activity.	[92]
<i>Cuminum cyminum</i>	Cumin	Antifungal activity ( <i>Candida albicans</i> , <i>Lachancea thermotolerans</i> , <i>Metschnikowia pulcherrima</i> )	MIC ≤ 0.5 mg/mL.	[16]

Table 2. Cont.

Botanical Name	Common Name	Biocidal Effect	Results/Mechanisms	References
		Insecticidal activity against peach-potato aphid ( <i>Myzus persicae</i> ), tobacco cutworm/cotton leaf worm ( <i>Spodoptera littoralis</i> ), and two insect vectors, common housefly ( <i>Musca domestica</i> ) and lymphatic filariasis and Zika virus vector <i>Culex quinquefasciatus</i>	LD <sub>50</sub> = 3.2 mL/L ( <i>M. persicae</i> ), 100 µg/larva ( <i>S. littoralis</i> ), 31.8 µg/adult ( <i>M. domestica</i> ), 40.5 µL/L ( <i>C. quinquefasciatus</i> ).	[49].
		Insecticidal activity ( <i>Tribolium confusum</i> )	100% efficacy in dose of 1 mL.	[93]
<i>Daucus carota</i>	Wild Carrot	Antibacterial activity ( <i>Agrobacterium tumefaciens</i> , <i>Erwinia carotovora</i> , <i>Pseudomonas fluorescens</i> , <i>Pseudomonas glycinea</i> )	28.66–100% of antibacterial activities in disc diffusion method.	[13]
<i>Deverra scoparia</i>		Acaricidal activity against <i>Tetranychus urticae</i> (two-spotted spider mite)	Toxicity LD <sub>50</sub> 1.79 mg/L and decreased fecundity.	[51]
		Antibacterial ( <i>Escherichia coli</i> , <i>Pseudomonas aeruginosa</i> ) and antifungal activity ( <i>Trichoderma viride</i> )	MIC in microdilution technique were 0.0625 ( <i>E. coli</i> ), 0.25 ( <i>P. aeruginosa</i> ), and 0.0625 ( <i>T. viride</i> ) mg/mL.	[52]
<i>Echinophora spinosa</i>	Prickly Parsnip	Antibacterial ( <i>Bacillus subtilis</i> , <i>Escherichia coli</i> , <i>Proteus mirabilis</i> ) and antifungal activity ( <i>Candida albicans</i> , <i>Aspergillus niger</i> )	Inhibition zone 2–30.5 mm diameter in disc diffusion assay.	[94]
		Insecticidal activity ( <i>Culex quinquefasciatus</i> , <i>Spodoptera littoralis</i> , <i>Musca domestica</i> )	Toxicity LC <sub>50</sub> 15.7 mg/L ( <i>C. quinquefasciatus</i> ), LD <sub>50</sub> 38.3 µg/adult ( <i>M. domestica</i> ), 86.3 µg/larva ( <i>S. littoralis</i> ).	[95]
<i>Eryngium alpinum</i>	Alpine Sea Holly	Antibacterial ( <i>Escherichia coli</i> , <i>Staphylococcus aureus</i> ), antifungal ( <i>Candida albicans</i> , <i>Microsporium gypseum</i> )	MIC 0.08–1.94 mg/mL using microdilution assay.	[53]
<i>Eryngium amethystinum</i>	Amethyst sea holly	Antibacterial ( <i>Escherichia coli</i> , <i>Staphylococcus aureus</i> ), antifungal ( <i>Candida albicans</i> )	MIC 0.06–1.94 mg/mL using microdilution assay.	[53]
<i>Eryngium foetidum</i>	Culantro, Mexican coriander, long coriander	Antibacterial activity against <i>Bacillus subtilis</i> , <i>Staphylococcus aureus</i> , <i>Escherichia coli</i> , <i>Pseudomonas aeruginosa</i> , antifungal activity against <i>Candida albicans</i>	Inhibition zone 12–28 mm in agar well diffusion method MIC 1.56–200 µg/mL in microbroth dilution method.	[96]
		Larvicidal activity ( <i>Aedes albopictus</i> )	24 h-LC <sub>50</sub> s 33.3 ppm.	[97]
<i>Eryngium triquetrum</i>	Choukzerk	Antibacterial ( <i>Staphylococcus aureus</i> , <i>Staphylococcus epidermidis</i> , <i>Bacillus subtilis</i> ) and antifungal ( <i>Candida albicans</i> ) activities	MIC using broth dilution method = 6.25–25 µg/mL.	[54]

Table 2. Cont.

Botanical Name	Common Name	Biocidal Effect	Results/Mechanisms	References
		Herbicidal activity ( <i>Lepidium sativum</i> ), antibacterial activities against <i>Pectobacterium atrosepticum</i> (potato blackleg disease), and Gram-negative soil bacterium ( <i>Pseudomonas cichorii</i> ) Antifungal and fumigant activity against <i>Fusarium graminearum</i> (cereal fusarium) and <i>Botrytis cinerea</i> (grey rot in tomatoes, strawberries)	Inhibition of germination, growth, and photosynthesis (43%) of <i>L. sativum</i> , inhibition of 85% <i>Pectobacterium</i> , 100% <i>Pseudomonas</i> , fumigant toxicity with lack of growth on <i>F. graminearum</i> and moderate inhibition (43%) on <i>B. cinerea</i> .	[55]
<i>Falcaria vulgaris</i>		Antifungal ( <i>Aspergillus fumigatus</i> )	0.140 mg/mL (MIC using microdilution method). Cholinesterase inhibition (AChE and BChE).	[98]
<i>Ferula caspica</i>		Antibacterial activity ( <i>Staphylococcus aureus</i> , <i>Enterococcus faecalis</i> )	MIC in broth microdilution method 32–64 µg/mL.	[99]
<i>Ferula orientalis</i>		Antibacterial ( <i>Staphylococcus aureus</i> ) and antifungal ( <i>Candida albicans</i> ) activities	Determined by direct bioautography	[56]
<i>Ferula pseudalliacea</i>		Insecticidal activity ( <i>Varroa destructor</i> )	Highest (30.72%) mortality was at 36 h.	[100]
<i>Ferula rigidula</i>		Antifungal activity ( <i>Aspergillus ochraceus</i> , <i>Trichoderma viride</i> ), Antibacterial ( <i>Pseudomonas aeruginosa</i> )	MIC determined by microdilution method were 0.27 (bacteria) and 0.10 (fungi) mg/mL. Enzymatic inhibitory activities against AChE, BChE, tyrosinase, α-amylase, and α-glucosidase.	[101]
<i>Ferulago angulata</i>		Antibacterial activity against <i>Bacillus thuringiensis</i> , <i>Erwinia amylovora</i> , <i>Xanthomonas oryzae</i> Antifungal activity against <i>Colletotrichum trichellum</i> , <i>Fusarium oxysporum</i>	MIC = 8 ( <i>B. thuringiensis</i> ), 12.5 ( <i>E. amylovora</i> ), and 12 ( <i>X. oryzae</i> ) µL/mL. 47.8–100% fungal growth inhibition in agar dilution method 52.5–100% in disc diffusion method at concentration of 800 µL/L.	[57]
<i>Ferulago cassia</i>		Anticholinesterase activity	Cholinesterase inhibition (AChE and BChE).	[58]
<i>Ferulago sandrasica</i>		Antibacterial ( <i>Staphylococcus aureus</i> ) and antifungal ( <i>Candida albicans</i> ) activity	Determined by direct bioautography.	[56]



Table 2. Cont.

Botanical Name	Common Name	Biocidal Effect	Results/Mechanisms	References
<i>Ferulago trachycarpa</i>		Antibacterial activity against <i>Staphylococcus aureus</i> and <i>E. faecalis</i> (dichloromethane and n-hexane extract), Antifungal activity against <i>Candida albicans</i> , <i>Candida tropicalis</i> , <i>Candida parapsilosis</i> (all extracts)	MIC in microdilution method 3.9–625 mg/L.	[59]
<i>Ferulago trifida</i>		Larvicidal activity ( <i>Anopheles stephensi</i> ) cytotoxic activity and inhibition of AChE activity	LC <sub>50</sub> = 116.54 ppm.	[60,102]
<i>Foeniculum vulgare</i>	Fennel	Insecticidal activity ( <i>Sitophilus granaries</i> , <i>Sitophilus oryzae</i> )	Fumigation toxicity 27.3 µL/L air ( <i>S. granaries</i> ) and 44.16 µL/L air ( <i>S. oryzae</i> ) (LC <sub>50</sub> ).	[103]
		Insecticidal activity (African cotton leafworm <i>Spodoptera littoralis</i> )	Fumigant toxicity LD <sub>50</sub> 51.2 µL/L air.	[86]
		Larvicidal, pupicidal, and insecticidal activity ( <i>Culex quinquefasciatus</i> )	Larvicidal activity with LC <sub>50</sub> = 0.10 mg/mL, LD <sub>50</sub> = 7.52/min.	[104]
		Larvicidal and ovicidal activities ( <i>Culex pipiens</i> )	Larvicidal activity with 24 h-LC <sub>50</sub> = 148.3 µg/mL.	[105]
		Insecticidal activity ( <i>Lymantria dispar</i> )	Antifeedant activity at concentration of 1% EO (77.68 Tot).	[79]
		Insecticidal activity ( <i>Sitophilus granaries</i> , <i>Tribolium castaneum</i> )	100% ( <i>T. castaneum</i> ) and 43.58% ( <i>S. granaries</i> ) fumigant toxicity.	[62]
		Antifungal activities against <i>Penicillium digitatum</i> and <i>Alternaria alternata</i>	Mycelial growth inhibition 13.94% ( <i>A. alternata</i> ) and 55.39% ( <i>P. digitatum</i> ).	[31]
		Insecticidal activity against <i>Tribolium castaneum</i> and <i>Sitophilus oryzae</i>	Fumigant toxicity [91.28 and 254.71 µL/L air, respectively (LC <sub>50</sub> )] and repellency effects.	[43]
		Herbicidal activity against weeds of wheats ( <i>Phalaris minor</i> , <i>Avena ludoviciana</i> , broad-leaved weeds, <i>Rumex dentatus</i> , and <i>Medicago denticulate</i> )	Inhibition of germination and seedling.	[63]
<i>Heracleum anisactis</i>		Antibacterial activity ( <i>Escherichia coli</i> , <i>Staphylococcus epidermidis</i> , <i>Staphylococcus aureus</i> )	11–13 mm inhibition zone in agar disc diffusion method, MIC in broth dilution method 1.1–1.5 v/v.	[106]
<i>Heracleum persicum</i>	Persian Hogweed	Larvicidal activity ( <i>Anopheles stephensi</i> )	24 h-LC <sub>50</sub> = 26–336 ppm	[107]

Table 2. Cont.

Botanical Name	Common Name	Biocidal Effect	Results/Mechanisms	References
		Insecticidal ( <i>Callosobruchus maculatus</i> )	Fumigant toxicity 24 h-LC <sub>50</sub> = 136.4 µL/L air and higher oviposition deterrence property.	[108]
<i>Hippomarathrum microcarpum</i>		Antibacterial ( <i>Staphylococcus aureus</i> ) and antifungal ( <i>Candida albicans</i> ) activity	Determined by direct bioautography.	[56]
<i>Levisticum officinale</i>	Lovage	Insecticidal activity ( <i>Tribolium confusum</i> )	100% efficacy in dose of 2 mL.	[93]
<i>Myrrhis odorata</i>	Sweet Cicely	Antifungal activity ( <i>Cladosporium cladosporioides</i> , <i>Aspergillus niger</i> , <i>Aspergillus flavus</i> , <i>Trichoderma viride</i> )	MIC for fungi were 0.5–1.0 mg/mL in microdilution test.	[109]
<i>Ostericum sieboldii</i>	Water Dropwort	Insecticidal activity against the red flour beetle ( <i>Tribolium castaneum</i> ) and maize weevil ( <i>Sitophilus zeamais</i> )	Contact toxicity against <i>T. castaneum</i> and <i>S. zeamais</i> with LD <sub>50</sub> values of 8.47 and 13.82 µg/adult, respectively. Fumigant toxicity LC <sub>50</sub> values of 20.92 and 27.39 mg/L air, respectively.	[66]
<i>Pastinaca sativa</i>	Parsnip	Antibacterial activity ( <i>Bacillus cereus</i> )	MIC in microwell dilution assay 0.72 mg/mL.	[110]
<i>Petroselinum crispum</i>	Parsley	Insecticidal activity against <i>Pseudaletia unipuncta</i> (Armyworm)	Antifeedant/feeding deterrence index (99.7%), and contact toxicant action (93% after 48 h).	[83]
<i>Peucedanum ostruthium</i>	Masterwort	Phytotoxic activity ( <i>Lolium multiflorum</i> , <i>Echinochloa oryzoides</i> ), nematocidal ( <i>Panagrolaimus rigidus</i> )	Decreased germination and significant impact on <i>L. multiflorum</i> root and shoot growth. Nematicidal activity of leaves extract: 85.6% and 90.5% mortality of larvae and adults of <i>P. rigidus</i> .	[111]
		Insecticidal activity ( <i>Lymantria dispar</i> )	Antifeedant activity at concentration of 1% EO (Tot 119.26).	[79]
<i>Pimpinella anisum</i>	Anise	Antibacterial ( <i>Bacillus subtilis</i> , <i>Klebsiella pneumonia</i> ) and antifungal ( <i>Aspergillus niger</i> , <i>Candida albicans</i> ) activities	11–40 mm inhibition in cup plate–agar diffusion method.	[112]
		Insecticidal activity against <i>Tribolium castaneum</i> and <i>Sitophilus oryzae</i>	Fumigant toxicity [43.75 and 292.04 µL/L air, respectively (LC <sub>50</sub> ), and repellency effects.	[43]

Table 2. Cont.

Botanical Name	Common Name	Biocidal Effect	Results/Mechanisms	References
		Insecticidal activity against peach-potato aphid ( <i>Myzus persicae</i> ), tobacco cutworm/cotton leaf worm ( <i>Spodoptera littoralis</i> ), and two insect vectors, common housefly ( <i>Musca domestica</i> ) and lymphatic filariasis and Zika virus vector <i>Culex quinquefasciatus</i>	<i>M. persicae</i> (4.3 mL/L), <i>S. littoralis</i> (57.3 µg/larva), <i>M. domestica</i> (54.8 µg/adult), <i>C. quinquefasciatus</i> (25.4 µL/L) (LD <sub>50</sub> ).	[49].
		Antifungal ( <i>Trichophyton rubrum</i> ) and antibacterial activities ( <i>Streptococcus mutans</i> , <i>Streptococcus pyogenes</i> , and <i>Staphylococcus aureus</i> )	MIC determined by twofold serial broth dilution method were 2000 ( <i>T. rubrum</i> ) and ≤1.19 (bacteria) µg/mL.	[67]
<i>Prangos peucedanifolia</i>		Antibacterial activity ( <i>Pseudomonas aeruginosa</i> , <i>Escherichia coli</i> , <i>Bacillus cereus</i> , <i>Listeria monocytogenes</i> , <i>Salmonella typhimurium</i> ), Antifungal activity ( <i>Aspergillus ochraceus</i> , <i>Trichoderma viride</i> , <i>Aspergillus versicolor</i> ), inhibition of α-glucosidase	MIC determined by microdilution method were 0.27–0.56 (bacteria) and 0.22–0.27 (fungi) mg/mL. Enzymatic inhibitory activities against AChE, tyrosinase, α-amylase, and α-glucosidase.	[101]
<i>Seseli gummiferum</i>	Moon Carrot	Antibacterial action against <i>Staphylococcus lugdunensis</i>	Enzymatic inhibitory activities against AChE, BChE, tyrosinase, and α-amylase MIC = 0.025–0.5 mg/mL.	[68]
<i>Sison amomum</i>	Stone Parsley	Antibacterial ( <i>Staphylococcus aureus</i> , <i>Klebsiella pneumonia</i> , <i>Acinetobacter baumannii</i> )	MIC using microdilution method = 2.0–4.1 mg/mL.	[69]
<i>Smyrniolum olusatrum</i>	Alexanders, Horse Parsley	Herbicidal activity ( <i>Lepidium sativum</i> ) Antifungal activity against <i>Fusarium graminearum</i> and <i>Zymoseptoria tritici</i> Fumigant toxicity ( <i>Fusarium graminearum</i> , <i>Botrytis cinerea</i> )	Inhibition of germination, growth, and photosynthesis (74%) of <i>L. sativum</i> , lack of growth on <i>F. graminearum</i> and 83% inhibition on <i>Z. tritici</i> mycelial growth at low concentration 0.142 µL/mL air. Fumigant toxicity with lack of growth on <i>F. graminearum</i> and 48% inhibition on <i>B. cinerea</i> at 0.142 µL/mL air.	[55]
<i>Trachyspermum ammi</i>	Ajwain, Ajowan caraway	Antibacterial activity ( <i>Bacillus cereus</i> , <i>Staphylococcus aureus</i> , <i>Listeria monocytogenes</i> , <i>Salmonella typhimurium</i> , <i>Escherichia coli</i> ), Antifungal activity ( <i>Penicillium citrinum</i> , <i>Penicillium chrysogenum</i> , <i>Aspergillus flavus</i> , <i>Aspergillus niger</i> , <i>Aspergillus parasiticus</i> )	Inhibition zone in disc diffusion method = 34–39 mm (bacteria) and >80 mm (fungi). MIC using broth microdilution method = 500 ppm (bacteria) and 1000–2000 ppm (fungi).	[70]

Table 2. Cont.

Botanical Name	Common Name	Biocidal Effect	Results/Mechanisms	References
		Insecticidal activity ( <i>Tribolium castaneum</i> )	Fumigant toxicity (LC <sub>50</sub> 11.6 µL), repellency, and reduces oviposition potential.	[78]
		Insecticidal activity against rice weevil ( <i>Sitophilus oryzae</i> ),	Repellent activity (95% at 1.2% EO), fumigant toxicity [0.37 µL/cm <sup>3</sup> (LC <sub>50</sub> )], AChE inhibition.	[15]
		Insecticidal activity ( <i>Plodia interpunctella</i> ) Antibacterial activity <i>S. aureus</i> , <i>E. coli</i> , <i>P. vulgaris</i> , <i>B. subtilis</i> , <i>K. pneumoniae</i> , and <i>B. megaterium</i>	Fumigant toxicity (LC <sub>50</sub> 4.33 µL/L air) MIC (dilution method) = 8–14 µL at 25% v/v EO.	[113]
		Insecticidal activity (maize weevil <i>Sitophilus zeamais</i> )	Repellent activity, fumigation (LC <sub>50</sub> 0.385 µL/cm <sup>3</sup> air), contact toxicity (0.317 µL/cm <sup>2</sup> ), and oviposition inhibitory effect.	[114]
		Insecticidal activity ( <i>Aethina tumida</i> )	Contact (LD <sub>50</sub> 66.64 µg/adult) and fumigant (LC <sub>50</sub> 89.03 mg/L air) toxicity may be attributed to the presence of thymol compound.	[115]

AChE, acetylcholinesterase; BChE, butyrylcholinesterase; LC<sub>50</sub>, median lethal concentrations; LD<sub>50</sub>, median lethal doses; MIC, minimum inhibitory concentration; EC<sub>50</sub>, half maximal effective concentration or concentration that reduces the infection in 50%; IC<sub>50</sub>, 50% inhibition of the AChE activity; Tot, total deterrence coefficients; MFC, minimum fungicidal concentration.

Botanical insecticides affect the pest with little or no side effects. These botanical insecticides kill the insects or affect their physiology in various ways, such as survival, behaviour, development, reproduction, and metabolic pathways [7].

The mechanism of insecticidal activity in secondary metabolites, including EOs, occurs via fumigant, contact, repellent, antifeedant, ovicidal, oviposition deterrent and larvicidal activity, and by inhibiting/altering neurotransmitters [acetylcholinesterase enzymes (AChE) and octopamine] and the neurotransmitter inhibitor ( $\gamma$ -aminobutyric acid (GABA)], impairment of the antioxidative defence systems, and cellular respiration [116–119]. Several earlier studies reported the inhibition of AChE by plant EOs. EOs components such as  $\alpha$ -pinene and  $\beta$ -pinene,  $\beta$ -phellandrene, carvacrol, limonene, menthol, menthone, 1,8-cineole, cis-ocimene, niloticin, and (L)-fenchone inhibited the AChE of insect pests [7,117]. Thymol and eugenol disrupt the functions of GABA and octopamine [118].

Phytotoxic plant extracts or metabolites delay or inhibit seed germination, as well as retarding the growth of weeds. The phytotoxic or herbicidal effect of EOs is associated with various mechanisms, including inhibition of DNA synthesis and cell proliferation, decreasing mitochondrial respiration, inducing reactive oxygen species (ROS) production, inhibition of enzymes, photosynthesis, seed germination, and seedling growth, altering membrane permeability and respiration. Oxygenated monoterpenes play a major role in most of the mechanisms. Monoterpenes, 1,8-cineole and camphor inhibit DNA synthesis, cell proliferation, and elongation [120–122].

The common antimicrobial mechanism of many EOs is attributed to the effect of membrane permeability or functioning, leading to cell death in fungi or bacteria. Other

than these, antifungal mechanisms of secondary metabolites, including EOs, involve inhibiting the fungi cell wall formation, inhibiting the mitochondrial electron transport, cell division, RNA or DNA and/or protein synthesis, and efflux pumps [123]. The sites and number of hydroxyl groups on the phenolic compounds (simple phenols, phenolic acids, quinones, flavones, flavonoids, flavonols, tannins, and coumarins) may be related to their relative toxicity to fungi. The antifungal mechanism of terpenoids (diterpenes, triterpenes, tetraterpenes, hemiterpenes, and sesquiterpenes) is related to membrane disruption by the lipophilic compounds, while the mechanisms of alkaloids (heterocyclic nitrogen compounds) are attributed to their ability to intercalate with DNA [12].

#### 4. Potential Use of Apiaceae Extracts as Agrochemicals

Synthetic agrochemicals are used to manage weeds, pests, and phytopathogens, thus increasing productivity and food safety. However, long-time usage of these synthetic chemicals may lead to environmental contamination, accumulation of toxic residues and biomagnification through the food chain, development of insect resistance, and threats to humans and non-target organisms [9,103]. These synthetic chemicals produce acute or chronic toxicity in living beings, depending on physicochemical characteristics, concentration and exposure time, route of entry, toxicodynamics and toxicokinetics (absorption, distribution, half-life, metabolism, and elimination), a combination of various pesticides, and the components of their formulation [9]. Secondary metabolites, including EOs, from aromatic plants, drew much attention due to their low toxicity to animals and non-target insects, fast degradation in the environment, and local availability. Other than the many health benefits, the biocidal potential of Apiaceae has great importance in agriculture for use as a natural pesticide, herbicide or phytotoxin, and antimicrobial [124].

##### 4.1. Insecticidal Activity for Crops and Stored Products Protection

###### 4.1.1. Insecticidal Activity to Protect Crops

Several Apiaceae genera are toxic to various insects [124]. *Tuta absoluta*, a major pest of tomato cropping, can be controlled by the EO of *Carum capticum* (Ajwain) [7]. Thymol,  $\gamma$ -terpinene, and  $\rho$ -cymene are the major components of *C. capticum* oil that showed AChE inhibition [7].

*Ammi visnaga* (toothpick-plant) seeds possess acaricidal activity against *Tetranychus urticae* Koch, which is a polyphagous species causing damage to numerous plants (around 1200 species), including major food crops and ornamental plants [125].

Cotton leafworm, *Spodoptera littoralis*, causes serious damage to important economic crops, such as maize, cotton, cereals, potatoes, tobacco, tomato, vegetables, and ornamental plants [47,86]. Apiaceae species, *C. carvi*, *F. vulgare*, *C. cyminum*, *C. caraway*, *C. sativum*, *D. carota* [86], *Helosciadium nodiflorum* (water celery) [126], and *Crithmum maritimum* [47] were demonstrated to possess toxicity against *S. littoralis* larvae. Benelli et al. [49] also suggest that cumin (*C. cyminum*) and anise (*P. anisum*) EOs can be used to protect crops from pests or to control insect vectors, as they have potential insecticidal activity against the peach-potato aphid (*Myzus persicae*) and the tobacco cutworm/cotton leafworm (*S. littoralis*), and insect vectors, such as the common housefly (*Musca domestica*). The crop pest *Spodoptera litura* is a threat to many crops and develops resistance to synthetic pesticides. This pest can be counteracted by sea fennel (*C. maritimum*) EOs [48].

###### 4.1.2. Insecticidal Activity to Protect Stored Products

Several species of storage insect pests infest agricultural products and stored granaries, causing substantial weight losses (around 10%) and quality [15]. They may carry pathogens and contaminate food [127]. *Sitophilus oryzae* (rice weevil), *Sitophilus zeamais* (maize weevil), *Tyrophagus putrescentiae* (mould or cheese mite), *Tribolium confusum* (confused flour beetle), *Callosobruchus maculatus* (cowpea weevil), *Periplaneta americana* (American cockroach), *Tribolium castaneum* (red flour beetle) are some of the pests that affect stored cereals, grains, flour, and other dried products [127]. Research is focused on producing safe and low-risk

natural compounds for pest control [43]. Several Apiaceae species were reported to carry insecticidal activities and have potential to protect stored grains from insect infestation.

Monoterpenes, such as (S)-(–)-linalool and (S)-(–)-menthone showed the most contact toxicity and fumigant activities, respectively, while  $\alpha$ -pinene, menthone, citronellal, and linalool exhibited repellent activities against the rice weevil (*S. oryzae*) [128]. Cuminaldehyde and (S)-carvone showed strong repellent activities, fumigation toxicity, and AChE inhibition [129].

Apiaceae species were demonstrated to possess insecticidal and repellent activity. EOs from *F. vulgare*, *Petroselinum hortense*, *C. sativum*, and *P. anisum* showed stronger fumigant toxicity against the storage pest, red flour beetle (*T. castaneum*) than against the rice weevil (*S. oryzae*) [43]. *C. carvi* EO and two isolated main compounds, (R)-carvone and D-limonene, exhibited strong contact and fumigant toxicity against the maize weevil (*S. zeamais*) and the red flour beetle (*T. castaneum*) [33]. *A. graveolens* (dill), *C. cyminum* (cumin), and *P. crispum* (parsley) EOs were reported as having contact toxicity, fumigant, and repellent activities against the stored grain pest *S. zeamais*. Moreover, AChE inhibition was also reported with *P. crispum* and *C. cyminum* EOs [129]. *Trachyspermum ammi* EO also showed repellent and fumigant activity against the rice weevil (*S. oryzae*). In addition, *T. ammi* EO showed significant inhibition of AChE activity in rice weevils [15].

Insecticidal activity of EO derived from *A. graveolens* (dill) seeds was reported against *T. confusum* (confused flour beetle), *C. maculatus* (cowpea weevils), *M. domestica*, *P. americana* (American cockroach), and *T. castaneum* (red flour beetles)] [130].

*Plodia interpunctella* (Indian meal moth), *Sitotroga cerealella* (grain moth), and *T. putrescentiae* (mould or cheese mite) affecting stored products, such as grains, flours, feeds, dried nuts, and fruits, globally. *P. interpunctella* moth continuously produces a silken web on the food; *S. cerealella* larvae invade grains and complete the larval and pupal stages within the grains, thus decreasing grain weight and nutritional value; and *T. putrescentiae* mites disseminate toxic fungi and induce allergic reactions among workers engaged in agriculture and food industries. Camphor and linalool found in *C. sativum* EO, extracted by steam distillation, exhibited acaricidal and insecticidal properties against *P. interpunctella*, *S. cerealella*, and *T. putrescentiae* [41].

#### 4.2. Herbicidal/Phytotoxic Activity against Weeds

Many studies were undertaken to investigate the herbicidal potential of Apiaceae species against weeds. Enzymes such as  $\alpha$ -amylase, catalase, and peroxidase enzymes play a key role in the physiological functions of the seeds and plants. Application of EOs increases oxidative stress in plant cells, which inhibits these enzymes and causes subsequent cell death. Moreover, the detrimental effect on the DNA and RNA of the cell results in decreased cell division, growth, and elongation [63].

Herbicidal effects of EOs derived from *C. sativum*, *C. carvi*, *F. vulgare*, and *P. anisum* were reported against various weeds [122]. *Eryngium triquetrum* EO and its major constituent falcarinol showed strong herbicidal activity against *Lepidium*, while *Smyrniolum olusatrum* EO showed moderate herbicidal activity. Therefore, *E. triquetrum* and *S. olusatrum* EOs can be used in crop protection to inhibit photosynthesis in weeds [55].

Caraway (*C. carvi*) EO (oil in water emulsion) are rich in oxygenated monoterpenes and exhibited herbicidal activity against *Echinochloa crus-galli* (barnyard grass, a typical maize weed) [29]. *C. carvi* EO was found to inhibit the germination of seeds of common weeds, including *Amaranthus retroflexus*, *Centaurea salsotitialis*, *Raphanus raphanistrum*, *Rumex nepalensis*, *Sonchus oleraceus*, and *Sinapis arvensis* [131].

Fennel (*F. vulgare*) EO strongly inhibited the seed germination and seedling growth of grass weeds, *Phalaris minor*, *Avena ludoviciana*, broad-leaved weeds, *Rumex dentatus* and *Medicago denticulate*, and can, therefore, be used in the biological management of weeds in wheat (*Triticum aestivum*) crops [63]. Fennel (*F. vulgare*) EO was also reported for its phytotoxic activity against field bindweed (*Convolvulus arvensis*) with significant inhibition of germination and early growth in field bindweed seedlings [132].

#### 4.3. Antimicrobial Activity against Phytopathogens

Plant pathogens, including fungi, bacteria, viruses, oomycetes, and nematodes, can cause diseases or damage to plants and result in significant crop yield and quality losses, of which fungi are the predominant pathogens, causing almost 30% of crop diseases during their cultivation or after harvest. Fungal disease symptoms on fruits, leaves, stems, and other plant parts include scabs, mouldy coatings, rust, smuts, powdery mildew, blotches, and rotted tissues. Plant diseases caused by phytopathogenic bacteria and fungi threaten human health and food security [13,120,133,134].

Various phytopathogenic bacteria, including *Xanthomonas* spp. (bacterial spots and blights), *Erwinia* spp. (soft rot, fire blight), *Pseudomonas* spp. (soft rot, bacterial canker), *Agrobacterium* spp. (crown gall), as well as *Ralstonia solanacearum* (bacterial wilt in banana), affect various agricultural plants [123].

Moreover, *Fusarium oxysporum* (vascular wilt of the banana tree), *Macrophomina phaseolina* (damping-off, seedling blight, and rot in peanuts, cabbage, pepper, chickpeas, soybeans, sweet potatoes, sesame, potatoes, sorghum, wheat, corn, etc.), *Alternaria alternata* (rot), *Penicillium digitatum* (green mould in citrus), and *Aspergillus flavus* (damping-off in peanut) are some of the fungi responsible for many diseases in various plants [134,135].

Apiaceae family plants were widely investigated for their potential biocidal activity against phytopathogens.

Apiaceae, such as *Torilis anthriscus*, *Aegopodium podagraria*, *D. carota*, *Heracleum sphondylium*, *Pimpinella saxifrage*, *P. sativa*, *Angelica silvestris*, and *F. vulgare* exhibited antimicrobial activity against phytopathogenic bacteria *Agrobacterium radiobacter* pv *tumefaciens*, *Erwinia carotovora*, *Pseudomonas fluorescens*, and *Pseudomonas glycinea* [13]. The EO of *C. maculatum* (poison hemlock), a poisonous Apiaceae species, has antibacterial activity against *Pseudomonas aeruginosa* [136].

*E. triquetrum* EO exhibited strong antibacterial activities against potato blackleg disease, *Pectobacterium atrosepticum*, and the soil bacterium *Pseudomonas cichorii*, which is responsible for disease in lettuce, celery, and chrysanthemum, while *S. olusatrum* EO showed moderate antibacterial activity [55].

The Apiaceae *Ferulago angulate* exhibited toxicity in variable degrees against phytopathogenic bacteria (*Erwinia amylovora*, *Xanthomonas oryzae*, *Pseudomonas syringae*, *Pectobacterium carotovorum*, *R. solanacearum*, *Bacillus thuringiensis*), and fungi (*A. alternata*, *Culvularia fallax*, *M. phaseolina*, *F. oxysporum*, *Cytospora sacchari*, *Colletotrichum tricbellum*) [57].

*A. alternata* and *P. digitatum* are two postharvest pathogens in tomato fruits. Based on in vivo studies, *Carum copticum* and *F. vulgare* EOs have the potential to control postharvest decay in tomatoes caused by *A. alternata* and *P. digitatum* [31].

*E. triquetrum* and *S. olusatrum* EOs have potent antifungal and fumigant activity against *Fusarium graminearum* (cereal fusarium) and moderate activity against *Botrytis cinerea* (grey rot on tomatoes, strawberries). *S. olusatrum* EO showed strong antifungal activity against *F. graminearum* and *Zymoseptoria tritici*, which are responsible for the septoria blight [55].

*A. graveolens* (dill) seeds EO mainly consists of carvone, limonene,  $\alpha$ -phellandrene,  $\beta$ -phellandrene and p-cymene and were proved to have antifungal (*Aspergillus niger*, *Aspergillus oryzae*, *A. flavus*, *A. alternata*) and antibacterial (*Escherichia coli*, *P. aeruginosa*, *Bacillus subtilis*, *Staphylococcus aureus*) activities [130].

Food poisoning is the most common cause of illness and death in both developed and developing countries. Most food poisoning diseases are caused by bacterial contamination. Pathogenic and food spoilage bacteria, such as *Salmonella typhi*, *E. coli*, *P. aeruginosa*, *S. aureus*, *Listeria monocytogenes*, and *Bacillus cereus* are the causal agents of foodborne diseases or food spoilage. Plant extracts are a good source of antimicrobial agents that can be used as food preservatives [137,138].

Apiaceae *C. cyminum* was reported to be effective against *S. aureus* [138]. Coriander (*C. sativum*) EO and its major compound linalool showed antibacterial activity against *Campylobacter jejuni* and *Campylobacter coli*, pathogens that cause foodborne diseases [139].

Coriander (*C. sativum*) was also reported to have strong antimicrobial activity against bacteria, *B. subtilis*, followed by *Stenotropomonas maltophilia*, and *Penicillium expansum* (fungi producing mycotoxin). Moreover, the strongest antibiofilm activity of coriander EO was also reported against *S. maltophilia* [140].

*Carum nigrum* (black caraway) seed EO has antibacterial activity against foodborne bacteria, *B. cereus* and *P. aeruginosa*, and antifungal activity against foodborne fungi, such as *P. purpurogenum*, *Acrophialophora fusispora*, *A. flavus*, and *A. niger* [35].

The antimicrobial activity of *T. ammi* (Ajwain) EO was demonstrated against various food spoilage and foodborne bacteria (*B. cereus*, *S. aureus*, *L. monocytogenes*, *Salmonella typhimurium*, *E. coli*) and fungi (*Penicillium citrinum*, *Penicillium chrysogenum*, *A. flavus*, *A. niger*, *Aspergillus parasiticus*) [70].

Moreover, certain fungi, including *Aspergillus* spp., *Fusarium* spp., and *Alternaria* spp., produce mycotoxins that can be harmful to human health due to hepatotoxic, nephrotoxic, and carcinogenic effects, or even cause death [120]. Mycotoxins are toxic secondary metabolites produced by certain fungal species in various agricultural and other food products, either in the field or during storage. Mycotoxin contamination in crops is potentially harmful to animals and human health. Aflatoxins B1, B2, G1, and G2 are the four major toxins generated in foods, of which the aflatoxin B1, secreted by *A. flavus*, *A. parasiticus* and *Aspergillus nomius*, is the most toxic and has potent teratogenic, mutagenic, hepatotoxic, and immune suppressive activities [141–143].

Apiaceae, *F. vulgare* EO extracted from flowers and roots, significantly inhibited the growth of *A. parasiticus* and the production of aflatoxins B1 and G1 [144]. Carvone and linalool compounds found in *C. carvi* and *C. sativum* seed EOs, respectively, showed an antifungal and aflatoxin-inhibition ability against *A. flavus*. Thus, they can be used as a preservative, particularly in post-harvest processing and the storage of agricultural products that are susceptible to aflatoxin contamination, such as cereals, dried fruits, and spices [143]. EOs of other Apiaceae plants, including *C. cyminum* [145], *C. carvi* [28,146], *C. sativum* [28], and *C. copticum* (*T. ammi*) were also reported to inhibit both growth and/or mycotoxin production. Table 3 summarizes the potential use of Apiaceae extracts as agrochemicals.

**Table 3.** Potential use of Apiaceae extracts as agrochemicals.

Insects/Weeds/Phytopathogens	Effects	Apiaceae	References
<b>Insecticidal activity to protect crops</b>			
<i>Myzus persicae</i> (Peach-potato aphid)	Agricultural pest	<i>C. cyminum</i> , <i>P. anisum</i>	[49]
<i>Spodoptera littoralis</i> (Tobacco cutworm/cotton leafworm)	Affects maize, cotton, cereals, potatoes, tobacco, tomato, vegetables, and ornamental plants.	<i>C. carvi</i> , <i>F. vulgare</i> , <i>C. cyminum</i> , <i>C. caraway</i> , <i>C. sativum</i> , <i>D. carota</i> , <i>P. anisum</i> , <i>C. maritimum</i> , <i>H. nodiflorum</i>	[47,49,86]
<i>Spodoptera litura</i> (Tobacco cutworm/cotton leafworm/armyworm)	Agricultural pest, affecting many crops.	<i>C. maritimum</i>	[48].
<i>Tetranychus urticae</i>	Affects numerous plants (around 1200 species), including major food crops and ornamental plants.	<i>A. visnaga</i>	[125]
<i>Tuta absoluta</i>	Tomato	<i>C. copticum</i>	[7]
<b>Insecticidal activity to protect stored products</b>			
<i>Culex quinquefasciatus</i>	Lymphatic filariasis and Zika virus vector	<i>P. anisum</i>	[49]



Table 3. Cont.

Insects/Weeds/Phytopathogens	Effects	Apiaceae	References
<i>Musca domestica</i> (Common housefly)	Storage pests, Insect vectors	<i>C. cyminum</i> , <i>P. anisum</i>	[49,130]
<i>Sitophilus oryzae</i> (Rice weevil)	Storage pests	<i>F. vulgare</i> , <i>P. hortense</i> , <i>C. sativum</i> , <i>P. anisum</i> , <i>T. ammi</i>	[15,43]
<i>Sitophilus zeamais</i> (Maize weevils)	Stored grain pest	<i>C. carvi</i> , <i>A. graveolens</i> , <i>C. cyminum</i> , <i>P. crispum</i>	[33,129]
<i>Tribolium castaneum</i> (Red flour beetles)	Storage pests	<i>F. vulgare</i> , <i>P. hortense</i> , <i>C. sativum</i> , <i>P. anisum</i> , <i>C. carvi</i> , <i>A. graveolens</i>	[33,43,130]
<i>Tyrophagus putrescentiae</i> (Mould or cheese mite)	Storage pests, disseminate toxic fungi and induce allergic reactions among workers engaged in agriculture and food industries.	<i>C. sativum</i>	[41]
<i>Callosobruchus maculatus</i> (Cowpea weevils) <i>Tribolium confusum</i> (Confused flour beetle) <i>Periplaneta americana</i> (American cockroach)	Storage pests	<i>A. graveolens</i>	[130]
<i>Plodia interpunctella</i> (Indian meal moth) <i>Sitotroga cerealella</i> (Grain moth)	Affect stored products, such as grains, flours, feeds, dried nuts, and fruits.	<i>C. sativum</i>	[41]
<b>Herbicidal/phytotoxic activity against weeds</b>			
<i>Convolvulus arvensis</i> (Field bindweed)	Common weed over the world	<i>F. vulgare</i>	[132].
<i>Echinochloa crus-galli</i> (Barnyard grass, a typical maize weed)	Maize weed	<i>C. carvi</i>	[29]
<i>Lepidium sativum</i> (watercress)		<i>E. triquetrum</i> , <i>S. olusatrum</i>	[55]
<i>Amaranthus retroflexus</i> , <i>Centaurea salsotitialis</i> , <i>Raphanus raphanistrum</i> , <i>Rumex nepalensis</i> , <i>Sonchus oleraceus</i> , and <i>Sinapis arvensis</i>	Common weeds	<i>C. carvi</i>	[131]
Grass weeds, <i>Phalaris minor</i> and <i>Avena ludoviciana</i> , broad-leaved weeds, <i>Rumex dentatus</i> and <i>Medicago denticulate</i>	Weeds of wheat ( <i>Triticum aestivum</i> )	<i>F. vulgare</i>	[63]
<b>Antimicrobial activity against phytopathogens</b>			
<i>Alternaria alternata</i>	Phytopathogens; postharvest pathogens in tomato fruits	<i>C. copticum</i> , <i>F. vulgare</i> , <i>A. graveolens</i> , <i>F. angulate</i>	[31,57,130]
<i>Penicillium digitatum</i>	Postharvest pathogens in tomato fruits	<i>C. copticum</i> , <i>F. vulgare</i>	[31]
<i>Pectobacterium atrosepticum</i>	Potato blackleg disease		
<i>Pseudomonas cichorii</i>	Soil bacterium causing disease of lettuce, celery, and chrysanthemum	<i>E. triquetrum</i> , <i>S. olusatrum</i>	[55]

Table 3. Cont.

Insects/Weeds/Phytopathogens	Effects	Apiaceae	References
Bacteria ( <i>Erwinia amylovora</i> , <i>Xanthomonas oryzae</i> , <i>Pseudomonas syringae</i> , <i>Pectobacterium carotovorum</i> , <i>Ralstonia solanacearum</i> , <i>Bacillus thuringiensis</i> ), and fungi ( <i>Culvularia fallax</i> , <i>Macrophomina phaseolina</i> , <i>Fusarium oxysporum</i> , <i>Cytospora sacchari</i> , <i>Colletotrichum trichellum</i> )	Phytopathogens	<i>F. angulate</i>	[57]
Bacteria; <i>Agrobacterium radiobacter</i> pv <i>tumefaciens</i> , <i>Erwinia carotowora</i> , <i>Pseudomonas fluorescens</i> , and <i>Pseudomonas glycinea</i>	Phytopathogenic bacteria	<i>T. anthriscus</i> , <i>A. podagraria</i> , <i>D. carota</i> , <i>H. sphondylium</i> , <i>P. saxifrage</i> , <i>P. sativa</i> , <i>A. silvestris</i> , <i>F. vulgare</i>	[13]
<i>Fusarium graminearum</i>	Cereal fusarium/wheat head blight fungus		
<i>Botrytis cinerea</i>	Grey rot on tomatoes, strawberries	<i>E. triquetrum</i> , <i>S. olusatrum</i>	[55]
<i>Zymoseptoria tritici</i>	Septoria blight		
<i>Aspergillus flavus</i>	Food spoilage and foodborne fungi, production of aflatoxins in cereals, dried fruits, and spices	<i>C. carvi</i> , <i>C. sativum</i> , <i>T. ammi</i> , <i>C. nigrum</i> , <i>A. graveolens</i>	[35,70,130,143]
<i>Aspergillus niger</i>	Food spoilage and foodborne fungus	<i>A. graveolens</i> , <i>C. nigrum</i> , <i>T. ammi</i>	[35,70,130]
<i>Aspergillus oryzae</i>	Foodborne pathogens	<i>A. graveolens</i>	[130]
<i>Aspergillus parasiticus</i>	Food spoilage and foodborne fungus, Production of aflatoxins B1 and G1	<i>F. vulgare</i> , <i>T. ammi</i>	[70,144]
<i>Bacillus cereus</i>	Food spoilage and foodborne pathogenic bacteria	<i>C. nigrum</i> , <i>T. ammi</i>	[35,70]
<i>Bacillus subtilis</i>	Foodborne pathogenic bacteria	<i>A. graveolens</i> , <i>C. sativum</i>	[130,140]
<i>Campylobacter jejuni</i> and <i>Campylobacter coli</i>	Foodborne pathogens	<i>C. sativum</i>	[139]
<i>Escherichia coli</i>	Food spoilage and foodborne bacteria	<i>T. ammi</i> , <i>A. graveolens</i>	[70,130]
<i>Pseudomonas aeruginosa</i>	Foodborne pathogens	<i>C. maculatum</i> , <i>A. graveolens</i> , <i>C. nigrum</i>	[35,130,136]
<i>Staphylococcus aureus</i>	Food spoilage and foodborne pathogenic bacteria	<i>C. cyminum</i> , <i>A. graveolens</i> , <i>T. ammi</i>	[70,130,138]
<i>Stenotropomonas maltophilia</i> (Bacteria), <i>Penicillium expansum</i> (fungi-producing mycotoxin)	Foodborne pathogens	<i>C. sativum</i>	[140]
<i>Penicillium purpurogenum</i> , <i>Acrophialophora fusispora</i>	Foodborne pathogenic fungi	<i>C. nigrum</i>	[35]
Bacteria ( <i>Listeria monocytogenes</i> , <i>Salmonella typhimurium</i> ), Fungi ( <i>Penicillium citrinum</i> , <i>Penicillium chrysogenum</i> )	Food spoilage and foodborne bacteria and fungi	<i>T. ammi</i>	[70]

## 5. Future Trends

Although EOs are an excellent alternative for synthetic chemicals, low water solubility, low persistence in the environment, chemical variability, low yield, and easy degradation under heat, humidity, light, and oxygen may limit the application of essential oils as biocides. To overcome these drawbacks, nanotechnologies are modern and eco-friendly approaches that enhance the effectiveness of natural products [48,63,118].

Nanoemulsion is a novel approach that increases the physical stability of active substances, decreases volatility, enhances solubility, and prevents interactions with the environment [48,63]. Nanoemulsions are oil-in-water dispersions with extremely small droplet diameters in the range of 10–100 nm and extended stability [48]. Size reduction to a nanoscale can improve the properties of materials, such as enhanced infusibility, effective release of the active ingredient, rapid interaction, enhanced bioactivity towards the target, and physical stability [63,147]. In agricultural applications, nanoemulsions have significant herbicidal and antimicrobial potential against weeds and pathogens that are resistant to synthetic biocides. This may be attributed to increased chemical stability and solubility, decreased evaporation, and the degradation of active essential oil components [63,148]. Nanotechnology also enhances efficacy and reduces the number of active ingredients currently employed [149].

The germination of *P. minor*, *A. ludoviciana*, *R. dentatus*, and *Medicago denticulate* was completely suppressed by nanoemulsions of *F. vulgare* EO at low concentrations. Further, the storage stability of *F. vulgare* nanoemulsion at ambient temperature was increased [63]. Significant antimicrobial activity of cumin (*C. cyminum*) and fennel (*F. vulgare*) EO nanoemulsions showed antibacterial/anti-biofilm properties against the foodborne pathogens *K. pneumoniae*, *C. violaceum*, and *S. typhimurium* [150]. Repellent activity by aniseed oil against *Rhopalosiphum padi* was also reported [149].

Other than nanotechnology, advanced extraction techniques, chemical synthesis of novel compounds, metabolic engineering, biotechnology, DNA sequencing, and recombinant DNA technologies can also enhance plant product effectiveness and commercial applications [118].

## 6. Conclusions

In conclusion, plant-based biocides can be a suitable candidate to substitute synthetic chemicals as they are cheap, safe, sustainable, eco-friendly, and biodegradable substances. In this context, secondary metabolites of Apiaceae species are widely used to control most agricultural pests, weeds, pathogenic foodborne, and food spoilage microbes to increase agriculture production and ensure food security. Several studies on various Apiaceae species for their potential biocidal activity proved their potent insecticidal, herbicidal, antibacterial, and antifungal activities. Therefore, Apiaceae-derived secondary metabolites, including EOs, can be used as a natural source in biocontrol products in agriculture, bio-pesticide, bio-herbicide, bio-fungicide, and antibacterial agents in organic farming and crop protection. However, there are several limitations in the application of secondary metabolites, including EOs, mainly due to their fast degradation and low stability in heat, light, or oxygen. Recently, nanotechnology research has attempted to overcome these limitations of natural products used as biocides.

**Author Contributions:** Writing—original draft, P.T., N.G., A.G., O.M. and T.M. Writing—review and editing, P.T., A.G., N.G., O.M. and T.M. All authors have read and agreed to the published version of the manuscript.

**Funding:** This research received no external funding

**Conflicts of Interest:** The authors declare no conflict of interest.

## References

1. Ahmad, B.S.; Talou, T.; Saad, Z.; Hijazi, A.; Ahmad, B.S.; Talou, T.; Saad, Z.; Hijazi, A.; Merah, O.; Apiaceae, T. The Apiaceae: Ethnomedicinal Family as Source for Industrial Uses. *Ind. Crops Prod.* **2017**, *109*, 661–671. [[CrossRef](#)]
2. Aćimović, M.G. Nutraceutical Potential of Apiaceae. In *Bioactive Molecules in Food*; Mérillon, J.-M., Ramawat, K.G., Eds.; Reference Series in Phytochemistry; Springer International Publishing: Cham, Switzerland, 2019; pp. 1311–1341. ISBN 978-3-319-78030-6.
3. Aćimović, M.G.; Milić, N.B. Perspectives of the Apiaceae Hepatoprotective Effects—A Review. *Nat. Prod. Commun.* **2017**, *12*, 309–317. [[CrossRef](#)] [[PubMed](#)]
4. Sousa, R.M.O.F.; Cunha, A.C.; Fernandes-Ferreira, M. The Potential of Apiaceae Species as Sources of Singular Phytochemicals and Plant-Based Pesticides. *Phytochemistry* **2021**, *187*, 112714. [[CrossRef](#)] [[PubMed](#)]
5. Christensen, L.P.; Brandt, K. Bioactive Polyacetylenes in Food Plants of the Apiaceae Family: Occurrence, Bioactivity and Analysis. *J. Pharm. Biomed. Anal.* **2006**, *41*, 683–693. [[CrossRef](#)]
6. Damalas, C.A.; Eleftherohorinos, I.G. Pesticide Exposure, Safety Issues, and Risk Assessment Indicators. *Int. J. Environ. Res. Public Health* **2011**, *8*, 1402–1419. [[CrossRef](#)]
7. Piri, A.; Sahebzadeh, N.; Zibae, A.; Sendi, J.J.; Shamakhi, L.; Shahriari, M. Toxicity and Physiological Effects of Ajwain (*Carum copticum*, Apiaceae) Essential Oil and Its Major Constituents against *Tuta absoluta* (Meyrick) (Lepidoptera: Gelechiidae). *Chemosphere* **2020**, *256*, 127103. [[CrossRef](#)]
8. Rajput, P.; Thakur, A.; Devi, P. Chapter 5—Emerging Agrochemicals Contaminants: Current Status, Challenges, and Technological Solutions. In *Agrochemicals Detection, Treatment and Remediation*; Prasad, M.N.V., Ed.; Butterworth-Heinemann: Oxford, UK, 2020; pp. 117–142. ISBN 978-0-08-103017-2.
9. Ramírez-Gómez, X.S.; Jiménez-García, S.N.; Campos, V.B.; Campos, M.L.G. *Plant Metabolites in Plant Defense Against Pathogens*; IntechOpen: London, UK, 2019; ISBN 978-1-78985-116-8.
10. Zaynab, M.; Fatima, M.; Abbas, S.; Sharif, Y.; Umair, M.; Zafar, M.H.; Bahadar, K. Role of Secondary Metabolites in Plant Defense against Pathogens. *Microb. Pathog.* **2018**, *124*, 198–202. [[CrossRef](#)]
11. Naboulsi, I.; Aboulmouhajir, A.; Kouisni, L.; Bekkaoui, F.; Yasri, A. Plants Extracts and Secondary Metabolites, Their Extraction Methods and Use in Agriculture for Controlling Crop Stresses and Improving Productivity: A Review. *Acad. J. Med. Plants* **2018**, *6*, 223–240.
12. Boulogne, I.; Petit, P.; Ozier-lafontaine, H.; Loranger-merciris, G.; Boulogne, I.; Petit, P.; Ozier-lafontaine, H.; Desfontaines, L.; Loranger-Merciris, G. Insecticidal and Antifungal Chemicals Produced by Plants: A Review. *Environ. Chem. Lett.* **2012**, *10*, 325–347. [[CrossRef](#)]
13. Duško, B.L.; Comiæ, L.; Sukdolak, S. Antibacterial Activity of Some Plants from Family Apiaceae in Relation to Selected Phytopathogenic Bacteria. *Kragujev. J. Sci.* **2006**, *28*, 65–72.
14. Ferrie, A.M.R.; Caswell, K.L. Chapter 13—Applications of Doubled Haploidy for Improving Industrial Oilseeds. In *Industrial Oil Crops*; McKeon, T.A., Hayes, D.G., Hildebrand, D.F., Weselake, R.J., Eds.; AOCS Press: Champaign, IL, USA, 2016; pp. 359–378. ISBN 978-1-893997-98-1.
15. Chaubey, M.K. Biological Effects of Essential Oils Against Rice Weevil *Sitophilus oryzae* L. (Coleoptera: Curculionidae). *J. Essent. Oil Bear. Plants* **2012**, *15*, 809–815. [[CrossRef](#)]
16. Petretto, G.L.; Fancello, F.; Bakhy, K.; Faiz, C.A.; Sibawayh, Z.; Chessa, M.; Zara, S.; Sanna, M.L.; Maldini, M.; Rourke, J.P.; et al. Chemical Composition and Antimicrobial Activity of Essential Oils from *Cuminum cyminum* L. Collected in Different Areas of Morocco. *Food Biosci.* **2018**, *22*, 50–58. [[CrossRef](#)]
17. El Karkouri, J.; Bouhrim, M.; Al Kamaly, O.M.; Mechchate, H.; Kchibale, A.; Adadi, I.; Amine, S.; Alaoui Ismaili, S.; Zair, T. Chemical Composition, Antibacterial and Antifungal Activity of the Essential Oil from *Cistus ladanifer* L. *Plants* **2021**, *10*, 2068. [[CrossRef](#)]
18. Gazim, Z.C.; Demarchi, I.G.; Lonardon, M.V.C.; Amorim, A.C.L.; Hovell, A.M.C.; Rezende, C.M.; Ferreira, G.A.; de Lima, E.L.; de Cosmo, F.A.; Cortez, D.A.G. Acaricidal Activity of the Essential Oil from *Tetradenia riparia* (Lamiaceae) on the Cattle Tick *Rhipicephalus (Boophilus) microplus* (Acari; Ixodidae). *Exp. Parasitol.* **2011**, *129*, 175–178. [[CrossRef](#)]
19. Abdulmanea, K.; Prokudina, E.A.; Lanková, P.; Vaničková, L.; Koblůvská, R.; Zelený, V.; Lapčík, O. Immunochemical and HPLC Identification of Isoflavonoids in the Apiaceae Family. *Biochem. Syst. Ecol.* **2012**, *45*, 237–243. [[CrossRef](#)]
20. Sadaoui, N.; Bec, N.; Barragan-Montero, V.; Kadri, N.; Cuisinier, F.; Larroque, C.; Arab, K.; Khetta, B. The Essential Oil of Algerian *Ammodaucus leucotrichus* Coss. & Dur. and Its Effect on the Cholinesterase and Monoamine Oxidase Activities. *Fitoterapia* **2018**, *130*, 1–5. [[CrossRef](#)]
21. Nikolić, M.; Marković, T.; Ćirić, A.; Glamočlija, J.; Marković, D.; Soković, M. Susceptibility of *Oral candida* Spp.: Reference Strains and Clinical Isolates to Selected Essential Oils of Apiaceae Species. *Lek. Sirovine* **2016**, *35*, 151–162. [[CrossRef](#)]
22. Singh, G.; Maurya, S.; de Lampasona, M.P.; Catalan, C. Chemical Constituents, Antimicrobial Investigations, and Antioxidative Potentials of *Anethum graveolens* L. Essential Oil and Acetone Extract: Part 52. *J. Food Sci.* **2005**, *70*, M208–M215. [[CrossRef](#)]
23. Aćimović, M.G.; Pavlović, S.D.; Varga, A.O.; Filipović, V.M.; Cvetković, M.T.; Stanković, J.M.; Čabarkapa, I.S. Chemical Composition and Antibacterial Activity of Angelica Archangelica Root Essential Oil. *Nat. Prod. Commun.* **2017**, *12*, 205–206. [[CrossRef](#)]
24. Sipailiene, A.; Venskutonis, P.R.; Sarkinas, A.; Cypiene, V. Composition and Antimicrobial Activity of Celery (*Apium graveolens*) Leaf and Root Extracts Obtained with Liquid Carbon Dioxide. *Acta Hort.* **2005**, *677*, 71–77. [[CrossRef](#)]

25. Alsalman, A.-H.; Aboalhaija, N.; Talib, W.; Abaza, I.; Afifi, F. Evaluation of the Single and Combined Antibacterial Efficiency of the Leaf Essential Oils of Four Common Culinary Herbs: Dill, Celery, Coriander and Fennel Grown in Jordan. *J. Essent. Oil Bear. Plants* **2021**, *24*, 317–328. [[CrossRef](#)]
26. Maggi, F.; Giuliani, C.; Fico, G.; Ricciutelli, M.; Bramucci, M.; Quassinti, L.; Petrelli, D.; Vitali, L.A.; Cianfaglione, K.; Tirillini, B.; et al. Secondary Metabolites, Secretory Structures and Biological Activity of Water Celery (*Apium nodiflorum* (L.) Lag.) Growing in Central Italy. *Plant Biosyst.—Int. J. Deal. All Asp. Plant Biol.* **2019**, *153*, 325–335. [[CrossRef](#)]
27. López, S.; Lima, B.; Aragón, L.; Espinar, L.A.; Tapia, A.; Zucchini, S.; Zygadlo, J.; Feresin, G.E.; López, M.L. Essential Oil of Azorella Cryptantha Collected in Two Different Locations from San Juan Province, Argentina: Chemical Variability and Anti-Insect and Antimicrobial Activities. *Chem. Biodivers.* **2012**, *9*, 1452–1464. [[CrossRef](#)] [[PubMed](#)]
28. Abou El-Soud, N.H.; Deabes, M.M.; Abou El-Kassem, L.; Khalil, M.Y. Antifungal Activity of Family Apiaceae Essential Oils. *J. Appl. Sci. Res.* **2012**, *8*, 4964–4973.
29. Synowiec, A.; Możdżeń, K.; Krajewska, A.; Landi, M.; Araniti, F. *Carum carvi* L. Essential Oil: A Promising Candidate for Botanical Herbicide against *Echinochloa crus-galli* (L.) P. Beauv. in Maize Cultivation. *Ind. Crops Prod.* **2019**, *140*, 111652. [[CrossRef](#)]
30. Hajlaoui, H.; Arraouadi, S.; Noumi, E.; Aouadi, K.; Adnan, M.; Khan, M.A.; Kadri, A.; Snoussi, M. Antimicrobial, Antioxidant, Anti-Acetylcholinesterase, Antidiabetic, and Pharmacokinetic Properties of *Carum carvi* L. and *Coriandrum sativum* L. Essential Oils Alone and in Combination. *Molecules* **2021**, *26*, 3625. [[CrossRef](#)]
31. Abdolahi, A.; Hassani, A.; Ghosta, Y.; Javadi, T.; Meshkatalasadat, M.H. Essential Oils as Control Agents of Postharvest Alternaria and Penicillium Rots on Tomato Fruits. *J. Food Saf.* **2010**, *30*, 341–352. [[CrossRef](#)]
32. Khalil, N.; Ashour, M.; Fikry, S.; Singab, A.N.; Salama, O. Chemical Composition and Antimicrobial Activity of the Essential Oils of Selected Apiaceous Fruits. *Future J. Pharm. Sci.* **2018**, *4*, 88–92. [[CrossRef](#)]
33. Fang, R.; Jiang, C.H.; Wang, X.Y.; Zhang, H.M.; Liu, Z.L.; Zhou, L.; Du, S.S.; Deng, Z.W. Insecticidal Activity of Essential Oil of *Carum carvi* Fruits from China and Its Main Components against Two Grain Storage Insects. *Molecules* **2010**, *15*, 9391–9402. [[CrossRef](#)]
34. Simic, A.; Rančić, A.; Sokovic, M.D.; Ristic, M.; Grujic-Jovanovic, S.; Vukojevic, J.; Marin, P.D. Essential Oil Composition of *Cymbopogon Winterianus*. and *Carum carvi*. and Their Antimicrobial Activities. *Pharm. Biol.* **2008**, *46*, 437–441. [[CrossRef](#)]
35. Singh, G.; Marimuthu, P.; de Heluani, C.S.; Catalan, C.A.N. Antioxidant and Biocidal Activities of *Carum nigrum* (Seed) Essential Oil, Oleoresin, and Their Selected Components. *J. Agric. Food Chem.* **2006**, *54*, 174–181. [[CrossRef](#)]
36. Oyedeji, O.A.; Afolayan, A.J. Chemical Composition and Antibacterial Activity of the Essential Oil of *Centella asiatica* Growing in South Africa. *Pharm. Biol.* **2005**, *43*, 249–252. [[CrossRef](#)]
37. Petrović, G.M.; Stamenković, J.G.; Kostevski, I.R.; Stojanović, G.S.; Mitić, V.D.; Zlatković, B.K. Chemical Composition of Volatiles; Antimicrobial, Antioxidant and Cholinesterase Inhibitory Activity of *Chaerophyllum aromaticum* L. (Apiaceae) Essential Oils and Extracts. *Chem. Biodivers.* **2017**, *14*, e1600367. [[CrossRef](#)]
38. Lakušić, B.; Slavkovska, V.; Pavlović, M.; Milenković, M.; Stanković, J.A.; Couladis, M. Chemical Composition and Antimicrobial Activity of the Essential Oil from *Chaerophyllum aureum* L. (Apiaceae). *Nat. Prod. Commun.* **2009**, *4*, 115–118. [[CrossRef](#)]
39. Hayta, S.; Celikeyen, F.C. Evaluation of Essential Oil Composition, Antioxidant and Antimicrobial Properties of *Chaerophyllum Crinitum* Boiss (Apiaceae) from Turkey: A Traditional Medicinal Herb. *J. Biol. Sci.* **2016**, *16*, 72–76. [[CrossRef](#)]
40. Shafaghat, A. Comparison of the Antimicrobial Activity and Chemical Constituents of the Essential Oil and Hexanic Extract from *Chaerophyllum Macropodium*. *J. Essent. Oil Bear. Plants* **2017**, *20*, 835–843. [[CrossRef](#)]
41. Lee, M.-J.; Lee, S.-E.; Kang, M.-S.; Park, B.; Lee, S.-G.; Lee, H.-S. Acaricidal and Insecticidal Properties of *Coriandrum sativum* Oils and Their Major Constituents Extracted by Three Different Methods against Stored Product Pests. *Appl. Biol. Chem.* **2018**, *61*, 481–488. [[CrossRef](#)]
42. Khani, A.; Rahdari, T. Chemical Composition and Insecticidal Activity of Essential Oil from *Coriandrum sativum* Seeds against *Tribolium confusum* and *Callosobruchus maculatus*. *ISRN Pharm.* **2012**, *2012*, 263517. [[CrossRef](#)]
43. Amini, S.; Tajabadi, F.; Khani, M.; Labbafi, M.R.; Tavakoli, M. Identification of the Seed Essential Oil Composition of Four Apiaceae Species and Comparison of Their Biological Effects on *Sitophilus oryzae* L. and *Tribolium castaneum* (Herbst.). *J. Med. Plants* **2018**, *17*, 68–76.
44. Alves-Silva, J.M.; Guerra, I.; Gonçalves, M.J.; Cavaleiro, C.; Cruz, M.T.; Figueirinha, A.; Salgueiro, L. Chemical Composition of *Crithmum maritimum* L. Essential Oil and Hydrodistillation Residual Water by GC-MS and HPLC-DAD-MS/MS, and Their Biological Activities. *Ind. Crops Prod.* **2020**, *149*, 112329. [[CrossRef](#)]
45. D’Agostino, G.; Giambra, B.; Palla, F.; Bruno, M.; Badalamenti, N. The Application of the Essential Oils of *Thymus vulgaris* L. and *Crithmum maritimum* L. as Biocidal on Two Tholu Bommalu Indian Leather Puppets. *Plants* **2021**, *10*, 1508. [[CrossRef](#)] [[PubMed](#)]
46. Pasiyas, I.N.; Ntakoulas, D.D.; Raptopoulou, K.; Gardeli, C.; Proestos, C. Chemical Composition of Essential Oils of Aromatic and Medicinal Herbs Cultivated in Greece—Benefits and Drawbacks. *Foods* **2021**, *10*, 2354. [[CrossRef](#)] [[PubMed](#)]
47. Pavela, R.; Maggi, F.; Lupidi, G.; Cianfaglione, K.; Dauvergne, X.; Bruno, M.; Benelli, G. Efficacy of Sea Fennel (*Crithmum maritimum* L., Apiaceae) Essential Oils against *Culex quinquefasciatus* Say and *Spodoptera littoralis* (Boisd.). *Ind. Crops Prod.* **2017**, *109*, 603–610. [[CrossRef](#)]
48. Suresh, U.; Murugan, K.; Panneerselvam, C.; Aziz, A.T.; Cianfaglione, K.; Wang, L.; Maggi, F. Encapsulation of Sea Fennel (*Crithmum maritimum*) Essential Oil in Nanoemulsion and SiO<sub>2</sub> Nanoparticles for Treatment of the Crop Pest *Spodoptera litura* and the Dengue Vector *Aedes Aegypti*. *Ind. Crops Prod.* **2020**, *158*, 113033. [[CrossRef](#)]

49. Benelli, G.; Pavela, R.; Petrelli, R.; Cappellacci, L.; Canale, A.; Senthil-Nathan, S.; Maggi, F. Not Just Popular Spices! Essential Oils from *Cuminum cyminum* and *Pimpinella anisum* Are Toxic to Insect Pests and Vectors without Affecting Non-Target Invertebrates. *Ind. Crops Prod.* **2018**, *124*, 236–243. [[CrossRef](#)]
50. Abdellaoui, M.; Bouhlali, E.D.T.; Rhaffari, L.E. Chemical Composition and Antioxidant Activities of the Essential Oils of Cumin (*Cuminum cyminum*) Conducted Under Organic Production Conditions. *J. Essent. Oil Bear. Plants* **2019**, *22*, 1500–1508. [[CrossRef](#)]
51. Attia, S.; Grissa, K.L.; Lognay, G.; Heuskin, S.; Mailleux, A.C.; Hance, T. Chemical Composition and Acaricidal Properties of Deverra Scoparia Essential Oil (Araliales: Apiaceae) and Blends of Its Major Constituents against *Tetranychus urticae* (Acari: Tetranychidae). *J. Econ. Entomol.* **2011**, *104*, 1220–1228. [[CrossRef](#)]
52. Glamoclija, J.M.; Sokovic, M.D.; Šiljegovic, J.D.; Ristic, M.S.; Ciric, A.D.; Grubišić, D.V. Chemical Composition and Antimicrobial Activity of *Echinophora spinosa* L. (Apiaceae) Essential Oil. *Rec. Nat. Prod.* **2011**, *5*, 319–323.
53. Kremer, D.; Zovko Končić, M.; Kosalec, I.; Košir, I.J.; Potočnik, T.; Čerenak, A.; Srećec, S.; Dunkić, V.; Vuko, E. Phytochemical Traits and Biological Activity of *Eryngium amethystinum* and *E. alpinum* (Apiaceae). *Horticulturae* **2021**, *7*, 364. [[CrossRef](#)]
54. Casiglia, S.; Bruno, M.; Rosselli, S.; Senatore, F. Chemical Composition and Antimicrobial Activity of the Essential Oil from Flowers of *Eryngium triquetrum* (Apiaceae) Collected Wild in Sicily. *Nat. Prod. Commun.* **2016**, *11*, 1019–1024. [[CrossRef](#)]
55. Merad, N.; Andreu, V.; Chaib, S.; de Carvalho Augusto, R.; Duval, D.; Bertrand, C.; Boumghar, Y.; Pichette, A.; Djabou, N. Essential Oils from Two Apiaceae Species as Potential Agents in Organic Crops Protection. *Antibiotics* **2021**, *10*, 636. [[CrossRef](#)]
56. Karakaya, S.; Göger, G.; Bostanlik, F.D.; Demirci, B.; Duman, H.; Kiliç, C.S. Comparison of the Essential Oils of *Ferula orientalis* L., *Ferulago Sandrasica* Peşmen and Quézel, and *Hippomarathrum microcarpum* Petrov and Their Antimicrobial Activity. *Turk. J. Pharm. Sci.* **2019**, *16*, 69–75. [[CrossRef](#)]
57. Moghaddam, M.; Mehdizadeh, L.; Mirzaei Najafgholi, H.; Ghasemi Pirbalouti, A. Chemical Composition, Antibacterial and Antifungal Activities of Seed Essential Oil of *Ferulago Angulata*. *Int. J. Food Prop.* **2018**, *21*, 158–170. [[CrossRef](#)]
58. Karakaya, S.; Koca, M.; Sytar, O.; Dursunoglu, B.; Ozbek, H.; Duman, H.; Guvenalp, Z.; Kılıç, C.S. Antioxidant and Anticholinesterase Potential of *Ferulago Cassia* with Farther Bio-Guided Isolation of Active Coumarin Constituents. *South Afr. J. Bot.* **2019**, *121*, 536–542. [[CrossRef](#)]
59. Dikpinar, T.; Süzgeç-Selçuk, S.; Çelik, B.Ö.; Uruşak, E.A. Antimicrobial Activity of Rhizomes of *Ferulago Trachycarpa* Boiss. and Bioguided Isolation of Active Coumarin Constituents. *Ind. Crops Prod.* **2018**, *123*, 762–767. [[CrossRef](#)]
60. Tavakoli, S.; Yassa, N.; Delnavazi, M.R.; Akhbari, M.; Hadjiakhoondi, A.; Hajimehdipoor, H.; Khalighi-Sigaroodi, F.; Hajiaghah, R. Chemical Composition and Biological Activities of the Essential Oils from Different Parts of *Ferulago trifida* Boiss. *J. Essent. Oil Res.* **2017**, *29*, 407–419. [[CrossRef](#)]
61. Sayed Ahmad, B.; Talou, T.; Saad, Z.; Hijazi, A.; Cerny, M.; Kanaan, H.; Chokr, A.; Merah, O. Fennel Oil and By-Products Seed Characterization and Their Potential Applications. *Ind. Crops Prod.* **2018**, *111*, 92–98. [[CrossRef](#)]
62. Teke, M.A.; Mutlu, Ç. Insecticidal and Behavioral Effects of Some Plant Essential Oils against *Sitophilus granarius* L. and *Tribolium castaneum* (Herbst). *J. Plant Dis. Prot.* **2021**, *128*, 109–119. [[CrossRef](#)]
63. Kaur, P.; Gupta, S.; Kaur, K.; Kaur, N.; Kumar, R.; Bhullar, M.S. Nanoemulsion of *Foeniculum Vulgare* Essential Oil: A Propitious Striver against Weeds of *Triticum Aestivum*. *Ind. Crops Prod.* **2021**, *168*, 113601. [[CrossRef](#)]
64. Torbati, M.; Nazemiyeh, H.; Lotfipour, F.; Asnaashari, S.; Nemati, M.; Fathiazad, F. Composition and Antibacterial Activity of *Heracleum Transcaasicum* and *Heracleum Anisactis* Aerial Parts Essential Oil. *Adv. Pharm. Bull.* **2013**, *3*, 415–418. [[CrossRef](#)]
65. Moradalizadeh, M.; Akhgar, M.R.; Rajaei, P.; Faghihi-Zarandi, A. Chemical Composition of the Essential Oils of *Levisticum Officinale* Growing Wild in Iran. *Chem. Nat. Compd.* **2012**, *47*, 1007–1009. [[CrossRef](#)]
66. Liu, Z.L.; Chu, S.S.; Jiang, G.H. Insecticidal Activity and Composition of Essential Oil of *Ostericum sieboldii* (Apiaceae) against *Sitophilus zeamais* and *Tribolium castaneum*. *Rec. Nat. Prod.* **2011**, *5*, 74–81.
67. Brusotti, G.; Ibrahim, M.F.; Dentamaro, A.; Gilardoni, G.; Tosi, S.; Grisoli, P.; Dacarro, C.; Guglielminetti, M.L.; Hussain, F.H.S.; Caccialanza, G.; et al. Chemical Composition and Antimicrobial Activity of the Volatile Fractions from Leaves and Flowers of the Wild Iraqi Kurdish Plant *Prangos Peucedanifolia* FENZL. *Chem. Biodivers.* **2013**, *10*, 274–280. [[CrossRef](#)] [[PubMed](#)]
68. Zengin, G.; Stojković, D.; Mahomoodally, M.F.; Jugreet, B.S.; Paksoy, M.Y.; Ivanov, M.; Gašić, U.; Gallo, M.; Montesano, D. Comprehensive Biological and Chemical Evaluation of Two Seseli Species (*S. Gummiferum* and *S. Transcausicum*). *Antioxidants* **2021**, *10*, 1510. [[CrossRef](#)] [[PubMed](#)]
69. Rosato, A.; Maggi, F.; Cianfaglione, K.; Conti, F.; Ciaschetti, G.; Rakotosaona, R.; Fracchiolla, G.; Clodoveo, M.L.; Franchini, C.; Corbo, F. Chemical Composition and Antibacterial Activity of Seven Uncommon Essential Oils. *J. Essent. Oil Res.* **2018**, *30*, 233–243. [[CrossRef](#)]
70. Gandomi, H.; Abbaszadeh, S.; Jebellijavan, A.; Sharifzadeh, A. Chemical Constituents, Antimicrobial and Antioxidative Effects of *Trachyspermum ammi* Essential Oil. *J. Food Processing Preserv.* **2014**, *38*, 1690–1695. [[CrossRef](#)]
71. Demirci, B.; Yasdikcioğlu, G.K.; Başer, K.H.C. Sesquiterpene Hydrocarbons of the Essential Oil of *Actinolema Macrolema* Boiss. *Turk. J. Chem.* **2013**, *37*, 917–926. [[CrossRef](#)]
72. Stefanovic, O.; Comic, L.; Stanojevic, D.; Solujic-Sukdolac, S. Antibacterial Activity of *Aegopodium podagraria* L. Extracts and Interaction between Extracts and Antibiotics. *Turk. J. Biol.* **2009**, *33*, 145–150. [[CrossRef](#)]
73. Pavela, R. Larvicidal Effects of Various Euro-Asiatic Plants against *Culex quinquefasciatus* Say Larvae (Diptera: Culicidae). *Parasitol. Res.* **2008**, *102*, 555–559. [[CrossRef](#)]

74. Fairouz, B.; Nora, C.; Salima, K.G. Insecticidal Effect of *Ammi visnaga* L. (Apiaceae: Apial) Methanolic Extract against a Citrus Pest, *Toxoptera aurantii* (Aphididae: Homoptera) under Controlled Conditions. *J. Entomol. Zool. Stud.* **2016**, *4*, 230–235.
75. Meepagala, K.M.; Estep, A.S.; Becnel, J.J. Mosquitocidal Activity of Extracts from *Ammi visnaga* (Apiaceae) Seeds. *J. Agric. Chem. Environ.* **2016**, *05*, 170–178. [[CrossRef](#)]
76. Ghoneim, K.; Mohammad, A.; Al-Daly, A.; Amer, M.; Khadrawy, F.; Mahmoud, M.A. Metabolic Responsiveness of Desert Locust *Schistocerca gregaria* (Forsk.) (Orthoptera: Acrididae) to the Khella Plant *Ammi visnaga* L. (Apiaceae) Extracts. *Int. J. Adv. Life Sci.* **2014**, *7*, 204–216.
77. Sebaa, A.; Marouf, A.; Kambouche, N.; Derdour, A. Phytochemical Composition, Antioxidant and Antimicrobial Activities of *Ammodaucus leucotrichus* Fruit from Algerian Sahara. *Orient. J. Chem.* **2018**, *34*, 519–525. [[CrossRef](#)]
78. Chaubey, M.K. Insecticidal Activity of *Trachyspermum ammi* (Umbelliferae), *Anethum graveolens* (Umbelliferae) and *Nigella Sativa* (Ranunculaceae) Essential Oils against Stored-Product Beetle *Tribolium castaneum* Herbst (Coleoptera: Tenebrionidae). *Afr. J. Agric. Res.* **2007**, *2*, 596–600. [[CrossRef](#)]
79. Kostić, I.; Lazarević, J.; Šešlija Jovanović, D.; Kostić, M.; Marković, T.; Milanović, S. Potential of Essential Oils from Anise, Dill and Fennel Seeds for the Gypsy Moth Control. *Plants* **2021**, *10*, 2194. [[CrossRef](#)]
80. Evergetis, E.; Michaelakis, A.; Haroutounian, S.A. Essential Oils of Umbelliferae (Apiaceae) Family Taxa as Emerging Potent Agents for Mosquito Control. In *Integrated Pest Management and Pest Control—Current and Future Tactics*; Soloneski, S., Ed.; InTech: London, UK, 2012; pp. 613–638. ISBN 978-953-51-0050-8.
81. Karakas, M. Use of Aromatic Plant Extracts as Bio-Insecticides for the Control of Stored-Product Insect, *Sitophilus granarius*. *Int. J. Entomol. Res.* **2017**, *2*, 27–29.
82. Din, Z.U.; Shad, A.A.; Bakht, J.; Ullah, I.; Jan, S. In Vitro Antimicrobial, Antioxidant Activity and Phytochemical Screening of *Apium graveolens*. *Pak. J. Pharm. Sci.* **2015**, *28*, 1699–1704.
83. Sousa, R.M.O.F.; Rosa, J.S.; Oliveira, L.; Cunha, A.; Fernandes-Ferreira, M. Activities of Apiaceae Essential Oils against Armyworm, *Pseudaletia unipuncta* (Lepidoptera: Noctuidae). *J. Agric. Food Chem.* **2013**, *61*, 7661–7672. [[CrossRef](#)]
84. Ebadollahi, A.; Mahboubi, M. Insecticidal Activity of the Essential Oil Isolated from *Azilia Eryngioides* (PAU) Hedge et Lamond against Two Beetle Pests. *Chil. J. Agric. Res.* **2011**, *71*, 406–411. [[CrossRef](#)]
85. Lima, B.; Sanchez, M.; Agüero, M.B.; Tapia, A.; Palermo, J.A.; Feresin, G.E. Antibacterial Activity of Extracts and Compounds Isolated from the Andean Medicinal Plant *Azorella Cryptantha* (Clos) Reiche, Apiaceae. *Ind. Crops Prod.* **2015**, *64*, 152–157. [[CrossRef](#)]
86. Ben-Khalifa, N.E.; Chaieb, I.; Laarif, A.; Haouala, R. Insecticidal Activity of Six Apiaceae Essential Oils against *Spodoptera littoralis* Biosduval (Lepidoptera: Noctuidae). *J. New Sci.* **2018**, *55*, 3603–3609.
87. Rajkumar, S.; Jebanesan, A. Larvicidal and Adult Emergence Inhibition Effect of *Centella asiatica* Brahmi (Umbelliferae) against Mosquito *Culex quinquefasciatus* Say (Diptera: Culicidae). *Afr. J. Biomed. Res.* **2005**, *8*, 31–33. [[CrossRef](#)]
88. Idris, F.N.; Nadzir, M.M. Antimicrobial Activity of *Centella asiatica* on *Aspergillus Niger* and *Bacillus Subtilis*. *Chem. Eng. Trans.* **2017**, *56*, 1381–1386. [[CrossRef](#)]
89. Shafaghat, A. Antibacterial Activity and Composition of Essential Oils from Flower, Leaf and Stem of *Chaerophyllum Macropodum* Boiss. from Iran. *Nat. Prod. Commun.* **2009**, *4*, 861–864. [[CrossRef](#)] [[PubMed](#)]
90. Matasyoh, J.C.; Maiyo, Z.C.; Ngure, R.M.; Chepkorir, R. Chemical Composition and Antimicrobial Activity of the Essential Oil of *Coriandrum sativum*. *Food Chem.* **2009**, *113*, 526–529. [[CrossRef](#)]
91. Pereira, C.G.; Moraes, C.B.; Franco, C.H.; Feltrin, C.; Grougnet, R.; Barbosa, E.G.; Panciera, M.; Correia, C.R.D.; Rodrigues, M.J.; Custódio, L. In Vitro Anti-Trypanosoma Cruzi Activity of Halophytes from Southern Portugal Reloaded: A Special Focus on Sea Fennel (*Crithmum maritimum* L.). *Plants* **2021**, *10*, 2235. [[CrossRef](#)]
92. Chaubey, M.K. Evaluation of Insecticidal Properties of *Cuminum cyminum* and Piper Nigrum Essential Oils against *Sitophilus zeamais*. *J. Entomol.* **2017**, *14*, 148–154. [[CrossRef](#)]
93. Hrudová, E.; Kocourková, B.; Zelená, V. Insecticidal Effect of Carrot (*Daucus Carota*) and Lovage (*Levisticum Officinale*) (Apiaceae) Extracts against *Tribolium confusum* Jacquelin Du Duval, 1868 (Coleoptera, Tenebrionidae). *Acta Univ. Agric. Silvic. Mendel. Brun.* **2014**, *54*, 165–168. [[CrossRef](#)]
94. Mouloud, G.; Rabah, B.; Khellaf, R. Antimicrobial and Antioxidant Activity of Methanol Extract of *Echinophora spinosa* L. from Jijel, Algeria. *Alger. J. Biosci.* **2020**, *1*, 24–29. [[CrossRef](#)]
95. Pavela, R.; Maggi, F.; Cianfaglione, K.; Canale, A.; Benelli, G. Promising Insecticidal Efficacy of the Essential Oils from the Halophyte *Echinophora spinosa* (Apiaceae) Growing in Corsica Island, France. *Environ. Sci. Pollut. Res.* **2020**, *27*, 14454–14464. [[CrossRef](#)]
96. Lingaraju, D.P.; Sudarshana, M.S.; Mahendra, C.; Rao, K.P. Phytochemical Screening and Antimicrobial Activity of Leaf Extracts of *Eryngium foetidum* L. (Apiaceae). *Indo Am. J. Pharm. Res.* **2016**, *6*, 4339–4344.
97. Sumitha, K.V.; Prajitha, V.; Sandhya, V.N.; Anjana, S.; Thoppil, J.E. Potential Larvicidal Principles in *Eryngium foetidum* L. (Apiaceae), An Omnipresent Weed, Effective Against *Aedes Albopictus* Skuse. *J. Essent. Oil-Bear. Plants* **2014**, *17*, 1279–1286. [[CrossRef](#)]
98. Zengin, G.; Mahomoodally, M.F.; Paksoy, M.Y.; Picot-Allain, C.; Glamocilja, J.; Sokovic, M.; Diuzheva, A.; Jekó, J.; Cziáky, Z.; Rodrigues, M.J.; et al. Phytochemical Characterization and Bioactivities of Five Apiaceae Species: Natural Sources for Novel Ingredients. *Ind. Crops Prod.* **2019**, *135*, 107–121. [[CrossRef](#)]
99. Kahraman, C.; Topcu, G.; Bedir, E.; Tatli, I.I.; Ekizoglu, M.; Akdemir, Z.S. Phytochemical Screening and Evaluation of the Antimicrobial and Antioxidant Activities of *Ferula Caspica* M. Bieb. Extracts. *Saudi Pharm. J.* **2019**, *27*, 525–531. [[CrossRef](#)]

100. Ramezanipour, O.; Yakhcha, M. Study on the Effect of *Ferula Pseudalliacea* (Family: Apiaceae) Extract on *Varroa Destructor* (Acari: Varroidae) Infestation in Honeybee (Hymenoptera: Apidae, Apis Melifera). *Iran. Vet. J.* **2020**, *15*, 85–92. [[CrossRef](#)]
101. Zengin, G.; Sinan, K.I.; Ak, G.; Mahomoodally, M.F.; Paksoy, M.Y.; Picot-Allain, C.; Glamocilja, J.; Sokovic, M.; Jekó, J.; Cziáky, Z.; et al. Chemical Profile, Antioxidant, Antimicrobial, Enzyme Inhibitory, and Cytotoxicity of Seven Apiaceae Species from Turkey: A Comparative Study. *Ind. Crops Prod.* **2020**, *153*, 112572. [[CrossRef](#)]
102. Goodarzi, S.; Tavakoli, S.; Abai, M.R.; Amini, Z.; Vatandoost, H.; Yassa, N.; Hadjiakhoondi, A.; Tofighi, Z. Strong Insecticidal Potential of Methanol Extract of *Ferulago trifida* Fruits against *Anopheles Stephensi* as Malaria Vector. *Environ. Sci. Pollut. Res.* **2019**, *26*, 7711–7717. [[CrossRef](#)]
103. Ebadollahi, A. Susceptibility of Two Sitophilus Species (Coleoptera: Curculionidae) to Essential Oils from *Foeniculum Vulgare* and *Satureja hortensis*. *Ecol. Balk.* **2011**, *3*, 1–8.
104. Modise, S.A.; Ashafa, A.O.T. Larvicidal, Pupicidal and Insecticidal Activities of *Cosmos Bipinnatus*, *Foeniculum Vulgare* and *Tagetes Minuta* against *Culex quinquefasciatus* Mosquitoes. *Trop. J. Pharm. Res.* **2016**, *15*, 965–972. [[CrossRef](#)]
105. Al-Mekhlafi, F.A.; Abutaha, N.; Al-Doaiss, A.A.; Ahmed Al-Keridis, L.; Alsayadi, A.I.; Ali El Hadi Mohamed, R.; Wadaan, M.A.; Elfaki Ibrahim, K.; Al-Khalifa, M.S. Target and Non-Target Effects of *Foeniculum Vulgare* and *Matricaria Chamomilla* Combined Extract on *Culex Pipiens* Mosquitoes. *Saudi J. Biol. Sci.* **2021**, *28*, 5773–5780. [[CrossRef](#)]
106. Torbati, M.; Nazemiyeh, H.; Lotfipour, F.; Nemati, M.; Asnaashari, S.; Fathiazad, F. Chemical Composition and in Vitro Antioxidant and Antibacterial Activity of *Heracleum Transcaucasicum* and *Heracleum Anisactis* Roots Essential Oil. *BiolImpacts* **2014**, *4*, 69–74. [[PubMed](#)]
107. Sedaghat, M.M.; Sanei Dehkordi, A.; Abai, M.R.; Khanavi, M.; Mohtarami, F.; Salim Abadi, Y.; Rafi, F.; Vatandoost, H. Larvicidal Activity of Essential Oils of Apiaceae Plants against Malaria Vector, *Anopheles Stephensi*. *Iran. J. Arthropod-Borne Dis.* **2011**, *5*, 51–59. [[PubMed](#)]
108. Izakmehri, K.; Saber, M.; Mehrvar, A.; Hassanpouraghdam, M.B.; Vojoudi, S. Lethal and Sublethal Effects of Essential Oils from *Eucalyptus Camaldulensis* and *Heracleum persicum* against the Adults of *Callosobruchus maculatus*. *J. Insect Sci.* **2013**, *13*, 152. [[CrossRef](#)] [[PubMed](#)]
109. Rančić, A.; Soković, M.; Vukojević, J.; Simić, A.; Marin, P.; Duletić-Laušević, S.; Djoković, D. Chemical Composition and Antimicrobial Activities of Essential Oils of *Myrrhis odorata* (L.) Scop, *Hypericum perforatum* L. and *Helichrysum arenarium* (L.) Moench. *J. Essent. Oil Res.* **2005**, *17*, 341–345. [[CrossRef](#)]
110. Matejić, J.S.; Džamić, A.M.; Mihajilov-krstev, T.; Randelović, V.N. Antimicrobial Potential of Essential Oil from *Pastinaca sativa* L. *Biol. Nyssana* **2014**, *5*, 31–35.
111. Vitalini, S.; Palmioli, A.; Orlando, F.; Scari, G.; Airoidi, C.; De Noni, I.; Bocchi, S.; Iriti, M. Phytotoxicity, Nematicidal Activity and Chemical Constituents of *Peucedanum ostruthium* (L.) W.D.J.Koch (Apiaceae). *Ind. Crops Prod.* **2021**, *166*, 113499. [[CrossRef](#)]
112. Mohamed, H.S.A.A.; Abdelgadir, W.S.; Almagboul, A.Z.I. In Vitro Antimicrobial Activity of Anise Seed (*Pimpinella anisum* L.). *Int. J. Adv. Res.* **2015**, *3*, 359–367.
113. Soni, R.; Sharma, G.; Jasuja, N.D. Essential Oil Yield Pattern and Antibacterial and Insecticidal Activities of *Trachyspermum ammi* and *Myristica Fragrans*. *Scientifica* **2016**, *2016*, 1428194. [[CrossRef](#)]
114. Chaubey, M.K. Study of Insecticidal Properties of *Trachyspermum ammi* and *Mentha Arvensis* Essential Oils against *Sitophilus zeamais* L. (Coleoptera: Curculionidae). *Curr. Life Sci.* **2017**, *4*, 10–17. [[CrossRef](#)]
115. Bisrat, D.; Jung, C. Insecticidal Toxicities of Three Main Constituents Derived from *Trachyspermum ammi* (L.) Sprague Ex Turrill Fruits against the Small Hive Beetles, *Aethina Tumida* Murray. *Molecules* **2020**, *25*, 1100. [[CrossRef](#)]
116. Chaudhari, A.K.; Singh, V.K.; Kedia, A.; Das, S.; Dubey, N.K. Essential Oils and Their Bioactive Compounds as Eco-Friendly Novel Green Pesticides for Management of Storage Insect Pests: Prospects and Retrospects. *Environ. Sci. Pollut. Res.* **2021**, *28*, 18918–18940. [[CrossRef](#)]
117. Jankowska, M.; Rogalska, J.; Wyzkowska, J.; Stankiewicz, M. Molecular Targets for Components of Essential Oils in the Insect Nervous System—A Review. *Molecules* **2017**, *23*, 34. [[CrossRef](#)]
118. Prakash, B.; Kumar, A.; Singh, P.P.; Das, S.; Dubey, N.K. Chapter 16—Prospects of Plant Products in the Management of Insect Pests of Food Grains: Current Status and Future Perspectives. In *Natural Bioactive Compounds*; Sinha, R.P., Häder, D.-P., Eds.; Academic Press: Cambridge, MA, USA, 2021; pp. 317–335. ISBN 978-0-12-820655-3.
119. Rattan, R.S. Mechanism of Action of Insecticidal Secondary Metabolites of Plant Origin. *Crop Prot.* **2010**, *29*, 913–920. [[CrossRef](#)]
120. Ivănescu, B.; Burlec, A.F.; Crivoi, F.; Roșu, C.; Corciovă, A. Secondary Metabolites from *Artemisia* Genus as Biopesticides and Innovative Nano-Based Application Strategies. *Molecules* **2021**, *26*, 3061. [[CrossRef](#)]
121. Radhakrishnan, R.; Alqarawi, A.A.; Abd\_Allah, E.F. Bioherbicides: Current Knowledge on Weed Control Mechanism. *Ecotoxicol. Environ. Saf.* **2018**, *158*, 131–138. [[CrossRef](#)]
122. Verdeguer, M.; Sánchez-Moreiras, A.M.; Araniti, F. Phytotoxic Effects and Mechanism of Action of Essential Oils and Terpenoids. *Plants* **2020**, *9*, 1571. [[CrossRef](#)]
123. Raveau, R.; Fontaine, J.; Lounès-Hadj Sahraoui, A. Essential Oils as Potential Alternative Biocontrol Products against Plant Pathogens and Weeds: A Review. *Foods* **2020**, *9*, 365. [[CrossRef](#)]
124. Ebadollahi, A. Plant Essential Oils from Apiaceae Family as Alternatives to Conventional Insecticides. *Ecol. Balk.* **2013**, *5*, 149–172.
125. Pavela, R. Acaricidal Properties of Extracts and Major Furanochromenes from the Seeds of *Ammi visnaga* Linn. against *Tetranychus urticae* Koch. *Ind. Crops Prod.* **2015**, *67*, 108–113. [[CrossRef](#)]



126. Benelli, G.; Pavela, R.; Ricciutelli, M.; Lupidi, G.; Maggi, F. Efficacy of the Volatile Oil from Water Celery (*Helosciadium nodiflorum*, Apiaceae) against the Filariasis Vector *Culex quinquefasciatus*, the Housefly *Musca Domestica*, and the African Cotton Leafworm *Spodoptera littoralis*. *Chem. Biodivers.* **2017**, *14*, e1700376. [[CrossRef](#)]
127. Bell, C.H. 15—Pest Control of Stored Food Products: Insects and Mites. In *Hygiene in Food Processing*, 2nd ed.; Lelieveld, H.L.M., Holah, J.T., Napper, D., Eds.; Woodhead Publishing Series in Food Science, Technology and Nutrition; Woodhead Publishing: Cambridge, UK, 2014; pp. 494–538. ISBN 978-0-85709-429-2.
128. Fouad, H.A.; de Souza Tavares, W.; Zanoncio, J.C. Toxicity and Repellent Activity of Monoterpene Enantiomers to Rice Weevils (*Sitophilus oryzae*). *Pest Manag. Sci.* **2021**, *77*, 3500–3507. [[CrossRef](#)] [[PubMed](#)]
129. Rosa, J.S.; Oliveira, L.; Sousa, R.M.O.F.; Escobar, C.B.; Fernandes-Ferreira, M. Bioactivity of Some Apiaceae Essential Oils and Their Constituents against *Sitophilus zeamais* (Coleoptera: Curculionidae). *Bull. Entomol. Res.* **2020**, *110*, 406–416. [[CrossRef](#)] [[PubMed](#)]
130. Kaur, V.; Kaur, R.; Bhardwaj, U. A Review on Dill Essential Oil and Its Chief Compounds as Natural Biocide. *Flavour Fragr. J.* **2021**, *36*, 412–431. [[CrossRef](#)]
131. Azirak, S.; Karaman, S. Allelopathic Effect of Some Essential Oils and Components on Germination of Weed Species. *Acta Agric. Scand. Sect. B Soil Plant Sci.* **2008**, *58*, 88–92. [[CrossRef](#)]
132. Sabzi Nojaded, M.; Pouresmaeil, M.; Younessi-Hamzkehlanlu, M.; Venditti, A. Phytochemical Profile of Fennel Essential Oils and Possible Applications for Natural Antioxidant and Controlling *Convolvulus arvensis* L. *Nat. Prod. Res.* **2021**, *35*, 4164–4168. [[CrossRef](#)] [[PubMed](#)]
133. Pujari, J.D.; Yakkundimath, R.; Byadgi, A.S. Image Processing Based Detection of Fungal Diseases in Plants. *Procedia Comput. Sci.* **2015**, *46*, 1802–1808. [[CrossRef](#)]
134. Silva, R.N.; Monteiro, V.N.; Steindorff, A.S.; Gomes, E.V.; Noronha, E.F.; Ulhoa, C.J. Trichoderma/Pathogen/Plant Interaction in Pre-Harvest Food Security. *Fungal Biol.* **2019**, *123*, 565–583. [[CrossRef](#)]
135. Carmona-Hernandez, S.; Reyes-Pérez, J.J.; Chiquito-Contreras, R.G.; Rincon-Enriquez, G.; Cerdan-Cabrera, C.R.; Hernandez-Montiel, L.G. Biocontrol of Postharvest Fruit Fungal Diseases by Bacterial Antagonists: A Review. *Agronomy* **2019**, *9*, 121. [[CrossRef](#)]
136. Di Napoli, M.; Varcamonti, M.; Basile, A.; Bruno, M.; Maggi, F.; Zanfardino, A. Anti-Pseudomonas Aeruginosa Activity of Hemlock (*Conium maculatum*, Apiaceae) Essential Oil. *Nat. Prod. Res.* **2019**, *33*, 3436–3440. [[CrossRef](#)]
137. Mith, H.; Duré, R.; Dalcenserie, V.; Zhiri, A.; Daube, G.; Clinquart, A. Antimicrobial Activities of Commercial Essential Oils and Their Components against Food-Borne Pathogens and Food Spoilage Bacteria. *Food Sci. Nutr.* **2014**, *2*, 403–416. [[CrossRef](#)]
138. Mostafa, A.A.; Al-Askar, A.A.; Almaary, K.S.; Dawoud, T.M.; Sholkamy, E.N.; Bakri, M.M. Antimicrobial Activity of Some Plant Extracts against Bacterial Strains Causing Food Poisoning Diseases. *Saudi J. Biol. Sci.* **2018**, *25*, 361–366. [[CrossRef](#)]
139. Duarte, A.; Luís, Â.; Oleastro, M.; Domingues, F.C. Antioxidant Properties of Coriander Essential Oil and Linalool and Their Potential to Control *Campylobacter* Spp. *Food Control* **2016**, *61*, 115–122. [[CrossRef](#)]
140. Kačániová, M.; Galovičová, L.; Ivanišová, E.; Vukovic, N.L.; Štefániková, J.; Valková, V.; Borotová, P.; Žiarovská, J.; Terentjeva, M.; Felšöciová, S.; et al. Antioxidant, Antimicrobial and Antibiofilm Activity of Coriander (*Coriandrum sativum* L.) Essential Oil for Its Application in Foods. *Foods* **2020**, *9*, 282. [[CrossRef](#)]
141. Das, S.; Kumar Singh, V.; Kumar Dwivedy, A.; Kumar Chaudhari, A.; Deepika; Kishore Dubey, N. Nanostructured *Pimpinella anisum* Essential Oil as Novel Green Food Preservative against Fungal Infestation, Aflatoxin B1 Contamination and Deterioration of Nutritional Qualities. *Food Chem.* **2021**, *344*, 128574. [[CrossRef](#)]
142. El-Soud, N.H.A.; Deabes, M.; El-Kassem, L.A.; Khalil, M. Chemical Composition and Antifungal Activity of *Ocimum basilicum* L. Essential Oil. *Open Access Maced. J. Med. Sci.* **2015**, *3*, 374. [[CrossRef](#)]
143. Lasram, S.; Zemni, H.; Hamdi, Z.; Chenenaoui, S.; Houissa, H.; Saidani Tounsi, M.; Ghorbel, A. Antifungal and Antiaflatoxinogenic Activities of *Carum carvi* L., *Coriandrum sativum* L. Seed Essential Oils and Their Major Terpene Component against *Aspergillus Flavus*. *Ind. Crops Prod.* **2019**, *134*, 11–18. [[CrossRef](#)]
144. Alinezhad, S.; Kamalzadeh, A.; Shams-Ghahfarokhi, M.; Rezaee, M.-B.; Jaimand, K.; Kawachi, M.; Zamani, Z.; Tolouei, R.; Razzaghi-Abyaneh, M. Search for Novel Antifungals from 49 Indigenous Medicinal Plants: *Foeniculum Vulgare* and *Platycladus Orientalis* as Strong Inhibitors of Aflatoxin Production by *Aspergillus Parasiticus*. *Ann. Microbiol.* **2011**, *61*, 673–681. [[CrossRef](#)]
145. Kedia, A.; Prakash, B.; Mishra, P.K.; Dubey, N.K. Antifungal and Antiaflatoxinogenic Properties of *Cuminum cyminum* (L.) Seed Essential Oil and Its Efficacy as a Preservative in Stored Commodities. *Int. J. Food Microbiol.* **2014**, *168–169*, 1–7. [[CrossRef](#)]
146. Maurya, A.; Kumar, S.; Singh, B.K.; Chaudhari, A.K.; Dwivedy, A.K.; Prakash, B.; Dubey, N.K. Mechanistic Investigations on Antifungal and Antiaflatoxinogenic Activities of Chemically Characterised *Carum carvi* L. Essential Oil against Fungal Infestation and Aflatoxin Contamination of Herbal Raw Materials. *Nat. Prod. Res.* **2021**, 1–6. [[CrossRef](#)]
147. Hazrati, H.; Saharkhiz, M.J.; Niakousari, M.; Moein, M. Natural Herbicide Activity of *Satureja hortensis* L. Essential Oil Nanoemulsion on the Seed Germination and Morphophysiological Features of Two Important Weed Species. *Ecotoxicol. Environ. Saf.* **2017**, *142*, 423–430. [[CrossRef](#)]
148. Chouhan, S.; Sharma, K.; Guleria, S. Antimicrobial Activity of Some Essential Oils—Present Status and Future Perspectives. *Medicines* **2017**, *4*, 58. [[CrossRef](#)] [[PubMed](#)]
149. Pascual-Villalobos, M.J.; Cantó-Tejero, M.; Vallejo, R.; Guirao, P.; Rodríguez-Rojo, S.; Cocero, M.J. Use of Nanoemulsions of Plant Essential Oils as Aphid Repellents. *Ind. Crops Prod.* **2017**, *110*, 45–57. [[CrossRef](#)]
150. Venkadesaperumal, G.; Rucha, S.; Sundar, K.; Shetty, P.H. Anti-Quorum Sensing Activity of Spice Oil Nanoemulsions against Food Borne Pathogens. *LWT Food Sci. Technol.* **2016**, *66*, 225–231. [[CrossRef](#)]