



**HAL**  
open science

## Multi-physics modelling of freeze-thaw cycles effects on tree branches

Cyril Bozonnet, M. Saudreau, Eric Badel, Thierry Améglio, Guillaume Charrier

► **To cite this version:**

Cyril Bozonnet, M. Saudreau, Eric Badel, Thierry Améglio, Guillaume Charrier. Multi-physics modelling of freeze-thaw cycles effects on tree branches. 10th Plant Biomechanics conference, Aug 2022, Lyon, France. , 2022, proceeding 10th Plant Biomechanics conference, Lyon-France, 22-26th Aug 2022. hal-03727914

**HAL Id: hal-03727914**

**<https://hal.inrae.fr/hal-03727914>**

Submitted on 19 Jul 2022

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Multi-physics modelling of freeze-thaw cycles effects on tree branches

Cyril Bozonnet, Marc Saudreau, Eric Badel, Thierry Améglio, Guillaume Charrier

Université Clermont Auvergne, INRAE, PIAF, 63000 Clermont-Ferrand, France

## Abstract

Frost hardiness is the main factor affecting plant species distribution at high latitudes and altitudes. The main effects of freeze-thaw cycles on trees are damages to living cells, as well as the formation of gas embolism in xylem vessels. Frost effect on trees is also quantified through changes in branch diameter.

In order to resorb embolism, some species (walnut, maple, birch, etc.) exhibit an increase in xylem sap pressure during successive freeze-thaw cycles. The modelling of such phenomenon is very challenging due to its multi-physics and multi-scale nature. In this work, we present a numerical model coupling heat transfer, phase change, water and osmotic fluxes, taking into consideration different cell types within walnut branch tissues. We show how diameter and pressure variations are inter-related, and we validate the model against experimental results from the literature. We eventually show how this work can be adapted in order to explore inter-species differences.