



**HAL**  
open science

## Deciphering the link between sugar transport and vascular system development: A SWEET story

Beate Hoffmann, Emilie Aubry, Françoise Vilaine, Catherine C. Bellini, Sylvie Dinant, Rozenn Le Hir

### ► To cite this version:

Beate Hoffmann, Emilie Aubry, Françoise Vilaine, Catherine C. Bellini, Sylvie Dinant, et al.. Deciphering the link between sugar transport and vascular system development: A SWEET story. *Plant Vascular Biology*, Jul 2022, Berlin, Germany. . hal-03731708

**HAL Id: hal-03731708**

**<https://hal.inrae.fr/hal-03731708>**

Submitted on 21 Jul 2022

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Deciphering the link between sugar transport and vascular system development: A SWEET story

Hoffmann B<sup>1</sup>, Aubry E<sup>1</sup>, Vilaine F<sup>1</sup>, Bellini C<sup>1,2</sup>, Dinant S<sup>1</sup> and Le Hir R<sup>1</sup>

<sup>1</sup> *Université Paris-Saclay, INRAE, AgroParisTech, Institut Jean-Pierre Bourgin (IJPB), 78000 Versailles, France*

<sup>2</sup> *Umeå Plant Science Centre, Department of Plant Physiology, Umeå University, 90183 Umeå, Sweden*

In plant cells, carbohydrates are the main building blocks for central metabolism, for soluble sugars storage and polysaccharides synthesis. Sugars also serve as signaling molecules and osmotica during abiotic stress responses. Intercellular and intracellular transport systems are therefore, not only critical for long-distance sugar allocation but also to modulate local sugar partitioning ranging from tissular to subcellular scales.

The vascular system including phloem, (pro)cambium and xylem, is an important tissue for sugar transport. A complex network of transcription factors controls its formation. At the cellular level, cytosolic sugar availability is regulated by sugar exchanges at the plasma membrane and the tonoplast, through the transport of sugars by plasmodesmata as well as active and/or facilitated transporters, suggesting a complexity in fine tuning of sugar transport and homeostasis in the vascular tissues. However, the balance between nutrient/sugars availability in the vascular cells and their use for the development of the vascular tissues remains to be addressed.

We are interested in the link between sugar transport and the plant vascular system development. Among the various families of sugar transporters, we are focusing our analysis on SWEET transporters, the latest sugar transporter family identified, that facilitate the movement of soluble sugars along the concentration gradient. By using an integrated approach, at the cell and the whole plant levels, we functionally characterized in *Arabidopsis thaliana* several members of the SWEET transporter family and show that they are important for the vascular system development and the secondary cell wall formation. Additionally, our work shed new light on the central role of vascular parenchyma cells in these processes and further highlights the importance of finely regulated intercellular and/or intracellular sugar transport to assure appropriate plant growth and development.