



HAL
open science

Impact of the physiological state of fungal spores on their inactivation by active chlorine and hydrogen peroxide

Karim Rigalma

► **To cite this version:**

Karim Rigalma. Impact of the physiological state of fungal spores on their inactivation by active chlorine and hydrogen peroxide. *Food Microbiology*, 2021, 100, pp.103850. 10.1016/j.fm.2021.103850 . hal-03736423

HAL Id: hal-03736423

<https://hal.inrae.fr/hal-03736423v1>

Submitted on 2 Aug 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial 4.0 International License

1 **Impact of the physiological state of fungal spores on their inactivation by active chlorine**
2 **and hydrogen peroxide**

3

4 Vincent VISCONTI, Karim RIGALMA, Emmanuel COTON and Philippe DANTIGNY*

5

6 Univ Brest, Laboratoire Universitaire de Biodiversité et Ecologie Microbienne, F-29280 Plouzané,
7 France

8

9 ***Corresponding author** : Philippe DANTIGNY

10 Laboratoire Universitaire de Biodiversité et Ecologie Microbienne,

11 Parvis Blaise-Pascal, Technopôle Brest-Iroise

12 29280 Plouzané, France

13 Tel : +33 (0)2.90.91.51.11

14 E-mail : philippe.dantigny@univ-brest.fr

15

16 **Keywords:** *Aspergillus*, *Cladosporium*, *Mucor*, *Penicillium*, fungicide, disinfectant, predictive
17 **mycology**

18

19

20 **Abstract**

21 This study aimed at assessing the impact of the physiological state of fungal spores on
22 inactivation by sodium hypochlorite, 0.1% and 0.2% active chlorine, and 3% hydrogen
23 peroxide. In this context, two physiological states were compared for 4 fungal species (5
24 strains). The first physiological state corresponded to fungal spores produced at 0.99 a_w and
25 harvested using an aqueous solution (laboratory conditions), while the second one
26 corresponded to fungal spores produced under a moderate water stress (0.95 a_w) and dry-
27 harvested (mechanical harvesting without use of any water, mimicking food plant conditions).
28 *Aspergillus flavus* “food plant” conidia were more resistant to all tested fungicide molecules
29 than the “laboratory” ones. The same phenomenon was observed for *Penicillium commune*
30 UBOCC-A-116003 conidia treated with hydrogen peroxide. However, this isolate did not
31 exhibit any inactivation difference between “laboratory” and “food plant” conidia treated with
32 sodium hypochlorite. Similarly, the physiological state of *Cladosporium cladosporioides*
33 conidia did not impact the efficacy of the tested biocides. *P. commune* UBOCC-A-112059
34 “food plant” and “laboratory” conidia were more resistant to hydrogen peroxide and sodium
35 hypochlorite, respectively. As for *Mucor circinelloides*, “laboratory” spores were more
36 resistant to all disinfectant than the “food plant” ones. Noteworthy, regardless of the
37 physiological state, all *M. circinelloides* and *C. cladosporioides* conidia were inactivated for 5
38 min treatment at 0.2% active chlorine and for 2.5 min treatment at 0.1% active chlorine, while
39 the conidia of all the other species remained viable for these treatments. The obtained data
40 indicate that the efficacy of disinfectant molecules depends not only on the encountered
41 fungal species and its intraspecific diversity but also on the spore physiological state.

42

44 1. Introduction

45 Efficacy of fungicide molecules depends on various treatment conditions, i.e., nature of the
46 disinfectant, contact time, concentration, application conditions (temperature, pH, relative
47 humidity, dirtiness of the surfaces, presence of surfactants...) and mode (e.g. fumigation,
48 liquid application) and obviously the targeted fungi. For sanitization efficacy tests, the
49 standard fungal strains are *Aspergillus brasiliensis* (ATCC 16404) and *Candida albicans*
50 (ATCC 10231). According to the European Committee for Standardization, other species can
51 be tested, i.e., *Absidia corymbifera* (IP 1129 75), *Candida albicans* (IP 1180 79),
52 *Cladosporium cladosporioides* (IP 1232 80), *Penicillium verrucosum* var. *cyclopium* (IP 1231
53 80) to assess fungicidal efficacy (KSG France, 2021). Beyond the fact that the proposed
54 species may not be the most representative of food plant contaminants and, as recently shown
55 by Scaramuzza et al. (2020), might even be more sensitive than the latter; in all cases, fungal
56 spores have to be produced under optimal growth conditions (i.e., temperature, time,
57 atmosphere, medium, (AFNOR NF T 72-281, 2014; European Standard 1275, 2006;
58 European Standard 1650, 2019; European Standard 13624, 2013; European Standard
59 13697+A1, 2019). These standards recommend Malt Extract Agar (pH 5.6, 0.99 a_w) to
60 produce fungal spores that are subsequently diluted into peptonized water. In contrast to these
61 spores produced in laboratory conditions, in food plant conditions, spores are airborne and not
62 rehydrated, and unlikely to be produced under optimal conditions.

63 The physiological state of *Penicillium* spp. strains was affected by temperature, water activity
64 and pH during conidia production (Nguyen van Long et al., 2017a). Moreover, the impact of
65 the harvesting protocol on the susceptibility to disinfectants can be important as, for example,
66 wet harvested *Rhizopus stolonifer* conidia had a different physiological state than dry-
67 harvested ones. Even brief exposure (30 min) to water during harvesting caused spores to

68 engage metabolic activity (Nicherson et al., 1981). The effects of environmental conditions
69 during sporogenesis on the germination time is well documented (Blaszyk et al., 1998; Judet
70 et al., 2008; Nanguy et al., 2010; Nguyen van Long et al., 2017a, 2017b; Lattab et al., 2012;
71 Ruijten et al., 2020; Stevenson et al., 2016). Less information on the impact of environmental
72 conditions on the susceptibility to disinfectants is available. Dry-harvested *Penicillium*
73 conidia exposed to ethanol vapours for 24h did not show any inactivation whereas aqueous
74 suspensions of *P. digitatum* and *P. italicum* conidia exhibited about 3 log and 2 log
75 inactivation, respectively (Dao and Dantigny, 2009). The same authors showed that the
76 contact time with ethanol vapours (0.67 kPa) required to inactivate 4 log *P. chrysogenum* and
77 *P. digitatum* conidia produced at either 0.99 a_w or in the 0.85-0.90 a_w range were about 30h
78 and 120h, respectively (Dao and Dantigny, 2009).

79 Production of conidia at reduced water activity increases the amount of protective compatible
80 solutes in the cytoplasm (Andersen et al., 2006; Chandler et al., 1984; de Lima Alves et al.,
81 2015; Hallsworth and Magan, 1995; Magan, 2006, 2007; Nesci et al., 2014; Rangel et al.,
82 2015; Wyatt et al., 2013, 2015) that can contribute to a certain resistance. It was also
83 hypothesized that both the production of conidia at reduced water activity and application of a
84 dry-harvesting protocol contributed to lower the intracellular water activity of fungal spores
85 (Dao and Dantigny, 2009), thus explaining their resistance to ethanol which is an hydrophilic
86 molecule. This hypothesis was strengthened by the observation that for 1 min exposure to 70%
87 ethanol, only dry-harvested conidia of four *Penicillium commune* strains produced at 0.95 a_w
88 exhibited survivors, while suspensions of conidia produced at 0.995 a_w were completely
89 inactivated (Visconti et al. 2020).

90 This study aimed at extending the comparison between “laboratory” conidia (production at
91 0.995 a_w and harvested using an aqueous solution) and “food plant” (production at 0.95 a_w
92 and dry-harvested) conidia to other fungal species and disinfectant molecules. The ethanol

93 resistant *P. commune* UBOCC-A-116003 and the ethanol sensitive *P. commune* UBOCC-A-
94 112059 (Visconti et al. 2020) were selected, in addition to species that are frequently isolated
95 from dairy product environments, namely *Cladosporium cladosporioides*, *Mucor*
96 *circinelloides* and *Aspergillus flavus*. The tested disinfectant molecules corresponded to
97 sodium hypochlorite (0.1 and 0.2% active chlorine) and hydrogen peroxide (3%).

98

99 **2. Material and methods**

100 *2.1. Molds*

101 The studied molds were provided by the Université de Bretagne Occidentale Culture
102 Collection (UBOCC, <https://www.univ-brest.fr/ubocc/>) and were originally isolated from
103 spoiled dairy products (Table 1). Molds were maintained on Potato Dextrose Agar (PDA)
104 medium at 4°C.

105

106 *2.2. Spores production*

107 “Laboratory” spores were produced from mycelium grown on PDA at 0.995 a_w for 7 days at
108 25°C and harvested in sterile saline solution (NaCl, 9 g/l of water) containing Tween 80
109 (0.015% v/v).

110 “Food plant” spores were produced from mycelium grown on PDA at 0.950 a_w for 7 days at
111 25°C. Water activity was adjusted by substituting a part of the water with an equal weight of
112 glycerol. The relative amount of glycerol was 20.6% (w/w) to obtain PDA media at 0.950 a_w .

113 Spores were harvested mechanically without contact with any liquid solution (dry-harvest)
114 according to Dao and Dantigny (2009). Briefly, ten sterile glass beads (diameter 3 mm) were
115 deposited on the mycelium and the Petri dish was shaken gently to detach the conidia. Then,
116 the plate was turned upside-down and the conidia were harvested on the lid by gently tapping
117 100 times on the bottom of the Petri dish.

118

119 2.3. *Disinfectants*

120 Sodium hypochlorite test solutions were prepared from a stock solution diluted in distilled
121 water containing Tween 80 (0.015%, v/v). The stock solution was prepared from 250 ml
122 bleach boxes (Eau de Javel La Croix, Bois-Colombes, France) that contained 4.8% active
123 chlorine (5% hypochlorite), diluted with distilled water according to the supplier's
124 recommendation to reach final concentrations of 0.1 and 0.2% active chlorine. The co-
125 formulants mentioned by the supplier were 5-15% chlorine bleach and sodium hydroxide. The
126 hydrogen peroxide test solution was prepared from 30% hydrogen peroxide solution (Sigma
127 Aldrich, Saint-Quentin Fallavier, France) diluted in distilled water containing Tween 80
128 (0.015%, v/v) to reach a 3% final concentration.

129

130 2.4. *Disinfectant testing procedures*

131 Treatment of “food plant” spores started by applying 10 ml of biocide solution to the dry-
132 conidia on the lid. For *M. circinelloides*, this volume was reduced to 5 ml as fewer spores
133 were harvested. Prior to the first sampling, the suspension (biocide + conidia) was transferred
134 to a Falcon tube and homogenized by vortexing for 10 s. Treatment of “laboratory” spores
135 began when 1ml of spore suspension was transferred in 9 ml of biocide solution and the mix
136 vortexed. For sodium hypochlorite, spores survival was measured every 2.5 min for 20 min
137 except for *M. circinelloides* for which an additional contact time of 1 min was done. For
138 hydrogen peroxide, spores survival was measured after 1, 5, 10, 15, 30 and 60 min of contact
139 time. Both biocides were neutralized by diluting ten-fold the suspension in 1% sodium
140 thiosulfate pentahydrate diluted in saline aqueous solution (NaCl, 9 g/l) containing Tween 80
141 (0.015% v/v). The initial amount of “food plant” spores (N_0) was evaluated by counting on a
142 Malassez cell according to Dao et al. (2008). The method was based on the principle that the

143 application of biocide does not affect the morphology of the spores during the time of the
144 experiment so it is possible to count under the microscope all the spores treated including
145 inactivated ones.

146

147 2.5. Viability assessment

148 After neutralization, the treated spores were grown on PDA at 25 °C for 3 days by spreading
149 100 µl of sample and four subsequent decimal dilutions. *M. circinelloides* spores were grown
150 only for 1 day. Only counts (N) in the range of 10 to 100 colonies per plate were considered.
151 Inactivation of conidia was expressed as the logarithmic reduction factor, $\log_{10} (N/N_0)$.

152

153 3. Results

154 3.1. Impact of disinfectant molecules on “laboratory” spores

155 At 0.1% active chlorine, *C. cladosporioides* was the most sensitive species (Fig. 1A), as for a
156 contact time of 2.5 min, *C. cladosporioides* showed 2.8 log inactivation. Inactivation were
157 less than or equal to 0.2 log for all the other tested species. At 12.5 min of exposure, *P.*
158 *commune* only exhibited survivors, the UBOCC-A-112059 isolate was more resistant than the
159 UBOCC-A-116003.

160 For a contact time of 2.5 min to 0.2% active chlorine, total inactivation of *C. cladosporioides*
161 and *M. circinelloides* spores was observed, (Fig. 1B). At 7.5 min of exposure, *P. commune*
162 only exhibited survivors. Inactivation values were 1.9 log and 3.1 log for the UBOCC-A-
163 112059 and UBOCC-A-116003 strains, respectively. Under these experimental conditions, all
164 *A. flavus* conidia were inactivated.

165 By increasing the active chlorine concentration from 0.1 to 0.2%, the contact time leading to a
166 complete inactivation of the conidia was reduced from 5 to 2.5 min and, from 15 to 10 min,

167 for *C. cladosporioides* and *P. commune*, respectively. Regardless of the tested concentration,
168 the former and the latter species were the most sensitive and most resistant to active chlorine,
169 respectively.

170 In contrast to active chlorine, *P. commune* was the most sensitive species to 3% hydrogen
171 peroxide (Fig. 1C). For a 10 min contact time, all spores of the two tested strains were
172 inactivated. After 30 min of exposure, all *A. flavus* spores were inactivated, inactivation being
173 greater than 4 log. In contrast to this species, inactivation was only 0.2 log for *C.*
174 *cladosporioides* and *M. circinelloides*, which were the most resistant species.

175

176 3.2. Impact of fungicide molecules on "food plant" spores

177 As for the physiological "food plant" state, *M. circinelloides* was the most susceptible mold to
178 0.1% active chlorine. At 2.5 min of exposure, all *M. circinelloides* spores were inactivated,
179 inactivation being greater than 4.3 log (Fig. 2A). For this contact time, the inactivation of *C.*
180 *cladosporioides* was equal to 2.7 log, and less than or equal to 0.2 log for *A. flavus* and *P.*
181 *commune*. For the "food plant" physiological state, *A. flavus* was the most resistant species,
182 the only one to exhibit survivors for a contact time of 17.5 min.

183 By increasing the active chlorine concentration, the contact time necessary to inactivate all the
184 spores was reduced. After 2.5 min of exposure, all *C. cladosporioides* conidia and *M.*
185 *circinelloides* spores were inactivated, while inactivation was lower than or equal to 0.4 log
186 for *A. flavus* and *P. commune* (Fig. 2B). At 0.2% active chlorine, *P. commune* UBOCC-A-
187 116003 was more resistant than the UBOCC-A-112059 strain. No *A. flavus* survivors were
188 observed for a contact time of 12.5 min.

189 *A. flavus* was the most sensitive species to 3% hydrogen peroxide. A complete inactivation
190 was observed for a contact time of 30 min (Fig. 2C). After 60 min exposure, only *C.*

191 *cladosporioides* (1.2 log inactivation) and *P. commune* UBOCC-A-112059 (2.4 log
192 inactivation) exhibited survivors. All spores of *M. circirnelloides* and *P. commune* UBOCC-A-
193 116003 were inactivated (> 4 and > 4.4 log, respectively).

194

195 3.3. Impact of the physiological state on the disinfectant efficacy

196 In order to assess the impact of the physiological state on the efficacy of the fungicide
197 molecules, inactivation were compared at the same time, although this time depended on the
198 considered molecule and species. Whenever possible, the shortest contact time leading to no
199 survivors, for either "laboratory" or "food plant" spores, was chosen (Table 2). Four different
200 cases were distinguished, i)"laboratory" spores were more resistant than "food plant" ones, ii)
201 no significant differences were observed between the two physiological states, iii)
202 impossibility to determine whether the two kind of spores exhibited significant differences
203 and iv)"food plant" spores were more resistant than "laboratory" ones.

204 For *M. circirnelloides*, the "laboratory" spores were more resistant than the "food plant"
205 spores to two disinfectants. For a contact time of 2.5 min at 0.1% active chlorine, inactivation
206 was 0.1 log and greater than 4.3 log for the "laboratory" and "food plant" spores, respectively.
207 The difference in inactivation between the two physiological states was greater than 4.2 log.
208 After exposure to 0.2% active chlorine for 1 min, inactivation was 0.2 log and greater than 4.3
209 log for "laboratory" and "food plant" spores, respectively. The difference in inactivation
210 between the two physiological states was greater than 4.1 log. For *P. commune* UBOCC-A-
211 112059, the "laboratory" conidia were more resistant than the "food plant" ones to active
212 chlorine. For a contact time of 12.5 min at 0.1% active chlorine, inactivation was 1.5 log and
213 3 log for the "laboratory" and "food plant" conidia, respectively. The difference in
214 inactivation between the two physiological states was 1.5 log. At 0.2% active chlorine for 1

215 min, inactivation was 1.9 and 4.6 log for "laboratory" and "food plant" conidia, respectively,
216 the difference in inactivation being 2.7 log between the two physiological states.

217 No significant impact was observed between the inactivation by sodium hypochlorite of the
218 two physiological states of *P. commune* UBOCC-A-116003. For a contact time of 12.5 min at
219 0.1% active chlorine, inactivation was 3 and 3.1 log for the "laboratory" and "food plant"
220 spores, respectively. For a contact time of 7.5 min at 0.2% active chlorine, inactivation was
221 3.1 and 3.5 log for "laboratory" and "food plant" physiological states, respectively. For *C.*
222 *cladosporioides*, no significant difference was observed between the two physiological states
223 at 0.1% active chlorine and 3% hydrogen peroxide. Indeed, at 0.1% active chlorine for 2.5
224 min, the inactivation was 2.7 and 2.8 log for the "food plant" and "laboratory" spores,
225 respectively, while, at 3% hydrogen peroxide for 60 min, inactivation was 1.2 and 1.4 log for
226 the same physiological states, respectively.

227 As for the impact of physiological state after exposure to 0.2% active chlorine, it was
228 impossible to determine as all conidia were inactivated from 2.5 min (inactivation > than 4.1
229 log).

230 For *A. flavus*, the "food plant" conidia were more resistant than the "laboratory" spores for the
231 two tested disinfectants. For a contact time of 12.5 min at 0.1% active chlorine, inactivation
232 was 0.7 log and greater than 3.7 log for the "food plant" and "laboratory" conidia,
233 respectively, difference in inactivation between the two physiological states being greater than
234 3 log. For 0.2% active chlorine for 7.5 min, inactivation was 0.9 log and greater than 4.3 log
235 for the "food plant" and "laboratory" conidia, respectively, and the difference in inactivation
236 between the two physiological states was greater than 3.4 log. For a contact time of 15 min
237 with 3% hydrogen peroxide, inactivation was 1.4 log and 3.6 log for the "food plant" and
238 "laboratory" conidia, respectively. The difference in inactivation between the two
239 physiological states was 2.2 log. Finally, for 3% hydrogen peroxide, the two *P. commune*

240 "food plant" conidia were more resistant than the "laboratory" ones. At a contact time of 10
241 min, inactivation of *P. commune* UBOCC-A-112059 was nil and greater than 4.1 log for the
242 "food plant" and "laboratory" conidia, respectively. The difference in inactivation between the
243 two physiological states was greater than 4.1 log. For the same contact time, inactivation of *P.*
244 *commune* UBOCC-A-116003 was 0.7 log and greater than 4.2 log for the "food plant" and
245 "laboratory" conidia, respectively (difference in inactivation between the two physiological
246 states being greater than 3.5 log).

247

248 *3.4. Efficacy of fungicide molecules on the species*

249 With the exception of *C. cladosporioides* and *M. circinelloides*, a contact time of 5 min to
250 0.1% active chlorine or 2.5 min to 0.2% active chlorine had almost no effect on the tested
251 species. However, inactivation obtained for a contact time of 7.5 min to 0.2% active chlorine
252 were of the same order of magnitude than treatments to 0.1% active chlorine for 12.5 min,
253 except for "food plant" *P. commune* UBOCC-A-112059 conidia. However, these conditions
254 were not sufficient to completely inactivate the two *P. commune* strains and the *A. flavus*
255 "food plant" conidia. These conidia were the most resistant ones to sodium hypochlorite.

256 A contact time of 15 min to 3% hydrogen peroxide did not inactivate the spores of the tested
257 species, except the "laboratory" *P. commune* conidia. In contrast to sodium hypochlorite, the
258 *A. flavus* "food plant" conidia were not the most resistant ones to hydrogen peroxide.
259 Surprisingly, *C. cladosporioides* and *M. circinelloides* exhibited a stronger resistance than the
260 other species to a 15 min treatment to 3% hydrogen peroxide. Overall, at the respective tested
261 concentrations, a longer time was required to inactivate the mold species with hydrogen
262 peroxide than with sodium hypochlorite.

263

264 4. Discussion

265 In order to assess a fungicidal efficacy, many studies examined the effect of disinfectant
266 molecules for many contact times, but very few studies described inactivation kinetics. For
267 chlorine inactivation, a classical linear decrease of the log survivors *versus* time with (for
268 *Aspergillus fumigatus*) or without (for *Aspergillus versicolor* and *Penicillium purpurogenum*)
269 an initial lag phase was observed (Ma and Bibby, 2017). In our study, a clear presence of an
270 initial lag phase was shown for the inactivation of “food plant” *P. commune* conidia with
271 0.1% active chlorine. Due to very fast inactivation kinetics leading to very few experimental
272 data, our results did not show a clear linear decrease in the log survivors. However,
273 inactivation curves exhibiting a tailing off (Pereira et al., 2013; Salomão et al., 2009) were not
274 observed in our study. Inactivation kinetics in drinking water were evaluated by means of Ct
275 (mg.min.l⁻¹), i.e., concentration x time (Ma and Bibby, 2017, Pereira et al., 2013), these
276 approaches assumed an inactivation rate proportional to the disinfectant concentration. For a
277 given Ct value, inactivation should be the same regardless of the time and the concentration.
278 In our study, inactivation for 0.1% active chlorine x 5 min and 0.2% active chlorine x 2.5 min
279 were not significantly different regardless of the considered species or physiological state.
280 Although this approach was not validated for other concentration x time conditions or other
281 disinfectants.

282 Although the concept of dry-harvested spores was introduced as early as 1981 by Nickerson
283 et al., almost all publications concern spores produced under optimal conditions and
284 suspended into aqueous solutions (designated as “laboratory” spores in this study). In this
285 context, Andrews (1986) found that 0.37% hypochlorite efficiently killed *A. flavus* after a 2
286 min contact time. In another study, seven *A. flavus* isolates conidia were treated with a diluted
287 commercial disinfectant for 72h. Five isolates did not show any survivors, i.e., log reduction
288 greater than or equal to 2.7, for sodium hypochlorite concentrations in the range 0.4-1.6%.

289 However, the two other isolates exhibited survivors at 1.6% (Mattei et al., 2013). Eventually,
290 4.3 log reduction for *A. flavus* conidia exposed to 3% hypochlorite (2.88% active chlorine) for
291 10 min was reported by Bundgaard-Nielsen and Nielsen (1995). In our study, no survivors
292 were detected after 10 min treatment to 0.2% active chlorine, thus suggesting that the *A.*
293 *flavus* isolate tested in our study was more sensitive than that of the other study. This is a
294 potential indication of behaviour heterogeneity within a given species.

295 The effect of sodium hypochlorite (0.1, 0.5 and 1%), was tested against *C.*
296 *cladosporioides* for 15 min (Bernardi et al., 2018) exhibiting log reductions greater than 4. In
297 our study no survivors (log reduction greater than 3.9) were detected for 5 min treatment at
298 0.1%. As Bernardi et al. (2018) did not carry out treatment at contact times different than 15
299 min, it is therefore difficult to compare. Noteworthy, these authors also tested *P. commune*
300 conidia that exhibited 2-2.9 log reduction and more than 4 log reduction at 0.1% and 0.5/1%
301 active chlorine, respectively. Our results showed 1.5 and 3 log reductions depending on the
302 considered *P. commune* strain for 12.5 min treatments at 0.1% active chlorine. Therefore, the
303 obtained results in this study are in accordance with those of Bernardi et al. (2018) despite
304 slightly different contact times and methods (i.e., carrier method in Bernardi et al. (2018) and
305 suspension method in this study). However, as previously mentioned for *A. flavus*,
306 intraspecific diversity may also be an important factor to explain contrasted inactivation
307 results, as 10 min treatments at 3% hypochlorite led to log inactivation in the range 1.7->5.2
308 for other *P. commune* isolates (Bundgaard-Nielsen and Nielsen, 1995). In agreement with
309 disinfectant susceptibility heterogeneity within a considered species, these authors also
310 reported great differences of behaviour of five *Penicillium roqueforti* isolates when exposed
311 to 0.3% hypochlorite for 10 min (log reductions ranged from 0.5 for the most resistant strain
312 to 4.5 for the most sensitive one, while for the other strains, reductions were in the 1-2 log
313 range).

314 For 15 min treatments with hydrogen peroxide (at 20°C), 3 log reduction in viable counts
315 were obtained for 12.5%, *Absidia corymbifera*, 3.9%, *Candida albicans*, 5.51%,
316 *Scopulariopsis brevicaulis*, 1.71%, *Aspergillus versicolor* and *Penicillium cyclopium*, and
317 0.39%, *Geotrichum candidum* (Martin and Maris, 2012). In this study, 3 log and 3.7 log
318 reduction of *A. flavus* conidia were obtained for 10 min and 15 min treatments with 3%
319 hydrogen peroxide, respectively. For this species, a hydrogen peroxide concentration close to
320 3% would have been required to obtain 3 log reduction in 15 min. According to our results,
321 concentrations lower than (for both *P. commune* strains) and greater than (for *C.*
322 *cladosporioides* and *M. circinelloides*) than 3% would have been necessary to obtain the same
323 reduction.

324 In contrast to peroxide hydrogen, *P. commune* was more resistant than *C. cladosporioides* to
325 active chlorine. This result is in accordance with the results reported by Bernardi et al. (2018)
326 indicating, for 15 min treatments at 0.1% sodium hypochlorite, 2-2.9 log and more than 4 log
327 reduction in *P. commune* and *C. cladosporioides* viable conidia, respectively. In accordance
328 with Bundgaard-Nielsen and Nielsen (1995), *P. commune* was more resistant than *A. flavus* to
329 active chlorine. Overall, both *P. commune* strains were more resistant to active chlorine than
330 *A. flavus*, followed by *C. cladosporioides* and *M. circinelloides*. Surprisingly, the rank of
331 resistance was the reverse than the one observed for hydrogen peroxide, *C. cladosporioides* >
332 *M. circinelloides* > *A. flavus* > *P. commune*.

333 Noteworthy, the resistance ranking differed slightly for “food plant” spores. *A. flavus* was the
334 more resistant and the more sensitive species to active chlorine and hydrogen peroxide,
335 respectively. The rank of resistance depended also on the considered strain; indeed, whereas
336 *P. commune* UBOCC-A-116003 was more resistant than the UBOCC-A-112059 to 70%
337 ethanol (Visconti et al., 2020), UBOCC-A-112059 was more resistant than UBOCC-A-
338 116003 to 3% hydrogen peroxide. This study also showed that differences in resistance

339 depended on the concentration of the disinfectant, at 0.1% active chlorine no difference was
340 noticed between the two *P. commune* strains, while, at 0.2% active chlorine, UBOCC-A-
341 116003 was more resistant than UBOCC-A-112059. At the tested concentrations in this study,
342 active chlorine was more efficient than hydrogen peroxide. In this study, *P. commune*
343 UBOCC-A-112059 “laboratory” conidia were more resistant than the other isolate to active
344 chlorine, but less resistant than the other isolate to hydrogen peroxide. The impact of the
345 physiological state was greater than 1.54 log inactivation for *P. commune* UBOCC-A-116003
346 treated with 70% ethanol (Visconti et al., 2020), but the impact was even greater than 3.5 log
347 for 10 min contact time to 3% peroxide hydrogen. “Food plant” *A. flavus* conidia were also
348 more resistant than “laboratory” ones, but *M. circinelloides* “laboratory” spores were more
349 resistant to all disinfectants than the “food plant” ones. In contrast to the other species, *M.*
350 *circinelloides* can hardly develop at water activities below 0.90 a_w (Hocking and Miscamble,
351 1995; Tresner and Hayes, 1971) and at 0.95 a_w , the growth rate was shown to about half the
352 optimum growth rate (Morin-Sardin et al., 2016). In contrast, *A. flavus* is a xerophilic species
353 that can grow at water activities less than 0.85 a_w , (Pitt and Miscamble, 1995; Sautour et al.,
354 2001, 2002). At 0.95 a_w the growth rate was almost the same than that observed at the
355 optimum water activity of 0.974 (Sautour et al., 2001). These observations suggested that the
356 increased sensitivity to disinfectants of *M. circinelloides* “food plant” spores may be due to
357 the fact that these spores were produced at a water activity (0.95 a_w) too close to the minimum
358 one.

359

360 **5. Conclusions**

361 Surface disinfection in food industries should be based on the use of most effective sanitizer,
362 for the best “contact time / concentration” combination and taking into account the most
363 resistant mold, in agreement with a “worst case scenario” approach. In many cases, a lag

364 phase prior to inactivation was observed. Therefore, shortening the contact time could lead to
365 the elimination of the disinfectant even before fungal spores are actually killed. What can be
366 considered as the most resistant mold depends not only on the encountered species and strains
367 within a given species (disinfectant susceptibility heterogeneity), but also, as demonstrated in
368 this study, on the physiological state of fungal spores and even on the considered disinfectant
369 molecule. Based on the obtained data, it is recommended to assess fungicidal activity of
370 disinfectants on “food plant” spores, not only because there were more resistant than
371 “laboratory” ones for the most resistant species, but because the “food plant” physiological
372 state was closer to airborne conidia that are responsible for actual food contamination and
373 spoilage.

374

375 **Acknowledgements**

376 This work was funded by the French Dairy Interbranch Organization (CNIEL) (Moisibio
377 Project, 2019_00621) and the French Association for Research and Technology (ANRT)
378 (CIFRE 2018/0036).

379

380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402

References

- Agence Française de Normalisation NF T 72-281, 2014. Procédés de désinfection des surfaces par voie aérienne - Détermination de l'activité bactéricide, fongicide, levuricide, mycobactéricide, tuberculocide sporicide et virucide incluant les bactériophages.
- Andersen, M., Magan, N., Mead, A., Chandler, D., 2006. Development of a population based threshold model of conidial germination for analysing the effects of physiological manipulation on the stress tolerance and infectivity of insect pathogenic fungi. *Environ. Microbiol.* 8, 1625–1634.
- Andrews, S., 1986. Optimisation of conditions for the surface disinfection of sorghum and sultanas using sodium hypochlorite solutions. *In: A.D. King, J.I. Pitt, L.R. Beuchat and J.E.L. Corry (editors), Methods for the Mycological Examination of Food.* Plenum Press, New York. pp. 28-32.
- Andrews, S., 1996. Evaluation of surface disinfection procedures for enumerating fungi in foods: a collaborative study. *Int. J. Food Microbiol.* 29, 177-184.
- Andrews, S., Pardoel, D., Harun, A., Treloar, T., 1997. Chlorine inactivation of fungal spores on cereal grains. *Int. J. Food Microbiol.* 35, 153-162.
- Bernardi, A.O., Stefanello, A., Garcia, M.V., Parussolo, G., Stefanello, R.F., Moro, C.B., Copetti, M.V., 2018. Efficacy of commercial sanitizers against fungi of concern in the food industry. *LWT – Food Sci. Technol.* 97, 25-30.

403 Blaszyk, M., Blank, G., Holley, R., Chong, J., 1998. Reduced water activity during
404 sporogenesis in selected *Penicillia*: impact on spore quality. Food Res. Int. 31, 503-509.

405 Bundgaard-Nielsen, K. and Nielsen, P.V., 1995. Fungicidal effect of 15 disinfectants against
406 25 fungal contaminants commonly found in bread and cheese manufacturing. J. Food
407 Prot. 59, 268-275.

408 Chandler, D., Crowe, L.M., Chapman, D., 1984. A population based threshold model to
409 demonstrate that physiological manipulation of endogenous reserves can increase the
410 virulence of insect pathogenic fungi. BCPC Crop Science and Technology 4B, 309–314.

411 Dao, T., Dantigny, P., 2009. Preparation of fungal conidia impacts their susceptibility to
412 inactivation by ethanol vapours. Int. J. Food Microbiol. 135, 268–273.

413 de Lima Alves, F., Stevenson, A., Baxter, E., Gillion, J.L.M., Hejazi, F., Hayes, S., Morrison,
414 I.E.G., Prior, B.A., McGenity, T.J., Rangel, D.E.N., Magan, N., Timmis, K.N.,
415 Hallsworth, J.E., 2015. Concomitant osmotic and chaotropicity-induced stresses in
416 *Aspergillus wentii*: compatible solutes determine the biotic window. Curr. Genet. 61,
417 457–477.

418 European Standard n. 1275, 2006. Chemical disinfectants and antiseptics Quantitative
419 suspension test for the evaluation of basic fungicidal or basic yeasticidal activity of
420 chemical disinfectants and antiseptics - Test method and requirements (Phase 1)

421 European Standard n. 13624, 2013. Chemical disinfectants and antiseptics - Quantitative
422 suspension test for the evaluation of fungicidal or yeasticidal activity in the medical area
423 - Test method and requirements (phase 2, step 1)

424 European Standard n. 13697+A1, 2019. Chemical Disinfectants and Antiseptics –
425 Quantitative Non-porous Surface Test for the Evaluation of Bactericidal and/or

426 Fungicidal Activity of Chemical Disinfectants Used in Food, Industrial, Domestic and
427 Institutional Areas –Test Method and Requirements Without Mechanical Action (Phase
428 2, Step 2).

429 European Standard n. 1650, 2019. Chemical disinfectants and antiseptics - Quantitative
430 suspension test for the evaluation of fungicidal or yeasticidal activity of chemical
431 disinfectants and antiseptics used in food, industrial, domestic and institutional areas -
432 Test method and requirements (Phase 2, step 1)

433 Hallsworth, J.E., Magan, N., 1995. Manipulation of intracellular glycerol and erythritol
434 enhances germination of conidia at low water availability. *Microbiology* 141, 1109–
435 1115.

436 Hocking, A.D. and Miscamble, B.F., 1995. Water relations of some Zygomycetes isolated
437 from food. *Mycol. Res.* 99, 1113-1119.

438 Judet, D., Bensoussan, M., Perrier-Cornet, J.-M., Dantigny, P., 2008. Distributions of the
439 growth rate of the germ tubes and germination time of *Penicillium chrysogenum* conidia
440 depend on water activity. *Food Microbiol.* 25, 902-907.

441 KSG France, 2021. Les Normes Afnor ou EN, les méthodes, les marquages et les étiquettes.
442 [https://www.ksg-france.fr/les-normes-afnor-ou-en-les-methodes-les-marquages-et-les-](https://www.ksg-france.fr/les-normes-afnor-ou-en-les-methodes-les-marquages-et-les-etiquettes/)
443 [etiquettes/](https://www.ksg-france.fr/les-normes-afnor-ou-en-les-methodes-les-marquages-et-les-etiquettes/) Accessed date: 26 March 2021.

444 Lattab, N., Kalai, S., Bensoussan, M., Dantigny, P., 2012. Effect of storage conditions
445 (relative humidity, duration, and temperature) on the germination time of *Aspergillus*
446 *carbonarius* and *Penicillium chrysogenum*. *Int. J. Food Microbiol.* 160, 80–84.

447 Ma, X. and Bibby, K., 2017. Free chlorine and monochloramine inactivation kinetics of
448 *Aspergillus* and *Penicillium* in drinking water. *Water Res.* 120, 265-271.

449 Magan, N., 2006. Ecophysiology of biocontrol agents for improved competence in the
450 phyllosphere. *In*: Bailey, M.J., Lilley, A.K., Timms-Wilson, T.M., Spenser-Phillips,
451 P.T.N. (Eds.), *Microbial Ecology of Aerial Plant Surfaces*. In CABI International, 149-
452 164.

453 Magan, N., 2007. Fungi in extreme environments. In *Environmental and Microbial*
454 *Relationships*, In: Kubicek, C.P., Druzhinina, I.S. (Eds.), *The Mycota IV*, 2nd edn.
455 Springer-Verlag, Berlin, Germany, pp. 85–103.

456 Martin, H. and Maris, P., 2012. Synergism between hydrogen peroxide and seventeen acids
457 against five agri-food-borne fungi and one yeast strain. *J. Appl. Microbiol.* 113, 1451-
458 1460.

459 Mattei, A.S., Madrid, I.M., Santin, R., Schuch, L.F.D., Meireles, M.C.A., 2013. *In vitro*
460 activity of disinfectants against *Aspergillus* spp. *Braz. J. Microbiol.* 44, 481-484.

461 Morin-Sardin, S., Rigalma, K., Coroller, L., Jany, J-L., Coton, E., 2016. Effect of
462 temperature, pH, and water activity on *Mucor* spp. growth on synthetic medium, cheese
463 analog and cheese. *Food Microbiol.* 56, 69-79.

464 Nanguy, S.P.-M., Perrier-Cornet, J.-M., Bensoussan, M., Dantigny, P., 2010. Impact of water
465 activity of diverse media on spore germination of *Aspergillus* and *Penicillium* species.
466 *Int. J. Food Microbiol.* 142, 273-276.

467 Nesci, A., Etcheverry, M., Magan, N., 2004. Osmotic and matric potential effects on growth,
468 sugar alcohol and sugar accumulation by *Aspergillus* section Flavi strains from
469 Argentina. *J. Appl. Microbiol.* 96, 965–972.

470 Nguyen Van Long, N., Rigalma, K., Coroller, L., Dadure, R., Debaets, S., Mounier, J.,
471 Vasseur, V., 2017b. Modelling the effect of water activity reduction by sodium chloride

472 or glycerol on conidial germination and radial growth of filamentous fungi encountered
473 in dairy foods. *Food Microbiol.* 68, 7-15.

474 Nguyen Van Long, N., Vasseur, V., Coroller, L., Dantigny, P., Le Panse, S., Weill, A.,
475 Mounier, J., Rigalma, K., 2017a. Temperature, water activity and pH during conidia
476 production affect the physiological state and germination time of *Penicillium* species.
477 *Int. J. Food Microbiol.* 241, 151-160.

478 Nickerson, K.W., Freer, S.N., Etten, J.L.V., 1981. *Rhizopus stolonifer* sporangiospores: a wet-
479 harvested spore is not a native spore. *Exp. Mycol.* 5, 189-192.

480 Pitt, J.I. and Miscamble, B.F., 1995. Water relations of *Aspergillus flavus* and related species.
481 *J. Food Prot.* 58, 86-90.

482 Rangel, D.E.N., Braga, G.U.L., Fernandes, É.K.K., Keyser, C.A., Hallsworth, J.E., Roberts,
483 D.W., 2015. Stress tolerance and virulence of insect-pathogenic fungi are determined by
484 environmental conditions during conidial formation. *Curr. Genet.* 61, 383–404.

485 Ruijten, P., Huinink, H.P., Adan, O.C.G., 2020. *Penicillium rubens* germination on desiccated
486 and nutrient-depleted conditions depends on the water activity during sporogenesis.

487 Salamão, B.C.M., Churey, J.J., Aragão, G.M.F., Worobo, R.W., 2009. Modeling *Penicillium*
488 *expansum* resistance to thermal and chlorine treatments. *J. Food Microbiol.* 72, 2618-
489 2622.

490 Scaramuzza, N., Mutti, P., Cigarini, M., Berni, E., 2020. Effect of peracetic acid on
491 ascospore-forming molds and test microorganisms used for bio-validations of sanitizing
492 processes in food plants. *Int. J. Food Microbiol.* 332:108772

493 Sautour, M., Dantigny, P., Divies, C., Bensoussan, M., 2001. A temperature-type model for
494 describing the relationship between fungal growth and water activity. Int. J. Food
495 Microbiol. 67, 63-69.

496 Sautour, M., Soares Mansur C., Divies, C., Bensoussan, M., Dantigny, P., 2001. Comparison
497 of the effects of temperature and water activity on growth rate of food spoilage moulds.
498 J. Ind. Microbiol. Biotechnol. 28, 311-315.

499 Stevenson, A., Hamill, P.G., Medina, A., Kminek, G., Rummel, J.D., Dijksterhuis, J., Timson,
500 D.J., Magan, N., Leong, S.-L.L., Hallsworth, J.E., 2016. Glycerol enhances fungal
501 germination at the water-activity limit for life. Environ. Microbiol. 19, 947-967.

502 Tresner, H.D. and Hayes, J.A., 1971. Sodium chloride tolerance of terrestrial fungi. Appl.
503 Microbiol. 22, 210-213.

504 Visconti, V., Rigalma, K., Coton, E., Dantigny, P., 2020. Impact of intraspecific variability
505 and physiological state on *Penicillium commune* inactivation by 70% ethanol. Int. J.
506 Food Microbiol. 332, 108782.

507 Wyatt, T.T., Golovina, E.A., van Leeuwen, M.R., Hallsworth, J.E., Wösten, H.A.B.,
508 Dijksterhuis, J., 2015. A decrease in bulk water and mannitol and accumulation of
509 trehalose and trehalose-based oligosaccharides define a two-stage maturation process
510 towards extreme stress resistance in ascospores of *Neosartorya fischeri* (*Aspergillus*
511 *fischeri*). Environ. Microbiol. 17, 383–394.



512 Wyatt, T.T., Wösten, H.A.B., Dijksterhuis, J., 2013. Fungal spores for dispersion in space and
513 time. Adv. Appl. Microbiol. 85, 43–91.

514

Figure legends

Fig. 1. Inactivation of “laboratory” spores (produced at 0.99 a_w and harvested using an aqueous solution) of *Aspergillus flavus* (●), *Cladosporium cladosporioides* (◆), *Mucor circinelloides* (■), *Penicillium commune* (UBOCC-A-112059) (▲) and *Penicillium commune* (UBOCC-A-116003) (▼). Treatment with A) 0.1% active chlorine, B) 0.2% active chlorine and C) 3% hydrogen peroxide. Linked symbols = presence of survivors and isolated symbol = absence of survivors. Error bars represent standard deviations.

Fig. 2. Inactivation of “food plant” spores (produced at 0.95 a_w and dry-harvested) of *Aspergillus flavus* (⊙), *Cladosporium cladosporioides* (◇), *Mucor circinelloides* (□), *Penicillium commune* (UBOCC-A-112059) (▲) and *Penicillium commune* (UBOCC-A-116003) (▼). Treatment with A) 0.1% active chlorine, B) 0.2% active chlorine and C) 3% hydrogen peroxide. Linked symbols = presence of survivors and isolated symbol = absence of survivors. Error bars represent standard deviations.








Figure 1






Figure 2

Table 1. List of strains provided by the Université de Bretagne Occidentale Culture Collection (UBOCC).

Species	Reference collection number	Origin
<i>Aspergillus flavus</i> var <i>columnaris</i>	UBOCC-A-108066	Butter
<i>Cladosporium cladosporioides</i>	UBOCC-A-111114	Fresh dairy product
<i>Mucor circinelloides</i>	UBOCC-A-112187	Yoghurt
<i>Penicillium commune</i>	UBOCC-A-112059	Blue-veined cheese
<i>Penicillium commune</i>	UBOCC-A-116003	Blue-veined cheese

Table 2. Log₁₀ reductions obtained after exposure to various disinfectants using the liquid method.

Mold (Strain)		Inactivation values (log ₁₀)									
		<i>Aspergillus flavus</i> (UBOCC-A-108066)		<i>Cladosporium cladosporioides</i> (UBOCC-A-111114)		<i>Mucor circinelloides</i> (UBOCC-A-112187)		<i>Penicillium commune</i> (UBOCC-A-112059)		<i>Penicillium commune</i> (UBOCC-A-116003)	
Physiological state		Labo- ratory	Food plant	Labo- ratory	Food plant	Labo- ratory	Food plant	Labo- ratory	Food plant	Labo- ratory	Food plant
Sodium hypochlorite 0.1% active chlorine	2.5 min	0.0 (± 0.1)	0.2 (± 0.1)	2.8 (± 0.2)	2.7 (± 0.2)	0.1* (± 0.1)	> 4.3*	0.0 (± 0.1)	0.1 (± 0.1)	0.2 (± 0.1)	0.1 (± 0.1)
	5 min	0.2 (± 0.1)	0.1 (± 0.1)	> 3.9	> 4.6	> 3.2	> 4.3	0.0 (± 0.2)	0.4 (± 0.1)	0.3 (± 0.1)	0.2 (± 0.1)
	12.5 min	> 3.7*	0.7* (± 0.2)						1.5* (± 0.1)	3.0* (± 0.2)	3.0 (± 0.2)
Sodium hypochlorite 0.2% active chlorine	1 min					0.2* (± 0.1)	> 4.3*				
	2.5 min	0.5 (± 0.1)	0.4 (± 0.1)	> 4.1	> 4.5	> 3.2	> 4.3	0.1 (± 0.1)	0.2 (± 0.1)	0.2 (± 0.1)	0.2 (± 0.1)
	7.5 min	> 4.3*	0.9* (± 0.1)					1.9* (± 0.1)	4.6* (± 0.9)	3.1 (± 0.1)	3.5 (± 0.2)
Hydrogen peroxide 3%	10 min	3.0* (± 0.1)	0.8* (± 0.1)	0.1 (± 0.1)	0.4 (± 0.1)	0.0* (± 0.1)	0.8* (± 0.1)	> 4.1*	0.0* (± 0.1)	> 4.2*	0.7* (± 0.1)
	15 min	3.6* (± 0.4)	1.4* (± 0.1)	0.1 (± 0.1)	0.4 (± 0.1)	0.1* (± 0.1)	1.1* (± 0.2)	> 4.1*	0.5* (± 0.1)	> 4.2*	1.5* (± 0.1)
	60 min	> 4.0	> 3.5	1.4 (± 0.1)	1.2 (± 0.1)	1.3* (± 0.2)	> 4.0*	> 4.1*	2.4* (± 0.1)	> 4.2	> 4.4

“Laboratory”: spores produced at 0.99 a_w and harvested using an aqueous solution. “Food plant”: spores produced at 0.95 a_w and dry-harvested. Shaded cells were used to compare the physiological states. *: inactivation of “laboratory” spores significantly different from that of “food plant” ones at p=0.05.