



HAL
open science

Influence of age on the spread of SARS-CoV-2 variant B.1.1.7

Chloé Dimeglio, Florence Nicot, Marcel Miedougé, Jean-Loup Chappert,
Cécile Donnadiou, Jacques Izopet

► **To cite this version:**

Chloé Dimeglio, Florence Nicot, Marcel Miedougé, Jean-Loup Chappert, Cécile Donnadiou, et al..
Influence of age on the spread of SARS-CoV-2 variant B.1.1.7. *Journal of Clinical Virology*, 2021, 141,
10.1016/j.jcv.2021.104872 . hal-03739196

HAL Id: hal-03739196

<https://hal.inrae.fr/hal-03739196>

Submitted on 13 Jun 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial | 4.0 International License

1 **Influence of age on the spread of SARS-CoV-2 variant B.1.1.7**

2 Chloé Dimeglio^{1,2*}, Florence Nicot¹, Marcel Miedougé¹, Jean-Loup Chappert³, Cécile Donnadieu⁴,
3 Jacques Izopet^{1,2}

4
5 ¹ CHU Toulouse, Hôpital Purpan, Virology Laboratory, 31300 Toulouse, France

6 ²INSERM UMR1291 – CNRS UMR5051, Toulouse Institute for Infectious and Inflammatory Diseases
7 (INFINITy), 31300 Toulouse, France

8 ³Santé Publique France, Cellule Occitanie, Direction des regions, Montpellier 34067, France

9 ⁴ Genotoul - Genome & Transcriptome - Plateforme Génomique (GeT-PlaGe), US INRAe 1426,
10 Castanet-Tolosan 31326, France

11

12 *To whom correspondence should be addressed: 330 avenue de Grande Bretagne, 31059 Toulouse
13 mail: dimeglio.c@chu-toulouse.fr

14 Tel : +33567690457

15

16 **Keywords:** SARS-CoV-2; variant B.1.1.7; age; public health

17

18 Dear Editor,

19 Previous studies of COVID-19 have indicated that children are less likely to be infected than
20 adults [1,2]. The rate at which the SARS-CoV-2 virus spreads may depend on the age of the
21 people infected. A new SARS-CoV-2 variant, B.1.1.7, emerged in the south-east of England in
22 October 2020 (3) and has subsequently spread to several countries including France. Recent
23 reports estimate that this new variant is up to 70% more transmissible than the original virus
24 and is responsible for an increase in the number of cases (4,5). We examined the

25 relationship between the circulation of the SARS-CoV-2 variant B.1.1.7 and the age of those
26 infected.

27 Nasopharyngeal swab samples were collected at drive-through testing centers and private
28 laboratories in the Toulouse area between January 8 and March 21, 2021. Samples were
29 tested using the ThermoFisher TaqPath RT-PCR assay (Scientific TaqPath COVID-19 Combo
30 Kit, Thermo Fisher, Waltham, USA). All positive samples with a cycle threshold (Ct) below 28
31 (N-gene) were re-tested using the Illumina CovidSeq test (COVIDSeq, Illumina, USA) on the
32 Illumina Nova Seq 6000 Sequencing System (Genotoul platform, GeTPlaGe, Toulouse). SARS-
33 CoV-2 sequences were analysed using the Illumina DRAGEN COVIDSeq test pipeline.

34 A total of 2133 positive samples were analyzed. The subjects' ages ranged from 0.25 years to
35 99 years (mean: 34.6 ± 19.09). The main variants in the Toulouse area on January 8, 2021
36 were B.1.177 (GV/20E.EU1), 42.1% of the total, and B.1.160 (GH/20A.EU2), 17.1%. The
37 SARS-CoV-2 variant B.1.1.7 was detected the following week (January 11-17) and became
38 the majority strain (over 70% of positive tests) during the first week of February (Figure 1A).
39 We compared the distribution of the B.1.1.7 variant to those of the other variants in young
40 (≤ 20 years) and older subjects (> 20 years). The proportion of the B.1.1.7 variant was
41 significantly greater in the young subjects in the period January 8 to March 21 (Figure 1B,
42 $p=0.04$, Wilcoxon's signed rank test). The greatest proportion of the B.1.1.7 variant was
43 found in the 10-20 year old subjects (data not shown).

44 The B.1.1.7 variant therefore seems to spread more rapidly in young people (< 20) than in
45 older people, unlike the classic forms of SARS-CoV-2 (6). This may be due to young people
46 adhering less rigorously to barrier measures (7). These findings could be relevant to the
47 design of public health measures intended to contain the spread of SARS-CoV-2.

48

49 **Conflict of Interest**

50 The authors declare no conflict of interest

51 **Funding/Support**

52 This work received no funding.

53 **Ethical Approval**

54 N/A

55

56 **Acknowledgments**

57 The English text was edited by Dr Owen Parkes.

58 We thank Charlène Chartron, Cécile Sergent, Pascal Chourrout, Patrick Caneiro, Nathalie
59 Wilhelm, Anne Bertrou, Frédéric Février, Brigitte Rivière, Marie Delpech, Catherine Pottier,
60 Philippe Gueudet, Alain Le Coustumier, Joseph Watine, Samir Brechiche, Elodie Carrere and
61 Margot Salse, Isabelle Le Goff, Audrey Brignoli and Roxane Steux, Anne Bruno, Pascal
62 Coudène, and Jean-Louis Galinier for their highly valuable implication.

63

64 **REFERENCES**

65

- 66 1. Nisha S Mehta, Oliver T Mytton, Edward W S Mullins, et al. SARS-CoV-2 (COVID-19): What Do
67 We Know About Children? A Systematic Review, *Clinical Infectious Diseases*, Volume 71,
68 Issue 9, 1 November 2020, Pages 2469–2479, <https://doi.org/10.1093/cid/ciaa556>
- 69 2. Statista Research Department. Population distribution in China in 2019, by broad age group.
70 Statista Research Department, March 5, 2020. Available at:
71 <https://www.statista.com/statistics/251524/population-distribution-by-age-group-in-china/>.
72 Accessed 23 March 2021.
- 73 3. Volz E. et al. Transmission of SARS-CoV-2 Lineage B.1.1.7 in England: Insights from linking
74 epidemiological and genetic data. *medRxiv* (2021)
- 75 4. Kirby, T. New variant of SARS-CoV-2 in UK causes surge of COVID-19. *Lancet Respiratory*
76 *Medicine*. 2020.DOI:[https://doi.org/10.1016/S2213-2600\(21\)00005-9](https://doi.org/10.1016/S2213-2600(21)00005-9)
- 77 5. Leung K, Shum MH, Leung GM, Lam TT, Wu JT. Early transmissibility assessment of the N501Y
78 mutant strains of SARS-CoV-2 in the United Kingdom, October to November 2020. *Euro*
79 *Surveill.* 2021 Jan;26(1):2002106. doi: 10.2807/1560-7917.ES.2020.26.1.2002106.

- 80 6. Lee P-I Hu Y-L Chen P-Y Huang Y-C Hsueh P-R. Are children less susceptible to COVID-19? J
81 Microbiol Immunol Infect. 2020; (published online Feb
82 25.)<https://doi.org/10.1016/j.jmii.2020.02.011>
83 7. Nivette A, Ribeaud D, Murray A, et al. Non-compliance with COVID-19-related public health
84 measures among young adults in Switzerland: Insights from a longitudinal cohort study. Soc
85 Sci Med. 2021;268:113370. doi:10.1016/j.socscimed.2020.113370
86

87

88

89 **Figure legend**

90 Figure 1: Proportion of circulating SARS-CoV-2 variant B.1.1.7 among all positive samples (1A) and among positive
91 samples by age category (1B) in the Toulouse area

92 Week 1: 4-10 January, 2021, Week 2: 11-17 January, 2021, Week 3: 18-24 January, 2021, Week 4: 25-31 January,
93 Week 5: 1-7 February, 2021, Week 6: 8-14 February, 2021, Week 7: 15-21 February, 2021, Week 8: 22-28
94 February, 2021, Week 9: 1-7 March, 2021, Week 10: 8-14 March, 2021, Week 11: 15-21 March, 2021

Figure 1

