



HAL
open science

Solving the grand challenge of phenotypic integration: allometry across scales

François Vasseur, Adrianus Johannes Westgeest, Denis Vile, Cyrille Violle

► To cite this version:

François Vasseur, Adrianus Johannes Westgeest, Denis Vile, Cyrille Violle. Solving the grand challenge of phenotypic integration: allometry across scales. *Genetica*, 2022, 150 (3-4), pp.161-169. 10.1007/s10709-022-00158-6 . hal-03745940

HAL Id: hal-03745940

<https://hal.inrae.fr/hal-03745940>

Submitted on 4 Aug 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License



Solving the grand challenge of phenotypic integration: allometry across scales

François Vasseur¹ · Adrianus Johannes Westgeest¹ · Denis Vile² · Cyrille Violle¹

Received: 3 December 2021 / Accepted: 13 June 2022
© The Author(s) 2022

Abstract

Phenotypic integration is a concept related to the cascade of trait relationships from the lowest organizational levels, i.e. genes, to the highest, i.e. whole-organism traits. However, the cause-and-effect linkages between traits are notoriously difficult to determine. In particular, we still lack a mathematical framework to model the relationships involved in the integration of phenotypic traits. Here, we argue that allometric models developed in ecology offer testable mathematical equations of trait relationships across scales. We first show that allometric relationships are pervasive in biology at different organizational scales and in different taxa. We then present mechanistic models that explain the origin of allometric relationships. In addition, we emphasized that recent studies showed that natural variation does exist for allometric parameters, suggesting a role for genetic variability, selection and evolution. Consequently, we advocate that it is time to examine the genetic determinism of allometries, as well as to question in more detail the role of genome size in subsequent scaling relationships. More broadly, a possible—but so far neglected—solution to understand phenotypic integration is to examine allometric relationships at different organizational levels (cell, tissue, organ, organism) and in contrasted species.

Keywords Allometry · Phenotypic integration · Genome size · Metabolic scaling · Trait relationships

Building complex organisms: an allometric perspective

The development of large and complex multicellular organisms requires the coordination and integration of a multitude of processes at different organizational levels (Pigliucci 2007; Zinner et al. 2020). Phenotypic integration (i.e. how complex organisms are built and how phenotypic traits are connected and coordinated) has been examined from a wide variety of approaches and research fields, including physiology, evolutionary biology, ecology and allometry (see for instance Schlichting and Pigliucci 1995; Pigliucci et al. 1996; Klingenberg 1996; Murren 2002; Pigliucci 2003;

Klingenberg and Marugan-Lobon 2013; Klingenberg 2014; Klingenberg 2016; Watanabe 2022). However, joint empirical efforts at the interface of those disciplines warrant further study. One reason is that the study of trade-offs and strategies in comparative ecology has largely occurred independently of the study of phenotypic integration in evolutionary biology. Indeed, comparative ecology has mainly focused on global trends and trait-trait relationships across species, taking advantage of large interspecific databases (Brosse et al. 2021; Kattge et al. 2011; Díaz et al. 2016; Mouillot et al. 2021; Toussaint et al. 2021); while evolutionary biology has mainly focused on investigating the mechanisms and consequences of genetic correlations within species (Steppan et al. 2002; Arnold et al. 2008; Penna et al. 2017; Milocco and Salazar-Ciudad 2022).

Following Arnold's definition (Arnold 1983), phenotypic integration can be represented as a pyramid, in which traits related to gene products (RNA, enzymes) are at the bottom, and performance-related traits (growth rate, fecundity) at the top (Fig. 1). Ecological theory proposes a similar hierarchy of trait integration from genes to fitness (Violle et al. 2007). Basically, a trait Y is said to be integrated when it results from the effect of one or several traits X_i (considered


This article is part of the Special Issue “The relationship between genotype and phenotype: new insights on an old question”.

✉ François Vasseur
francois.vasseur@cefe.cnrs.fr

¹ CEFE, University Montpellier, CNRS, EPHE, IRD, Montpellier, France

² LEPSE, University Montpellier, INRAE, Institut Agro, Montpellier, France

Fig. 1 Arnold's view of phenotypic integration. The pyramidal cascade of trait relationships from genes to organism's fitness is represented. Here, the trait-trait relationships between two successive organizational levels are hypothetical



at “lower degrees of phenotypic integration”). Traits at different organizational levels are assumed to be connected by scaling relationships or functions (Fig. 1). For instance, gene expression impacts RNA content, which in turn impacts cell metabolism and size, which in turn impacts organ physiology and size, and so on. Following Arnold's view of phenotypic integration, a genetic variation impacting a trait should also impact the traits that are at higher levels and that result from the effect of the focal trait, with possible feedback effects between consecutive levels of integration. This is corroborated by recent findings, in which the rate of adaptive substitution (i.e. genetic variation that impacts organismal fitness) along the genome increases with the degree of phenotypic integration (Zhang 2018). However, the pyramidal view of phenotypic integration proposed by Arnold largely remains theoretical: the ‘cause-and-effect’ cascades, as well as the relationships involved, are notoriously difficult to establish, so is the hierarchy of phenotypic integration. Traits are generally organized into groups of interacting features, called modules, which are relatively independent from each others (e.g. floral traits versus vegetative traits in plants) (Wagner et al. 2007; Klingenberg 2008; Murren 2012; Diggle 2014). However, how modules and traits at different scales are connected to each other remains unclear.

One of the main consequences of a change in the size of an object is the modification of its surface area to volume ratio. This is explained by simple geometric considerations: the surface area, which allows exchanges of resources and energy between the internal medium and the environment, increases according to a square function; while the volume, a priori linked to the number of cells and thus to energy needs,

increases according to a cubic function. As a result, the surface-to-volume ratio decreases with volume and mass, whatever the shape of the object (but see Bestova et al. 2021). In living things, these geometric constraints have profound consequences for resource use strategies. First, the rate of metabolic activity—measured through basal respiration rate—is expected to increase non-proportionally with organism size. Having a smaller surface-to-volume ratio, large individuals are expected to have a lower metabolic rate (Kleiber 1947) and to lose less heat per unit biomass (Bergmann's rule) (Bergmann 1847; Blackburn et al. 1999). For example, an increase of only 10% in height and width of a cone representing a simplified tree would result in an increase of c. 20% in its surface area, and an increase of c. 33% in its volume. A direct consequence is that surface-related processes, such as carbon fixation and water loss, increase twice as fast as the increase in height and width. A tree with increased size will therefore be likely to use significantly more surface-related resources (e.g., transpiration of water, acquisition of carbon through photosynthesis). Moreover, volume-related processes, such as carbon consumption via cellular respiration, increase even more significantly (33%). This example is of course an unrealistic simplification (a tree is never as simple as a cone) of the physiological processes involved in complex organisms, but it illustrates the major, and asymmetric, influence of size variations on resource use, growth, and development (Ohlberger 2013; Malerba et al. 2017; Lindmark et al. 2019; Malerba and Marshall 2019).

Considering that the size of an object is the main determinant of its internal properties is the central tenet of allometry research, which, in its broad definition, is the

study of the relationships between the size of a biological entity (e.g., genome, cell, organ, organism, population) and its morphological, physiological and metabolic traits (Niklas 1994). Allometric equations aim at modelling trait-trait relationships across organizational scales. They typically take the form of power-law functions $Y = \alpha X^\beta$, where X is a size-related trait and Y a morphological or physiological trait (Niklas 1994). Strikingly, the same equations are used to scale up from genome size to cell size (Kozłowski et al. 2003b; Beaulieu et al. 2008; Šímová and Herben 2011; Gonzalez-de-Salceda and Garcia-Pichel 2021), cell size to organ size (Gregory et al. 2000; Tisné et al. 2008; John et al. 2013), organ size to organism size (Stahl 1965; Gould 1971; Stevenson et al. 1995; Lindstedt and Schaeffer 2002; Shingleton et al. 2007; Poorter et al. 2015), body size to energy consumption (Huxley 1932; Kleiber 1932, 1947; DeLong et al. 2010; Capellini et al. 2010), as well as from organism size to population density (Enquist et al. 1998; Malerba and Marshall 2019) (see Fig. 2 for examples of allometric relationships at different organizational levels). We advocate that a promising approach to understand phenotypic integration is to study scaling relationships, because modelling such relationships is built from a mathematical toolkit aimed at linking one level of integration to another.

Integrating traits across scales: biological regularity versus genetic diversity

The allometric equations used to connect one phenotypic level to another are generally linearized by log-transformation, and β becomes the slope of the relationships $\log(Y) = \log(\alpha) + \beta \log(M)$ (Fig. 2). Simple geometric models predict $\beta = 2/3$ because β must be determined by the ratios between surface area (square function) and volume (cubic function) (Rubner 1883; White and Seymour 2003). However, in 1932, Kleiber observed that energy consumption in animals—which spanned several orders of magnitude in size—scales with body mass with an allometric exponent $\beta = 3/4$ (Kleiber 1932). This observation yielded many studies on the value that β actually takes for different traits in different taxa. Moreover, the question of a biological constant that applies to virtually all organisms has been the subject of intense debates in the scientific literature. Indeed, the sole existence of a biological constant would suggest that organisms are built under the same developmental constraints, which translates into universal scaling relationships. Moreover, if the same constants are observed at different levels of phenotypic integration (from genomes to whole organisms), this would have strong implications for our understanding, and the potential prediction, of biological structures and organization.

The introduction of fractal geometries into mechanistic models has revolutionized the field of allometry research. In the late 1990s and early 2000s, a series of papers laid the theoretical foundations for the Metabolic Scaling Theory (MST) (Enquist et al. 1999, 2007b; West et al. 1999a, b, 2001, 2002; Niklas and Enquist 2001; Brown et al. 2004; Price and Enquist 2007), which invokes the fractal-like geometry of vascular networks as the main determinant of several scaling relationships in both plants and animals. MST is primarily based on two central hypotheses: the network is space-filling and the branching is area-preserving, i.e. the sum of cross-sectional areas of daughter branches equals the cross-sectional area of the mother branch. These hypotheses led to predictions concerning the variation of organism morphology, the allocation of biomass and the physiological rates at the organism level (Fig. 2) (Savage et al. 2008). For instance, the organism's basal metabolic rate is expected to scale with $\beta = 3/4$ to the organism's body mass due to the optimization of the internal transport distance for resources and energy (West et al. 1999b). In plants, the metabolic rate is expected to scale isometrically (i.e. with exponent $\beta = 1$, which directly translates strict proportionality) with total leaf area and total leaf biomass, and allometrically (with exponent $\beta = 3/4$) with the mass of the plant (Enquist et al. 1998; West et al. 1999a; Price et al. 2007). These scaling expectations represent the master equations of the MST, from which other scaling relationships at different levels of phenotypic integration have been derived, such as metabolic activity, relative growth rate, xylem fluxes and net photosynthetic rate in plants (Enquist 2002; Price and Enquist 2007; Enquist et al. 2007b; Savage et al. 2008). MST predictions even extend to population dynamics and demography, as illustrated by the expected exponent of $-3/4$ for the relationship between maximum plant density and plant size (Enquist et al. 1998). However, some of the mechanistic assumptions of MST remain controversial, in large part because the universality of the scaling exponent has been challenged theoretically and empirically (Dodds et al. 2001; Kozłowski and Konarzewski 2004; Glazier 2005; Coomes 2006; Russo et al. 2007).

The theoretical underpinnings of MST are based on the idea that natural selection has optimized vascular networks in order to minimize the energy required for resource distribution, which would shape quasi-universal allometric relationships within and between species (West et al. 1999b). Interspecific comparisons have revealed global patterns of covariation that generally confirm the predictions of MST, in both animals and plants (Fig. 2). Moreover, recent intraspecific studies in plants also demonstrated an allometric exponent of $3/4$ in wild tomatoes and in *A. thaliana* (Vasseur et al. 2012; Muir and Thomas-Huebner 2013, 2015). This suggests that allometric relationships result from strong




Fig. 2 Examples of allometric relationships between scales. **A** Relationship between leaf mass (M_L) and stem mass (M_S) (Price et al. 2010). **B** Relationship between cell area (A_c) and genome size (S) (Beaulieu et al. 2008). **C** Relationship between plant mass (M) and plant density (N) (Enquist et al. 1998). **D** Relationship between organism's basal metabolic rate (B) and body mass (M) (West et al. 2002).

E Relationship between the number of 16S/18S ribosomal genes per cell (R_c) and cell volume (V_c) (Gonzalez-de-Salceda and Garcia-Pichel 2021). **F** Relationship between xylem flux (F) and plant mass (M) (Enquist and Niklas 2001). **G** Relationship between flatworm body volume (V) and genome size (S) (Gregory et al. 2000)

constraints, which operate at different organizational levels independently from the taxonomic scale and which, on average, adhere to the predictions of MST. Yet, intraspecific variability in allometries are also frequently observed in animals (Kozłowski et al. 2003a; Sieg et al. 2009; Summers and Ord 2021), and in plants (Vasseur et al. 2012; Muir and Thomas-Huebner 2013, 2015). For instance, the rate of respiratory metabolism has been found to scale isometrically (*i.e.* with

exponent $\beta \approx 1$) with plant biomass in herbaceous plants and tree seedlings (Reich et al. 2006; Cheng et al. 2010). Other studies in plants (Enquist et al. 2007a; Mori et al. 2010) and mammals (Kolokotronis et al. 2010), showed that respiration rate scales differently for organisms with small body mass and large body mass. In plants, competition for light results in significant variations in allometric relationships in trees (Russo et al. 2007; Lines et al. 2012). Strikingly, the

studies reporting that intraspecific plant allometries follow the $\frac{3}{4}$ rule on average, also showed that the allometric exponent can vary significantly around $\frac{3}{4}$ because of evolutionary adaptation to stressful environments (Vasseur et al. 2018b). Studies in plants also indicated that variation in allometric exponents are genetically associated with different carbon and water-use strategies (Vasseur et al. 2012, 2014; Muir and Thomas-Huebner 2015), as well as reproductive output and resistance to stress (Muir and Thomas-Huebner 2013; Vasseur et al. 2018b). Together, these recent findings support the idea that scaling relationships vary between populations as a consequence of evolutionary adaptation to different environments, but they generally center around $\frac{3}{4}$. Overall, it suggests that phenotypic integration obeys fundamental laws that are translated into the relative constancy of allometric relationships at different organizational levels. However, a key question related to the integration of phenotypic traits is how adaptation to contrasting environments is the result of, or is dependent on, substantial variations in allometric relationships (Pigliucci et al. 1996; Pélabon et al. 2014).

Current prospects in allometry: toward a genetic synthesis?

Exploring the genetic determinism of intraspecific allometries represents a promising avenue to better understand how organisms are built. Recent efforts have been devoted to the identification and characterization of genetic diversity in allometric relationships (Long et al. 2006; Qin et al. 2012, 2013; Vasseur et al. 2012, 2018b; Muir and Thomas-Huebner 2015). These works suggest that selection must operate on the genes that impact allometric parameters such as β , because the same biomechanical constraints (notably those related to surface area-to-volume ratio) are expected to operate at different scales. Identifying those genes should be a research priority to determine which molecular pathways are involved in the integration of phenotypic traits with size. In addition, the genetic architecture of intraspecific variability in allometry is directly informative about how organisms are expected to evolve. The predominance of few pleiotropic genes with strong effects on allometries, for instance genes related to life history, would suggest evolution by abrupt increments and sharp variations (Paaby and Rockman 2013). By contrast, if allometries are determined by a multitude of genes along the genomes, each having only a small effect, then a more gradual and continuous evolution of organisms is expected, which is directly related to the integration of multiple traits in complex organisms, and which has been called the “cost of complexity” (Orr 2000; Wagner et al. 2008). Genome-wide association studies (GWAS) might help to clarify the oligo- and/or the polygenic nature of allometric variations by the screening of plant and animal

populations at different organizational levels. Furthermore, the recent rise of next-generation sequencing technologies allows extensive genomic analyses of the identified genes. The genetic survey of natural populations through space and time can reveal genomic signatures of selection and clarify the demographic history of the investigated populations (Weigel 2012). For instance, comparing genetic and phenotypic differentiation between populations through Q_{ST}/F_{ST} approaches (Leinonen et al. 2013) allows testing whether allometric parameters such as β vary between populations due to neutral differentiation such as genetic drift, or because of directional or balancing selection. This kind of approach coupling allometry with quantitative genetics and genomics has recently been applied to the model plant *Arabidopsis thaliana*. It first identified two major pleiotropic genes as responsible for allometric variation between plant growth rate and plant dry mass (Vasseur et al. 2012), suggesting that plants evolve through strong allometric increments. However, this study was performed on a population of recombinant inbred lines (RIL) artificially created by crossing two contrasted ecotypes. A similar study performed six years later on a collection of 450 ecotypes (Vasseur et al. 2018b) revealed by contrast that plant allometry has a more polygenic architecture, with many contributing genes and with the major gene identified affecting only around 1% of the phenotypic variance. Taking advantage of the extensive genetic information available for *A. thaliana*, this study also showed that genes related to allometric variations exhibited signatures of selection and they were clustered geographically along a latitudinal gradient. Despite these encouraging findings, our knowledge of the genetic determinism of organism allometries remains limited. Yet, genetic studies of allometric variations could shed light on the complex genotype to-phenotype relationships involved in the coordination of complex phenotypes. A more systematic analysis of intraspecific relationships at different levels of integration (genome, cell, organ, and whole organism) is needed, in different species and different environmental conditions. Moreover, testing the effect of natural or artificial selection in real time with experimental evolution is a promising avenue to examine the evolution of allometry and its role for organism adaptation to contrasted environments.

Allometric scaling from genome size to whole organisms: the grand challenge

An alternative approach to explore phenotypic integration through the lens of allometry is to link scaling relationships at different organizational levels (i.e. from genome to cell, organ and whole organism). Each level has its own set of allometric expectations, with its own theoretical corpus. For instance, there are allometric expectations for the allometry

of genome size versus cell size (Šímová and Herben 2011), and cell size versus organism size (Gregory et al. 2000), as well as organism size versus metabolic rate or growth rate (Brown et al. 2004; Savage et al. 2008). Connecting these expectations, made at different levels and from different biological considerations, around a unifying model represents the ultimate challenge in allometry research. However, and as said above, the questions surrounding the links of cause-and-effects between relationships and organizational levels are particularly difficult to address. If in theory we could build allometric relationships to directly connect the lower level of integration—genome size—to the upper—organism size and metabolism—, in practice allometric relationships are generally strong between two successive organizational levels—e.g., genome size versus cell size, organism size versus organ size—but they decrease in predictive power with increasing phenotypic scale (Knight and Beaulieu 2008).

We advocate here that a promising avenue to solve this issue relies on the measurement of genome size. Genome size is usually estimated by DNA content in pg, which is expected to vary a lot among species because of different sources of variability in genome composition (including the number of genes but above all the amount of non-coding DNA, transposable elements, structural variations, repeated sequences, etc.) (Schubert and Vu 2016). However, this measurement of genome size misses the key point: what really matters is not the total amount of DNA, but the number of different genes that are effectively expressed during organism growth and development, as well as the amount (and repetition) of DNA segments regulating gene transcription and translation (mostly ribosomal DNA). The amount of DNA and the level of gene expression might differ a lot due to different processes. First, large portions of the genome exhibit presence/absence polymorphisms between genotypes of the same species (Tan et al. 2012; Bush et al. 2013; González et al. 2013; Zhang et al. 2014; Jiang et al. 2015), which suggests strong variability in the number of expressed genes between these genotypes. To what extent this variability is related to phenotypic variation remains unknown so far (Weisweiler et al. 2019). Second, the level of heterozygosity could positively impact the number of expressed genes and/or alleles. This could partly explain the increase in organism vigour and size frequently reported in hybrids between homozygous lines (Charlesworth and Charlesworth 1999). Indeed, the number of heterozygous genes, as well as presence/absence alleles has been shown to be positively correlated to hybrid vigour in plants (Springer et al. 2009; Paschold et al. 2012; Hochholdinger and Baldauf 2018). The links between expressed genome size and plant vigour are also supported by the higher size of organs and whole organisms in polyploid lines (containing 4 N, 6 N, 8 N etc.) versus their diploid relatives (Chen 2010; Sattler et al. 2016). Consistent with allometric expectations, polyploids might exhibit

larger organs and body size due to their enlarged genomes (Otto 2007; Lavania 2016). Accordingly, endopolyploidy at the cellular level is also associated with increased cell size in both plants and animals (Leitch and Dodsworth 2001; Neiman et al. 2017). However, polyploidy could also uncouple trait covariations and, thus, alter the pattern of phenotypic integration observed in diploids. This is notably the case for the traits related to leaf physiology and resource use (López-Jurado et al. 2022). Yet, to what extent polyploidy might break allometric rules and alter scaling equations at different organization scales remains to be elucidated. Finally, epigenetic marks along the genome can repress the expression of a large number of genes. For instance, methylated regions, which have reduced RNA transcription, represent a significant portion of the genome (Seymour et al. 2014; Seymour and Becker 2017). Consistent with a link between genome methylation level and organism growth and size, the re-analysis of the amount of methylated regions measured on more than 1000 natural accessions of *A. thaliana* (Kawakatsu et al. 2016) with the growth-related traits measured on 409 accessions in common (Vasseur et al. 2018a) (but on different individuals cultivated in an independent experiment), revealed a negative correlation between individual growth rate and the fraction of methylated genome ($r = -0.044$, slope = -0.02 , $P = 0.002$; unpublished data), which suggests that larger expressed genomes are associated with bigger plants and higher absolute growth rate. The potential allometric linkage between expressed genome size and growth rate could also rely on stoichiometric considerations. Indeed, individuals with large genomes are expected to produce more RNA (Kozłowski et al. 2003b), which is phosphorus consuming and could explain the positive relationship between growth rate and RNA/protein as well as phosphorus/protein ratios observed empirically (Elser et al. 2003). Consistently, a strong relationship has been observed between the number of ribosomal genes per cell and cell size across a large diversity of prokaryote and eukaryote organisms (Gonzalez-de-Salceda and Garcia-Pichel 2021). Collectively, these findings suggest that the number of expressed genes in the genome could be the key piece to link allometric relationships at different levels of phenotypic integration, as well as to reconcile allometry with other disciplines such as biological stoichiometry, quantitative genetics and physiology.

Conclusion

How multiple traits at different organizational levels are phenotypically integrated to ensure the proper development of complex organisms in a coordinated way is a key question for modern biology. We advocate here that allometric

relationships offer a promising avenue to tackle this issue because (i) they rely on well-documented empirical observations, (ii) their mechanistic bases have been studied at different scales, and (iii) they propose a mathematical toolkit to infer hypotheses and make predictions. We argue that future directions for allometry research should focus in priority on the genetic determinism of scaling variation, taking advantage of the development of next-generation sequencing technologies, as well as on how the expressed genome size scales with higher orders of phenotypic integration, from cell to whole organism and metabolism. We hope that this article will encourage novel experiments and interdisciplinary investigations to address allometric questions at different levels of phenotypic integration.

Acknowledgements This work was supported by CNRS, INRAE, the French Agency for Research (ANR grant ANR-17-CE02-0018-01, ‘AraBreed’ to FV, DV and CV), and the European Research Council (ERC) (‘CONSTRAINTS’: grant ERCStG-2014-639706 to CV; ‘PHENOVI GOUR’: grant ERC-StG-2020-949843 to FV and AJW).

Author contributions All authors equally contributed to the writing of the paper.

Declarations

Conflict of interest The authors declare no conflict of interest.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

References

- Arnold SJ (1983) Morphology, performance and fitness. *Integr Comp Biol* 23:347–361
- Arnold SJ, Bürger R, Hohenlohe PA, Ajie BC, Jones AG (2008) Understanding the evolution and stability of the G-matrix. *Evolution* 62(10):2451–2461
- Beaulieu JM, Leitch IJ, Patel S et al (2008) Genome size is a strong predictor of cell size and stomatal density in angiosperms. *New Phytol* 179:975–986
- Bergmann C (1847) Über die Verhältnisse der Wärme ökonomie der Thiere zu ihrer Grösse. *Göttinger Studien* 3:595–708
- Bestová H, Segrestin J, von Schwartzberg K et al (2021) Biological scaling in green algae: the role of cell size and geometry. *Sci Rep* 11(1):1–9
- Blackburn TM, Gaston KJ, Loder N (1999) Geographic gradients in body size: a clarification of Bergmann's rule. *Divers Distrib* 5:165–174
- Brosse S, Charpin N, Su G, Toussaint A, Herrera-R GA, Tedesco PA, Villéger S (2021) FISHMORPH: a global database on morphological traits of freshwater fishes. *Global Ecol Biogeogr* 30(12):2330–2336
- Brown JH, Gillooly JF, Allen AP et al (2004) Toward a metabolic theory of ecology. *Ecology* 85:1771–1789
- Bush SJ, Castillo-Morales A, Tovar-Corona JM et al (2013) Presence-absence variation in *A. Thaliana* is primarily associated with genomic signatures consistent with relaxed selective constraints. *Mol Biol Evol* 31:59–69
- Capellini I, Venditti C, Barton RA (2010) Phylogeny and metabolic scaling in mammals. *Ecology* 91:2783–2793
- Charlesworth B, Charlesworth D (1999) The genetic basis of inbreeding depression. *Genet Res* 74:329–340
- Chen ZJ (2010) Molecular mechanisms of polyploidy and hybrid vigor. *Trends Plant Sci* 15:57–71
- Cheng D-L, Li T, Zhong Q-L, Wang G-X (2010) Scaling relationship between tree respiration rates and biomass. *Biol Lett* 6:715–717
- Coomes DA (2006) Challenges to the generality of WBE theory. *Trends Ecol Evol* 21:593–596
- DeLong JP, Okie JG, Moses ME et al (2010) Shifts in metabolic scaling, production, and efficiency across major evolutionary transitions of life. *Proc Natl Acad Sci U S A* 107:12941–12945
- Díaz S, Kattge J, Cornelissen JH, Wright IJ, Lavorel S, Dray S, Reu B, Kleyer M, Wirth C, Colin Prentice I, Garnier E (2016) The global spectrum of plant form and function. *Nature* 529(7585):167–171
- Diggle PK (2014) Modularity and intra-floral integration in metameric organisms: plants are more than the sum of their parts. *Philos Trans Royal Soc B Biol Sci* 369(1649):20130253
- Dodds PS, Rothman DH, Weitz JS (2001) Re-examination of the “3/4-law” of Metabolism. *J Theor Biol* 209:9–27
- Elser JJ, Acharya K, Kyle M et al (2003) Growth rate–stoichiometry couplings in diverse biota. *Ecol Lett* 6:936–943
- Enquist BJ (2002) Universal scaling in tree and vascular plant allometry: toward a general quantitative theory linking plant form and function from cells to ecosystems. *Tree Physiol* 22:1045–1064
- Enquist BJ, Niklas KJ (2001) Invariant scaling relations across tree-dominated communities. *Nature* 410:655–660
- Enquist BJ, Brown JH, West GB (1998) Allometric scaling of plant energetics and population density. *Nature* 395:163–165
- Enquist BJ, West GB, Charnov EL, Brown JH (1999) Allometric scaling of production and life-history variation in vascular plants. *Nature* 401:907–911
- Enquist BJ, Allen AP, Brown JH et al (2007a) Biological scaling: does the exception prove the rule? *Nature* 445:E9–10
- Enquist BJ, Kerkhoff AJ, Stark SC et al (2007b) A general integrative model for scaling plant growth, carbon flux, and functional trait spectra. *Nature* 449:218–222
- Glazier DS (2005) Beyond the “3/4-power law”: variation in the intra- and interspecific scaling of metabolic rate in animals. *Biol Rev* 80:611–662
- González VM, Aventín N, Centeno E, Puigdomènech P (2013) High presence/absence gene variability in defense-related gene clusters of *Cucumis melo*. *BMC Genomics* 14:782
- Gonzalez-de-Salceda L, Garcia-Pichel F (2021) The allometry of cellular DNA and ribosomal gene content among microbes and its use for the assessment of microbiome community structure. *Microbiome* 9:173
- Gould SJ (1971) Geometric similarity in allometric growth: a contribution to the problem of scaling in the evolution of size. *Am Nat* 105:113–136

- Gregory TR, Hebert PD, Kolasa J (2000) Evolutionary implications of the relationship between genome size and body size in flatworms and copepods. *Heredity* 84(Pt 2):201–208
- Hochholdinger F, Baldauf JA (2018) Heterosis in plants. *Curr Biol* 28:R1089–R1092
- Huxley JS (1932) *Problems of relative growth* Methuen & Co. LTD, London
- Jiang L, Lv Y, Li T et al (2015) Identification and characterization of presence/absence variation in maize genotype Mo17. *Genes Genomics* 37:503–515
- John GP, Scoffoni C, Sack L (2013) Allometry of cells and tissues within leaves. *Am J Bot* 100:1936–1948
- Kattge J, Diaz S, Lavorel S, Prentice IC, Leadley P, Böniš G, Garnier E, Westoby M, Reich PB, Wright IJ, Cornelissen JHC (2011) TRY—a global database of plant traits. *Glob Chang Biol* 17(9):2905–2935
- Kawakatsu T, Huang S-SC, Jupe F et al (2016) Epigenomic diversity in a global collection of *Arabidopsis thaliana* accessions. *Cell* 166:492–505
- Kleiber M (1932) Body size and metabolism. *Energy* 1:E9
- Kleiber M (1947) Body size and metabolic rate. *Physiol Rev* 27:511–541
- Klingenberg CP (1996) Multivariate allometry. In *Advances in morphometrics*, Springer, Boston, MA
- Klingenberg CP (2008) Morphological integration and developmental modularity. *Annual Rev Ecol Evol Syst* 115–132
- Klingenberg CP (2014) Studying morphological integration and modularity at multiple levels: concepts and analysis. *Phil Trans R Soc B: Biol Sci* 369(1649):20130249
- Klingenberg CP (2016) Size, shape, and form: concepts of allometry in geometric morphometrics. *Dev Genes Evol* 226(3):113–137
- Klingenberg CP, Marugán-Lobón J (2013) Evolutionary covariation in geometric morphometric data: analyzing integration, modularity, and allometry in a phylogenetic context. *Syst Biol* 62(4):591–610
- Knight CA, Beaulieu JM (2008) Genome size scaling through phenotype space. *Ann Bot* 101:759–766
- Kolokotronis T, Savage V, Deeds EJ, Fontana W (2010) Curvature in metabolic scaling. *Nature* 464:753–756
- Kozłowski J, Konarzewski M (2004) Is West, Brown and Enquist's model of allometric scaling mathematically correct and biologically relevant? *Funct Ecol* 18:283–289
- Kozłowski J, Konarzewski M, Gawelczyk AT (2003b) Cell size as a link between noncoding DNA and metabolic rate scaling. *Proc Natl Acad Sci U S A* 100:14080–14085
- Kozłowski J, Konarzewski M, Gawelczyk AT (2003a) Intraspecific body size optimization produces interspecific allometries. *Macroecology: concepts and consequences* 299–320
- Leinonen T, Scott McCairns RJ, O'Hara RB, Merilä J (2013) QST–FST comparisons: evolutionary and ecological insights from genomic heterogeneity. *Nat Rev Genet* 14:179–190
- Leitch IJ, Dodsworth S (2001) Endopolyploidy in Plants. In: Wiley J, Ltd S (eds) eLS. John Wiley & Sons Ltd, Chichester, UK, pp 1–10
- Lindmark M, Ohlberger J, Huss M, Gårdmark A (2019) Size-based ecological interactions drive food web responses to climate warming. *Ecol Lett*. <https://doi.org/10.1111/ele.13235>
- Lindstedt SL, Schaeffer PJ (2002) Use of allometry in predicting anatomical and physiological parameters of mammals. *Lab Anim* 36:1–19
- Lines ER, Zavala MA, Purves DW, Coomes DA (2012) Predictable changes in aboveground allometry of trees along gradients of temperature, aridity and competition. *Glob Ecol Biogeogr* 21:1017–1028
- Long F, Chen YQ, Cheverud JM, Wu R (2006) Genetic mapping of allometric scaling laws. *Genet Res* 87:207–216
- López-Jurado J, Mateos-Naranjo E, Balao F (2022) Polyploidy promotes divergent evolution across the leaf economics spectrum and plant edaphic niche in the *Dianthus broteri* complex. *J Ecol* 110(3):605–618
- Malerba ME, Marshall DJ (2019) Size-abundance rules? Evolution changes scaling relationships between size, metabolism and demography. *Ecol Lett* 22:1407–1416
- Malerba ME, White CR, Marshall DJ (2017) Eco-energetic consequences of evolutionary shifts in body size. *Ecol Lett* 21:54–62
- Milocco L, Salazar-Ciudad I (2022) Evolution of the G matrix under nonlinear genotype–phenotype maps. *Am Nat* 199(3):420–435
- Mori S, Yamaji K, Ishida A et al (2010) Mixed-power scaling of whole-plant respiration from seedlings to giant trees. *Proc Natl Acad Sci U S A* 107:1447–1451
- Mouillot D, Loiseau N, Grenié M, Algar AC, Allegra M, Cadotte MW, Casajus N, Denelle P, Guéguen M, Maire A, Maitner B (2021) The dimensionality and structure of species trait spaces. *Ecol Lett* 24(9):1988–2009
- Muir CD, Thomas-Huebner M (2013) Small interspecific variation in metabolic scaling mediates tradeoff between drought tolerance and fecundity in wild tomatoes (*Solanum* sect. *Lycopersicon*; Solanaceae). *Ecophysiology And Genetics Of Phenotypic Diversification In Solanum* 61
- Muir CD, Thomas-Huebner M (2015) Constraint around quarter-power allometric scaling in wild tomatoes (*Solanum* sect. *Lycopersicon*; Solanaceae). *Am Nat* 186:421–433
- Murren CJ (2002) Phenotypic integration in plants. *Plant Species Biol* 17(2–3):89–99
- Murren CJ (2012) The integrated phenotype. *Integr Comp Biol* 52(1):64–76
- Neiman M, Beaton MJ, Hessen DO et al (2017) Endopolyploidy as a potential driver of animal ecology and evolution. *Biol Rev Camb Philos Soc* 92:234–247
- Niklas KJ, Enquist BJ (2001) Invariant scaling relationships for interspecific plant biomass production rates and body size. *Proc Natl Acad Sci U S A* 98:2922–2927
- Niklas KJ (1994) *Plant Allometry: The Scaling of Form and Process*. University of Chicago Press
- Ohlberger J (2013) Climate warming and ectotherm body size—from individual physiology to community ecology. *Funct Ecol* 27:991–1001
- Orr HA (2000) Adaptation and the cost of complexity. *Evolution* 54:13–20
- Paaby AB, Rockman MV (2013) The many faces of pleiotropy. *Trends Genet* 29:66–73
- Paschold A, Jia Y, Marcon C et al (2012) Complementation contributes to transcriptome complexity in maize (*Zea mays* L.) hybrids relative to their inbred parents. *Genome Res* 22:2445–2454
- Pélabon C, Firmat C, Bolstad GH, Voje KL, Houle D, Cassara J, Rouzic AL, Hansen TF (2014) Evolution of morphological allometry. *Ann N Y Acad Sci* 1320(1):58–75
- Penna A, Melo D, Bernardi S, Oyarzabal MI, Marroig G (2017) The evolution of phenotypic integration: How directional selection reshapes covariation in mice. *Evolution* 71(10):2370–2380
- Pigliucci M (2003) Phenotypic integration: studying the ecology and evolution of complex phenotypes. *Ecol Lett* 6(3):265–272
- Pigliucci M (2007) Finding the way in phenotypic space: the origin and maintenance of constraints on organismal form. *Ann Bot* 100(3):433–438
- Pigliucci M, Schlichting CD, Jones CS, Schwenk K (1996) Developmental reaction norms: the interactions among allometry, ontogeny and plasticity. *Plant Species Biol* 11(1):69–85
- Poorter H, Jagodzinski AM, Ruiz-Peinado R et al (2015) How does biomass distribution change with size and differ among species? An analysis for 1200 plant species from five continents. *New Phytol* 208:736–749

- Price CA, Enquist BJ (2007) Scaling mass and morphology in leaves: an extension of the WBE model. *Ecology* 88:1132–1141
- Price CA, Enquist BJ, Savage VM (2007) A general model for allometric covariation in botanical form and function. *Proc Natl Acad Sci U S A* 104:13204–13209
- Price CA, Gilooly JF, Allen AP et al (2010) The metabolic theory of ecology: prospects and challenges for plant biology. *New Phytol* 188:696–710
- Qin X, Niklas KJ, Qi L, Xiong Y, Li F (2012) The effects of domestication on the scaling of below-vs. aboveground biomass in four selected wheat (*Triticum*; Poaceae) genotypes. *Am J Botany* 99(6):1112–1117
- Qin XL, Weiner J, Qi L, Xiong YC, Li FM (2013) Allometric analysis of the effects of density on reproductive allocation and Harvest Index in 6 varieties of wheat (*Triticum*). *Field Crop Res* 144:162–166
- Reich PB, Tjoelker MG, Machado JL, Oleksyn J (2006) Universal scaling of respiratory metabolism, size and nitrogen in plants. *Nature* 439:457–461
- Rubner M (1883) Ueber den einfluss der korpergrösse auf stoffund kraftwechsel. *Z Biol* 19:535–562
- Russo SE, Wiser SK, Coomes DA (2007) Growth–size scaling relationships of woody plant species differ from predictions of the Metabolic Ecology Model. *Ecol Lett* 10:889–901
- Sattler MC, Carvalho CR, Clarindo WR (2016) The polyploidy and its key role in plant breeding. *Planta* 243:281–296
- Savage VM, Deeds EJ, Fontana W (2008) Sizing up allometric scaling theory. *PLoS Comput Biol* 4:e1000171
- Schlichting CD, Pigliucci M (1995) Gene regulation, quantitative genetics and the evolution of reaction norms. *Evol Ecol* 9(2):154–168
- Schubert I, Vu GT (2016) Genome stability and evolution: attempting a holistic view. *Trends Plant Sci* 21(9):749–757
- Seymour DK, Becker C (2017) The causes and consequences of DNA methylome variation in plants. *Curr Opin Plant Biol* 36:56–63
- Seymour DK, Koenig D, Hagmann J et al (2014) Evolution of DNA methylation patterns in the brassicaceae is driven by differences in genome organization. *PLoS Genet* 10:e1004785
- Shingleton AW, Frankino WA, Flatt T et al (2007) Size and shape: the developmental regulation of static allometry in insects. *BioEssays* 29:536–548
- Sieg AE, O'Connor MP, McNair JN et al (2009) Mammalian metabolic allometry: do intraspecific variation, phylogeny, and regression models matter? *Am Nat* 174:720–733
- Šímová I, Herben T (2011) Geometrical constraints in the scaling relationships between genome size, cell size and cell cycle length in herbaceous plants. *Proceedings of the Royal Society of London B: Biological Sciences* rspb20111284
- Springer NM, Ying K, Fu Y et al (2009) Maize inbreds exhibit high levels of copy number variation (CNV) and presence/absence variation (PAV) in genome content. *PLoS Genet* 5:e1000734
- Stahl WR (1965) Organ weights in primates and other mammals. *Science* 150:1039–1042
- Steppan SJ, Phillips PC, Houle D (2002) Comparative quantitative genetics: evolution of the G matrix. *Trends Ecol Evol* 17(7):320–327
- Stevenson RD, Hill MF, Bryant PJ (1995) Organ and cell allometry in Hawaiian *Drosophila*: how to make a big fly. *Proc Biol Sci* 259:105–110
- Summers TC, Ord TJ (2021) The stabilising impact of natural selection on the allometry of sexual ornaments: Fish that escape locomotor constraints exhibit extravagant ornamentation. *Funct Ecol*. <https://doi.org/10.1111/1365-2435.13968>
- Tan S, Zhong Y, Hou H et al (2012) Variation of presence/absence genes among *Arabidopsis* populations. *BMC Evol Biol* 12:86
- Tisné S, Reymond M, Vile D et al (2008) Combined genetic and modeling approaches reveal that epidermal cell area and number in leaves are controlled by leaf and plant developmental processes in *Arabidopsis*. *Plant Physiol* 148:1117–1127
- Toussaint A, Brosse S, Bueno CG, Pärtel M, Tamme R, Carmona CP (2021) Extinction of threatened vertebrates will lead to idiosyncratic changes in functional diversity across the world. *Nat Commun* 12(1):1–12
- Vasseur F, Violle C, Enquist BJ et al (2012) A common genetic basis to the origin of the leaf economics spectrum and metabolic scaling allometry. *Ecol Lett* 15:1149–1157
- Vasseur F, Bontpart T, Dauzat M et al (2014) Multivariate genetic analysis of plant responses to water deficit and high temperature revealed contrasting adaptive strategies. *J Exp Bot* 65:6457–6469
- Vasseur F, Bresson J, Wang G et al (2018a) Image-based methods for phenotyping growth dynamics and fitness components in *Arabidopsis thaliana*. *Plant Methods* 14:63
- Vasseur F, Exposito-Alonso M, Ayala-Garay OJ et al (2018b) Adaptive diversification of growth allometry in the plant *Arabidopsis thaliana*. *Proc Natl Acad Sci USA* 99:748
- Violle C, Navas ML, Vile D et al (2007) Let the concept of trait be functional! *Oikos* 116:882–892
- Wagner GP, Pavlicev M, Cheverud JM (2007) The road to modularity. *Nat Rev Genet* 8(12):921–931
- Wagner GP, Kenney-Hunt JP, Pavlicev M et al (2008) Pleiotropic scaling of gene effects and the “cost of complexity.” *Nature* 452:470–472
- Watanabe J (2022) Statistics of eigenvalue dispersion indices: quantifying the magnitude of phenotypic integration. *Evolution* 76(1):4–28
- Weigel D (2012) Natural variation in *Arabidopsis*: from molecular genetics to ecological genomics. *Plant Physiol* 158:2–22
- Weisweiler M, de Montaigu A, Ries D et al (2019) Transcriptomic and presence/absence variation in the barley genome assessed from multi-tissue mRNA sequencing and their power to predict phenotypic traits. *BMC Genomics* 20:787
- West GB, Brown JH, Enquist BJ (1999a) A general model for the structure and allometry of plant vascular systems. *Nature* 400:664–667
- West GB, Brown JH, Enquist BJ (1999b) The fourth dimension of life: fractal geometry and allometric scaling of organisms. *Science* 284:1677–1679
- West GB, Brown JH, Enquist BJ (2001) A general model for ontogenetic growth. *Nature* 413:628–631
- West GB, Enquist BJ, Brown JH (2002) Modelling universality and scaling. *Nature* 420:626–627
- White CR, Seymour RS (2003) Mammalian basal metabolic rate is proportional to body mass^{2/3}. *Proc Natl Acad Sci U S A* 100:4046–4049
- Zhang J (2018) Neutral theory and phenotypic evolution. *Mol Biol Evol* 35:1327–1331
- Zhang L-M, Luo H, Liu Z-Q et al (2014) Genome-wide patterns of large-size presence/absence variants in sorghum. *J Integr Plant Biol* 56:24–37
- Zinner M, Lukonin I, Liberali P (2020) Design principles of tissue organisation: how single cells coordinate across scales. *Curr Opin Cell Biol* 67:37–45

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.