



HAL
open science

Differential Genetic Strategies of *Burkholderia vietnamiensis* and *Paraburkholderia kururiensis* for Root Colonization of *Oryza sativa* subsp. *japonica* and *O. sativa* subsp. *indica* , as Revealed by Transposon Mutagenesis Sequencing

Adrian Wallner, Nicolas Busset, Joy Lachat, Ludivine Guigard, Eoghan King, Isabelle Rimbault, Peter Mergaert, Gilles Béna, Lionel Moulin

► To cite this version:

Adrian Wallner, Nicolas Busset, Joy Lachat, Ludivine Guigard, Eoghan King, et al.. Differential Genetic Strategies of *Burkholderia vietnamiensis* and *Paraburkholderia kururiensis* for Root Colonization of *Oryza sativa* subsp. *japonica* and *O. sativa* subsp. *indica* , as Revealed by Transposon Mutagenesis Sequencing. *Applied and Environmental Microbiology*, 2022, 88 (14), 10.1128/aem.00642-22 . hal-03747731v1

HAL Id: hal-03747731

<https://hal.inrae.fr/hal-03747731v1>

Submitted on 28 Oct 2022 (v1), last revised 18 Jan 2024 (v2)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

1 Title: **Differential genetic strategies of *Burkholderia vietnamiensis* and *Paraburkholderia kururiensis* for root**
2 **colonization of *Oryza sativa* ssp. *japonica* and ssp. *indica*, as revealed by Tn-seq**

3 Running title: Tn-seq of burkholderia genes for rice root colonization

4 Adrian Wallner^{a,*}, Nicolas Busset^a, Joy Lachat^{b,*}, Ludivine Guigard^a, Eoghan King^a, Isabelle Rimbault^a,
5 Peter Mergaert^b, Gilles Béna^a, Lionel Moulin^{a,#}

6 ^aPHIM Plant Health Institute, Univ Montpellier, IRD, CIRAD, INRAE, Institut Agro, Montpellier, France

7 ^bUniversité Paris-Saclay, CEA, CNRS, Institute for Integrative Biology of the Cell (I2BC), 91198, Gif-
8 sur-Yvette, France

9 *Present addresses: AW, Université de Reims Champagne-Ardenne, RIBP EA4707 USC INRAE 1488,
10 SFR Condorcet FR CNRS 3417, Reims, France

11 JL, Center for Immunology and Infectious Diseases, Cimi-Paris, Inserm, Sorbonne Université, Paris,
12 France

13

14 #Corresponding author: Lionel Moulin, PHIM, 911 Avenue Agropolis 34394 Montpellier cedex 5. E-
15 mail: lionel.moulin@ird.fr

16 Keywords: Rice, *Paraburkholderia kururiensis*, *Burkholderia vietnamiensis*, Root-colonization, Tn-seq,
17 PGPR.

18 Abstract

19 *Burkholderia vietnamiensis* LMG10929 (*Bv*) and *Paraburkholderia kururiensis* M130 (*Pk*) are bacterial
20 rice growth-promoting models. Besides this common ecological niche, species of the *Burkholderia* genus
21 are also found as opportunistic human pathogens while *Paraburkholderia* are mostly environmental and
22 plant-associated species. Here, we compared the genetic strategies used by *Bv* and *Pk* to colonize two
23 subspecies of their common host, *Oryza sativa* ssp. *japonica* (cv. Nipponbare) and ssp. *indica* (cv. IR64).
24 We used high-throughput screening of transposon insertional mutant libraries (Tn-seq) to infer which
25 genetic elements have the highest fitness contribution during root surface colonization at 7 days post
26 inoculation. Overall, we detected twice more genes in *Bv* involved in rice roots colonization compared to
27 *Pk*, including genes contributing to the tolerance of plant defenses, which suggests a stronger adverse
28 reaction of rice towards *Bv* compared to *Pk*. For both strains, the bacterial fitness depends on a higher
29 number of genes when colonizing *indica* rice compared to *japonica*. These divergences in host pressure
30 on bacterial adaptation could be partly linked to the cultivar's differences in nitrogen assimilation. We

31 detected several functions commonly enhancing root colonization in both bacterial strains e.g., Entner-
32 Doudoroff (ED) glycolysis. Less frequently and more strain-specifically, we detected functions limiting
33 root colonization such as biofilm production in *Bv* and quorum sensing in *Pk*. The involvement of genes
34 identified through the Tn-seq procedure as contributing to root colonization i.e., ED pathway, c-di-GMP
35 cycling and cobalamin synthesis, was validated by directed mutagenesis and competition with WT strains
36 in rice root colonization assays.

37 Importance

38 Burkholderiaceae are frequent and abundant colonizers of the rice rhizosphere and interesting candidates
39 to investigate for growth promotion. Species of *Paraburkholderia* have repeatedly been described to
40 stimulate plant growth. However, the closely related *Burkholderia* genus hosts both beneficial and
41 phytopathogenic species, as well as species able to colonize animal hosts and cause disease in humans.
42 We need to understand to what extent the bacterial strategies used for the different biotic interactions
43 differ depending on the host and if strains with agricultural potential could also pose a threat towards
44 other plant hosts or humans. To start answering these questions, we used here transposon sequencing to
45 identify genetic traits in *Burkholderia vietnamiensis* and *Paraburkholderia kururiensis* that contribute to
46 the colonization of two different rice varieties. Our results revealed large differences in the fitness gene
47 sets between the two strains and between the host plants, suggesting a strong specificity in each
48 bacterium-plant interaction.

49 Introduction

50 Species of the *Burkholderia* and closely related *Paraburkholderia* genera are highly prolific rhizosphere
51 colonizers (1, 2). Their persistence and competitiveness in the rhizosphere environment can be explained
52 by strong secondary metabolite production as well as efficient nitrogen cycling, mineral solubilization and
53 phytohormone biosynthesis (3–5). Furthermore, several *Paraburkholderia* and at least one *Burkholderia*
54 species can fix atmospheric nitrogen. Two well studied models, *Paraburkholderia kururiensis* strain
55 M130 (hereafter called *Pk*) and *Burkholderia vietnamiensis* strain LMG10929 (hereafter called *Bv*)
56 demonstrate strong rice root colonization, endophytic lifestyles and significant plant growth promotion
57 through transfer of fixed nitrogen (6–9). Despite this convergence in their plant beneficial features, both
58 bacteria belong to distinct genetic backgrounds. While *Paraburkholderia* species are often found in
59 beneficial relationships and symbioses with plants (10–12), *Burkholderia* members comprise human
60 pathogens and opportunists as well as fungal and plant pathogens (13).

61 Bacterial genes used for plant colonization have been screened in several model bacteria but also on a
62 broader scale using microbiome approaches to reveal plant-associated functions (1). A few studies have

63 compared plant-adapted *Burkholderia s.l.* (*sensu largo*, e.g. the former genus that is now regrouping the
64 newly defined *Burkholderia*, *Paraburkholderia*, *Caballeronia* and others) at the genomic level (4, 11), or
65 profiled the transcriptome of bacteria stimulated by root exudates (14, 15). However, there is no record of
66 a comparison between the strategies used by plant-adapted *Burkholderia* and *Paraburkholderia* species.
67 The host plant genotype's impact on bacterial colonization strategies also remains poorly explored in
68 these bacterial genera. Rice is an interesting model to assess *Burkholderia s.l.* adaptation to the plant
69 environment as it is hosting plant growth promoting model strains from both *Burkholderia* and
70 *Paraburkholderia* genera. It was repeatedly demonstrated that rice genotypes influence the composition
71 of their microbiome at the rhizosphere and rhizoplane levels (16, 17). In particular, a study on 95 different
72 *Oryza sativa* subsp. *indica* and subsp. *japonica* varieties showed significant differences in microbiome
73 composition between both rice subspecies, which was related to their nitrogen use efficiency and the
74 presence of a particular nitrate transporter in *indica* varieties (18).

75 Transposon mutagenesis sequencing (Tn-seq), is a high-throughput screening method that combines
76 transposon insertional mutagenesis followed by sequencing of the insertion sites (19). It is a powerful tool
77 that leads to immediate identification of genes of interest improving or reducing the bacteria's fitness in a
78 tested condition. This methodology has been successfully used to unravel important genes and functions
79 in plant-pathogenic or plant-symbiotic interactions (20–22). Commonly between pathogens and
80 mutualists, genes functioning in amino acid and purine metabolism as well as in cell motility were
81 detected to be required for root colonization (21, 23, 24).

82 In the present study, we used Tn-seq, to perform a genome-wide identification of genes involved in rice-
83 root colonization in *Bv* and *Pk*. In detecting which genes influence the fitness of *Bv* and *Pk* we aim at
84 unraveling their commonalities and differences in root colonization strategies. We also analyzed the
85 association strategies of *Bv* and *Pk* with the two rice genotypes *Oryza sativa* subsp. *japonica* (cv.
86 Nipponbare) and *indica* (cv. IR64) to understand how the host-factor can influence colonization
87 strategies. Overall, we identified a total of 1,404 and 540 genes that influence the fitness of *Bv* and *Pk*
88 respectively during rice root colonization. Our results underline the importance of motility, amino acid
89 and vitamin metabolism, stress response as well as biofilms for the efficient association of these bacteria
90 with rice roots.

91 Results

92 Quality of Tn-Seq libraries and essential genomes in *Pk* and *Bv*

93 To generate a genome-wide library of insertion mutants for *Pk* and *Bv* strains we used a mariner
94 transposon that targets genomic thymine-adenosine (TA) sites (Materials and methods). The genome of

95 *Bv* contains 95,075 TA sites and we estimated the mutant population at 1.6×10^8 cfu, which represents a
96 1683x coverage of the total TA sites. The saturation level of the *Pk* library was lower although still
97 significant at a 38x coverage given the 4.0×10^6 cfu obtained after transposon mutagenesis for a total of
98 106,136 total TA sites contained in the genome.

99 To further assess the quality of the Tn-seq libraries we determined the essential genomes required by both
100 bacteria for optimal development in a rich liquid growth medium. Both bacteria have similar genome
101 sizes (6,820 and 6,436 genes for *Bv* and *Pk* respectively) and comparable proportions of genes (661 or
102 9.7% and 638 or 9.9%, respectively) that are required for optimal growth in a rich, liquid medium (**Figure**
103 **1A; Supplemental dataset S1**). The size of the essential genomes determined in a controlled liquid
104 medium are in the order of magnitude of what has been observed for other prokaryotes, including
105 *Burkholderia spp.* (26–30). We used the Minimal Gene Set tool implemented in the MicroScope platform
106 (Materials and methods) to extract a core list of the predicted minimal bacterial gene sets from *Bv* and *Pk*.
107 On a total of 206 core bacterial genes defined by their conservation among multiple bacterial genomes
108 (25), 151 and 150 were identified as essential by our approach in *Bv* and *Pk*, respectively. In 10 and 9
109 cases of genes belonging to the minimal gene set but predicted as non-essential by our approach in *Bv* and
110 *Pk* respectively, there are duplicates present in the genome.

111 According to their distribution in COG categories, little variance differentiates the essential genomes of
112 *Pk* and *Bv* (**Figure 1B**). As expected in a liquid, rich medium, genes involved in motility, defense and
113 nutrition are largely unessential. Conversely, structural components of the cell and the general replicative
114 machinery are predictably essential (**Supplemental dataset S1**).

115 Given the strong saturation level of the TA insertion sites and the coherent essentiality results observed in
116 the control setting, we can safely assume that both mutant libraries allow a reliable analysis of the impact
117 of genes on the bacterial fitness.

118 Colonization of the two rice genotypes by *Pk* and *Bv*

119 Prior to genetic analyses, we assessed the colonization efficiencies of *Bv* and *Pk* on Nipponbare
120 (*japonica*) compared to IR64 (*indica*) rice genotypes (**Figure 2**). Overall, both bacteria display a similar
121 colonization dynamic with an increasing population density during the first week and a decrease in the
122 second week. A host-genotype effect is observed in the colonization phenotype displayed by *Pk* as
123 significantly different root colonizing populations are observed on IR64 and Nipponbare at 3 dpi and 14
124 dpi (**Figure 2**). *Bv* on the other hand displays a steady colonization pattern between both plant genotypes
125 at all measured time points. These observations confirm that the rice-colonizing populations of *Bv* and *Pk*
126 can be compared, especially at 7 dpi, which was selected for further analyses. In our following analyses,

127 we consider that the bacterial adaptations observed result purely from root surface colonization, as the
128 endophytic populations at 7 dpi are inferior by several log scales to the surface colonizing population (6).

129 Identification of Rice colonization genes

130 To assess gene fitness for rice root colonization, and its host-dependent variation in the two model
131 bacteria, we inoculated the Tn-Seq mutant libraries on Nipponbare and IR64 rice genotypes. Seven days
132 after inoculation, we harvested and pooled five root systems per replicate, representing a total of 1.2×10^7
133 colonization events (**Figure 2**) and a more than 100-fold coverage of the available mutant diversity. We
134 performed a first Tn-seq analysis by pooling the reads of IR64 and Nipponbare isolates together to infer
135 the genes globally required for the association with rice. The read frequencies were compared to a control
136 condition grown in a rich medium with limited growth generations (see Material and Methods) to
137 establish a root fitness score for each gene. In this manner, we identified 1,404 and 540 genes as
138 significantly impacted (enriched or depleted) after root-colonization by *Bv* and *Pk*, respectively (**Figure**
139 **3A**). Colonization-depleted genes will be our major focus as they are positively associated with bacterial
140 fitness. Inversely, colonization-enriched genes diminish the bacterial fitness during root colonization. We
141 organized the identified colonization genes according to their general function based on their clusters of
142 orthologous groups (COG) annotation (**Figure 3B**). Colonization-enriched and -depleted genes share a
143 similar distribution with amino acid metabolism, cell wall/membrane/envelope biogenesis, transcription
144 and cell motility being amongst the most frequent categories, consistent with the expected implication of
145 nutrition, motility and morphological adaptations involved in plant colonization.

146 We identified a total of 2,071 core gene families for *Bv* and *Pk*, sharing homologues in both strains
147 (**Supplemental dataset S2**). In *Bv* and *Pk*, 192 genes, respectively 24% and 41% of the colonization-
148 depleted genes are part of the core-genome and equally depleted in both strains (**Figure 4**). *Bv* displays
149 68% more colonization-depleted genes than *Pk* (**Figure 3**). Interestingly, a majority of these *Bv* specific
150 colonization-depleted genes (53%) are part of the core genome (**Figure 4**). Similarly, a lower, although
151 substantial portion of the genome shared with *Bv* (37%), is required by *Pk* specifically for efficient root
152 colonization (**Figure 4**). Thus, although a large portion of the core genes contribute to root colonization in
153 both strains, many core genes are only required for colonization in one or the other strain, indicating that
154 the impact of these genes on root colonization is dependent on the genome context.

155 Rice cultivar dependent colonization specialization

156 Next, we performed a separate analysis of the reads for each rice cultivar and then compared the lists of
157 significantly colonization-depleted and -enriched genes to infer cultivar-specificities. For both strains, the
158 majority (~65%) of identified genes impacted the bacterial fitness on both IR64 and Nipponbare rice

159 (Table 1). However, both bacteria displayed a greater number of colonization-depleted genes (41% and
160 49% higher for *Bv* and *Pk* respectively) on IR64 compared to Nipponbare rice (Table 1).

161 Core genes of *Bv* and *Pk* that are colonization-depleted on both cultivars belong primarily to
162 housekeeping categories such as central metabolism, cell cycle control and motility (Figure 5A). The
163 additional genes, depleted during IR64 colonization are chiefly attributed to amino acid metabolism,
164 transcription, cell wall/membrane/envelope biogenesis and energy production in both bacterial strains
165 (Figure 5B & 5C). Interestingly, contrary to the global trend, in *Pk* the COG categories “ replication,
166 recombination & repair”, “coenzyme transport & metabolism”, “carbohydrate metabolism”, and
167 “inorganic ion transport” are more strongly impacted on Nipponbare than IR64 (Figure 5C). The same
168 trend is observed in *Bv* for the COG categories “signal transduction mechanisms” and “coenzyme
169 transport and metabolism” (Figure 5B).

170 Tn-seq reveals bacterial functions involved in early rice colonization

171 Many genes significantly impacting the colonization fitness of *Bv* and *Pk* are clustered together within
172 operons supporting the validity of our results. In these cases, there is furthermore a strong conservation of
173 either gene enrichment or depletion within the same operon. In several cases explored hereafter, the lack
174 of detection of a complete operon can be explained by the presence of homologues for some of the genes
175 resulting in functional complementation.

176 Functions required for bacterial fitness during rice-root colonization

177 As expected in this kind of colonization assay, mutants affected in motility and chemotaxis functions (*flg*,
178 *flh*, *fli*, *mot*) were depleted from the root colonizing population (Table 2 & Supplemental dataset S3).
179 For both bacteria, we further detected many genes involved in amino acids (*arg*, *his*, *ilv*, *leu*, *lys*, *met*, *pro*,
180 *ser* and *trp*) and nucleotide synthesis (*pur* and *pyr*) that, when mutated, negatively impacted the bacteria's
181 fitness on plants. Genes involved in the synthesis of enzymatic cofactors for amino acid metabolism such
182 as vitamin B1 (thiamin) were also colonization-depleted in both strains. Multiple Tn-seq studies reported
183 that auxotrophy for certain amino acids is disadvantageous for root colonization and can limit plant
184 growth promotion and biocontrol potential (23, 31).

185 Additionally, genes involved in potassium nutrition (*kdpA-F*) were similarly depleted. Mutants of both *Bv*
186 and *Pk* affected in central elements of the Entner-Doudoroff glycolysis (ED) pathway (*edd*, *zwf*) suffered
187 a significant fitness decrease on rice. This pathway is involved in the metabolism of gluconate which is
188 not a dominant sugar of rice exudates (32) suggesting an alternative role than sole sugar assimilation for
189 this pathway. The activation of the ED pathway was suggested to be a tolerance strategy towards
190 oxidative stress through the generation of NADPH as essential cofactor for thioredoxins (33, 34).

191 Genes of the type 2 secretion system (T2SS; *gspD-M*) are the only ones belonging to a macromolecular
192 secretory pathway to be colonization-depleted in both bacteria. While *Bv* possesses a single T2SS, *Pk*
193 bears two copies (4) out of which only one is colonization-depleted suggesting that these systems are not
194 redundant but are used differently in specific conditions.

195 Several genes involved in DNA maintenance and repair, i.e., *ruvABC*, *xerCD* and *recABCD* were
196 colonization-depleted in both *Bv* and *Pk* while their absence was tolerated in the relatively stress-free
197 control medium (**Supplemental dataset S1**). Thus, rice appears to be inflicting considerable genotoxic
198 stress during the process of colonization. Furthermore, the presence of osmotic stress is exemplified by
199 the depletion of *Bv* and *Pk* mutants involved in the synthesis of the osmoprotectant trehalose (*otsAB*).

200 *Bv* seems to suffer additional stress as multiple functions maintaining cell wall and membrane integrity
201 are colonization-depleted. We detected several genes involved in hopanoid synthesis (*hpnDEFHJKN*),
202 peptidoglycan synthesis (*murAI*) and maintenance (*tolAQR*, *pal*) as well as lytic murein
203 transglycosylation (*mltA*, *rlpA* and *mtgA*) to be depleted specifically in *Bv* mutant populations. This
204 indicates that *Bv* has an increased requirement to maintain its cellular integrity compared to *Pk*.
205 Consistent with an increased need in membrane maintenance, the loss of vitamin B7 (biotin) synthesis
206 genes (*bioABCDF*) had a negative impact on the colonization of *Bv*. Biotin is a cofactor for many
207 enzymes, especially those involved in fatty acid biosynthesis and amino acid metabolism (35, 36). Roots
208 are known to secrete aromatic phenolic compounds which are toxic to various soil microbes. Accordingly,
209 *Bv* mutants were depleted during colonization when impacted in genes of the β -ketoadipate pathway
210 (*pcaBCDK*) which allow to metabolize 4-hydroxybenzoate and protocatechuate. A final sign of the stress
211 *Bv* is exposed to during colonization is found in the depletion of mutants for the queuosine synthesis
212 pathway (*queACEF*, *tgt*). This hypermodified nucleoside improves translation accuracy, a need that only
213 arose in *Bv* during rice colonization.

214 While no genes annotated as coding for autotransporter proteins (T5SS) are identified through our
215 approach, the genes *tamAB* were colonization-depleted in *Pk*. TamA and TamB can be involved in outer
216 membrane assembly, allowing surface structuration which is essential for adhesion and host-invasion by
217 bacteria in several pathogenic models (37). Genes for vitamin B12 (cobalamine) synthesis are present in
218 both *Bv* and *Pk* but were only colonization-depleted in *Pk* (*cobBDHIKLMNQSTUW*, *cbiB*, *btur*).
219 Intriguingly, several genes of this synthesis pathway were actually colonization-enriched in *Bv*,
220 suggesting an adverse role in colonization for this co-factor involved in a multitude of enzymatic
221 reactions.

222 Functions detrimental for bacterial fitness during rice-root colonization

223 Only two genes were identified to be colonization-enriched in *Bv* and *Pk* simultaneously, one of which
224 (AK36_1927/ANSKv1_30041) has an unknown function. The other (*pdeR*) is involved in cyclic di-GMP
225 (c-di-GMP) cycling. In *Bv*, four other genes having a homologous role in c-di-GMP degradation
226 (phosphodiesterases; **Supplemental dataset S3**) and two involved in c-di-GMP production (diguanylate
227 cyclases; **Supplemental dataset S3**) were also colonization-enriched. C-di-GMP levels can play a role in
228 colonization as they were demonstrated to influence biofilm production and motility in *Burkholderia*
229 species (38). Interestingly, while several genes of the putrescine catabolism (*puuBD*, *speG*) were
230 colonization-depleted in *Bv*, the entire cluster responsible for putrescine uptake (*potFGHI*, *puuP*) is
231 colonization-enriched. Together, this suggests that putrescine accumulation has a negative impact on root
232 colonization. An accumulation of intracellular putrescine could have an impact on biofilm production as
233 recently suggested (23). Among additional biofilm-related colonization-enriched genes in *Bv*, we
234 identified a cluster involved in cellulose synthesis (*bcsABCE*). Another *Bv* surface element that increases
235 colonization efficiency when mutated is a Tad pilus (*tadABCEVZ*, *rcpAB*, *flp*) suggesting that this
236 attachment or motility mechanism is suitable for an alternative condition than rice colonization. While
237 many genes involved in carbohydrate metabolism in both bacteria were colonization-depleted, *Bv* mutants
238 were colonization-enriched when impacted in ribose import and catabolism genes (*rbsABCK*).

239 *Pk* mutants impacted in the ability to sense the bacterial community through quorum sensing (*braR*, *rsaL*)
240 were enriched during colonization as can be expected for organisms that lost the ability to sense
241 population density and reduce growth rates accordingly. Genes belonging to a region enriched in elements
242 of phage origin (ANSKV1_30083-120) had the same effect when mutated, indicating that the prophage
243 might be induced by the plant colonization process and that its inactivation through mutation benefits the
244 bacteria's fitness.

245 Rice cultivar specific adaptations

246 Genes which are colonization-depleted or -enriched specifically on one cultivar can be more rarely
247 grouped in metabolic pathways and operon structures (**Supplemental dataset S4**). Still, single genes can
248 have determining impacts on a strain's ecology and metabolism. There are two genes that were
249 colonization-depleted for both *Bv* and *Pk* on Nipponbare rice. One is an outer membrane porin of the
250 OmpC family (AK36_1494 / ANSKv1_11218) and the other, *amtB*, is involved in ammonium uptake
251 (**Table 3 & Supplemental dataset S4**). *Bv* mutants for genes involved in phosphate transport (*pstAC*)
252 and glycerol (*ugpABQ*) metabolism were depleted on IR64 rice, further suggesting that the metabolic
253 requirements of the plant force adaptations on rhizosphere microbes.

254 Several *Bv* genes involved in type 6 secretion (*tssGHIJK*) were colonization-depleted on IR64. For *Pk*,
255 mutants impacted in O-antigen biosynthesis (*rfbABD*) were depleted on IR64. In both cases we can
256 hypothesize that surface elements have significant and diverging repercussions depending on the type of
257 colonized rice-host.

258 Validation of three candidate genes

259 We underlined the importance of various genes involved in the ED pathway, c-di-GMP cycling and
260 cobalamin synthesis among others. To validate their involvement in the colonization of rice roots, we
261 used an insertional mutagenesis approach in *Bv* and *Pk* by targeting single copy genes that are key in the
262 respective processes. The 6-phosphogluconate dehydratase *Edd* is essential for the first step of the ED
263 pathway (39). The metal delivery protein *CobW* is essential for vitamin B12 synthesis (40). Finally, we
264 chose *PdeR*, an enzyme with domains predicted to be involved in both c-di-GMP production and
265 degradation. The genomic context of each of these genes, including their fold change between root &
266 rich-medium conditions, is presented in **Supplementary Figure S1**.

267 Mutants and wild-type strains were used jointly in a competition assay and colonization efficiencies were
268 surveyed at 7 dpi. In *Bv*, each mutation had the observable impact that had been predicted by the Tn-seq
269 analysis (**Figure 6**). Disruption of the colonization-depleted gene *edd* reduced *Bv*'s colonization capacity
270 while the deletion of the colonization-enriched gene *pdeR* had the opposite effect. The disruption of *cobW*
271 had been identified in Tn-seq data as having a deleterious impact on *Pk* but did not significantly alter the
272 root colonization capacity of *Bv*. The mutagenesis approach confirms that the disruption of *cobW* has no
273 negative impact on *Bv* but is required by *Pk* for efficient root colonization. The only inconsistent
274 observation occurred for the *pdeR* mutant in *Pk*, which was predicted as slightly colonization-enriched by
275 the Tn-seq approach, but its disruption resulted in a colonization deficiency compared to the WT strain.

276 Discussion

277 Understanding which genetic bases are involved in PGPR-cereal interaction is pivotal for a controlled and
278 informed selection of beneficial organisms and a gateway to efficient and directed strain improvement. In
279 rice, the microbiome composition was demonstrated to be significantly modulated by the plant-genotype
280 (16). This suggests a variance in selective pressure forced by the plant onto colonizing microbes. The
281 present Tn-seq analysis demonstrates which genes have the strongest contribution to bacterial fitness in
282 the early steps of rhizosphere colonization of two rice cultivars, Nipponbare and IR64 by two bacteria *Pk*
283 and *Bv* (**Figure 7**). At this stage, we cannot exclude that additional genes might be required to colonize
284 more mature plants as the bacteria progress and potentially reach different plant compartments. It is also
285 known that Tn-seq approaches are insensitive to genes whose function is complemented by functional

286 homology either within the same bacteria (gene duplicates, paralogues) or by the surrounding bacterial
287 community, e.g., secretion systems and secreted molecules.

288 The present study compares insertion mutant libraries that were grown on plant roots for 7 days to
289 libraries grown in a rich liquid medium for a brief period. Thus, caution needs to be taken when
290 interpreting the role of the detected genes in plant interaction. Especially detected nutritional function
291 could be general prerequisites for a sound bacterial development in a poor medium and not specifically
292 involved in a plant colonization task. Still, we find several examples of genes involved in plant interaction
293 that are supported by independent approaches, as illustrated hereafter.

294 *Bv* and *Pk* have different strategies for rice root colonization

295 As was previously suggested by microscopy observations and host transcriptomics, *Bv* and *Pk* have
296 different approaches to rice colonization (6). The present analysis highlights the respective genetic
297 requirements of *Bv* and *Pk* during their rice colonizing process. Strikingly, a considerable higher number
298 of genes is necessary for a successful colonization by *Bv* than by *Pk*, despite similar genome size. This
299 requirement could result from increased plant defenses that *Bv* is exposed to compared to *Pk* (6).
300 Supposedly, the pathogenic genetic background and potential of *Bv* could be responsible for this adverse
301 plant response and necessitate a higher number of genes to act against. Moreover, as *Bv* is known to
302 colonize multiple eukaryotic hosts, both plants and animals (41–44), we can also hypothesize that *Bv*
303 expresses many genes which are not of use for plant colonization, thus a mutation in these genes will
304 increase the bacteria's fitness as the ensuing metabolic cost can be spared. This could explain that plant-
305 detrimental genes were found in relatively similar amounts to plant-essential genes in *Bv* while they are
306 considerably less frequent in *Pk* (**Figure 3A**). One striking example is the case of vitamin B12
307 production. This cofactor is among the most complex in bacteria and requires at least 19 enzymatic steps
308 to produce while many enzymes further depend on B12 for their activity (45). Here, the synthesis of B12
309 is beneficial for the fitness of *Pk* during colonization while it has an adverse effect on *Bv*. This
310 observation is supported both by the Tn-seq and the insertional mutagenesis approaches. We can
311 hypothesize that the cofactor is involved in enzymatic activities that play a crucial role for colonization in
312 *Pk* but not in *Bv*. As the production of B12 is associated with a high metabolic cost, sparing its expense
313 could explain the fitness advantage displayed by *Bv* when the B12 synthesis pathway is mutated.

314 Defects in secretory activities are readily complemented in bacterial communities making the responsible
315 secretion systems opaque for detection by Tn-seq approaches. Indeed, secretion systems are not detected
316 by most studies relying on Tn-seq except when focusing on more sparsely colonized environment such as
317 the apoplast (31). In the present study however, a T2SS is colonization-depleted in *Bv* and *Pk* and the type
318 6 secretion system (T6SS) in *Bv* during IR64 colonization. While the T6SS can be used for cell adhesion

319 (46), a role beyond complementable secretion remains elusive for the T2SS. Still, as only one T2SS is
320 supposedly involved in colonization it suggests a separate role from the second one present in *Pk*.

321 The only gene with an identified function that is predicted by the Tn-seq approach to be negatively
322 involved in root colonization in both *Bv* and *Pk* is *pdeR*, Site-directed mutagenesis and a competition
323 assay revealed that its mutation was highly beneficial for the root colonization activity of *Bv* as expected
324 from the Tn-seq results, but detrimental for *Pk*, which was inconsistent with the Tn-seq analysis. PdeR is
325 involved in the turnover of the secondary messenger c-di-GMP. In *Xanthomonas oryzae*, the deletion of
326 *pdeR* results in a decreased virulence on rice (47). Through its association in two component signal
327 transduction systems, PdeR could have a variety of indirect roles beside c-di-GMP cycling (48). The
328 divergence observed between the Tn-seq and directed mutagenesis for *Pk* could also be linked to the
329 inoculum's concentration which is likely to have been superior in the latter approach and might have
330 triggered a different plant response with the observed detrimental effects on the *Pk* mutant population
331 (**Figure 6**).

332 Tn-seq identifies common plant-colonization traits

333 To date, there is no transcriptome data available for *Bv* that would allow a comparison with our data.
334 However, the transcriptomic response of *Pk* to rice root macerates has been assessed before (14). Out of
335 the 471 colonization-depleted genes identified by the present approach, 267 are differentially expressed in
336 *Pk* when stimulated with root macerate (**Supplemental dataset S5**). Dominantly, these genes are
337 involved in amino acid metabolism, cell motility and cell wall/membrane biogenesis. The prevalence of
338 these functions has further been reported by Tn-seq approaches in other root colonizing models such as
339 *Dickeya dadantii* and *Pseudomonas spp.* (21, 23, 24). Our results are congruent with most plant-microbe
340 colonization studies, underlining the importance of genes involved in the production of surface
341 components for cellular attachment to the host (49). Our analysis also presents substantial overlap of
342 colonization-depleted genes with what has been found in the plant growth promoting species *Azoarcus*
343 *olearius* and *Herbaspirillum seropedicae* (50). Notably genes involved in peptidoglycan (*ampD*) and cell
344 wall (*murAI*) formation, chemotaxis (*cheARW*), iron uptake (ferredoxins) and cobalamin synthesis (*cbiA*,
345 *cobIO*) were detected.

346 Disruption of the ED pathway was demonstrated to reduce root colonization capacity in *Pseudomonas*
347 *chlororaphis* with a subsequent loss of ISR stimulation (51). Later works on *Pseudomonas fluorescens*
348 observed that the expression of *edd* was enhanced in the rhizosphere of *Arabidopsis* compared to liquid
349 growing medium (52). Mutants for *edd* in this latter study also failed to stimulate ISR but without the
350 colonization defect observed in *P. chlororaphis*. For pseudomonads, the importance of the ED pathway

351 might be linked to its direct product, pyruvate, that is required for the synthesis of the ISR-eliciting
352 butanediol (51). Other advantages might arise from the ED pathway such as the production of NAD(P)H
353 which is not generated by the EMP glycolysis pathway. This cofactor is used by thioredoxins and could
354 be involved in plant colonization and associated oxidative stress tolerance (33, 34).

355 IR64 rice has more stringent requirements for colonization by *Bv* and *Pk* than Nipponbare rice.
356 For both *Bv* and *Pk*, a stronger host-genotype effect on the number of colonization-impacted genes was
357 observed during the association with IR64 compared to Nipponbare rice. For *Pk*, this genotype-effect is
358 also observed on the root colonization profiles of the two cultivars (**Figure 2**). One specificity of *indica*
359 rice (IR64) over *japonica* (Nipponbare) is its improved nitrate uptake and assimilation capacity (53),
360 which is linked with the presence of the nitrate transporter NRT1.1B in the former, and proved to impact
361 its microbiota (18). Nipponbare rice preferably imports the alternative nitrogen source ammonium, which
362 can be correlated with the depletion of bacterial mutants impacted in the import of this nutrient (*amtB*) in
363 the Nipponbare environment in our experimental set-up, as bacteria would compete with rice for the same
364 nitrogen source) (**Table 3**).

365 It was interesting to observe that the T6SS, a major macromolecular system of *Bv*, was only colonization-
366 depleted during inoculation on IR64 roots. The T6SS can be employed by bacteria for competition with
367 other prokaryotes but also interaction with eukaryotes (54). As secretory function should be
368 complemented by the bacterial community, the T6SS of *Bv* appears to be rather involved in adherence to
369 eukaryotic cells, in a host-dependent manner.

370 Still, most colonization-impacted genes are conserved between rice cultivars and several functions such as
371 motility, amino acid metabolism and biofilm production have been repeatedly described for their role in
372 the general association of bacteria with plants (**Figure 7**). We have found evidence of an increased stress
373 to which bacteria are exposed in the near vicinity of plants through the depletion of mutants involved in
374 osmoprotection, toxic compound degradation and DNA reparation. The identification of several functions
375 that are part of the core-genome shared by both bacteria but are only colonization-impacted in one strain
376 reinforces the validity of these observations and the different adaptations that both bacteria must undergo
377 during colonization of rice. This is especially true in the present system as *Bv* and *Pk* induce different
378 levels of plant defenses (6).

379 An increasing amount of plant microbiome studies rely on Tn-seq to reveal the nature of genes underlying
380 root colonization (20, 21, 23, 24, 31, 50). Tn-seq further offers the benefit over more common methods
381 such as RNAseq, to inform on the genes obstructing colonization (mutants with higher colonization
382 fitness), thus presenting a more complete catalogue of factors driving host-bacteria compatibility.

383 Together with other trending methods such as microfluidic visualization technics and synthetic bacterial
384 communities (49), Tn-seq proves here to be a powerful tool for a better understanding of the genetic bases
385 underlying colonization of different hosts.

386 Materials and Methods

387 Bacteria and plant culture conditions

388 The strains used in this study are *Burkholderia vietnamiensis* LMG 10929 (*Bv*) and *Paraburkholderia*
389 *kururiensis* M130 (*Pk*), either wild type, or modified by insertion of the pIN29 plasmid conferring
390 chloramphenicol (Cm) resistance and DsRed fluorescence, or spontaneous rifampicin (Rif) and
391 spectinomycin (Spt) resistant strains, or insertional mutants of a non-replicative plasmid in several
392 candidate genes in each strain (**Table 4**) (6, 55). Spontaneous antibiotic resistant strains for LMG 10929
393 and M130 were obtained by plating 100 μL of bacterial liquid culture at $\text{OD}_{600}=1.0$ on Luria's low salt
394 LB (LBm; Sigma-Aldrich) with either rifampicin ($30 \mu\text{g}\cdot\text{mL}^{-1}$) or spectinomycin ($50 \mu\text{g}\cdot\text{mL}^{-1}$). After 48h
395 incubation at 28°C , single colonies were selected and streaked on fresh LBm plates with rifampicin or
396 spectinomycin, then grown in broth LBm with antibiotics and stored in 20% glycerol at -80°C . From
397 glycerol stocks, bacterial strains were cultured as follows: bacterial cells conserved at -80°C were plated
398 on LBm agar plates (with antibiotic for mutants) and incubated for 72 h at 28°C . Single colonies were
399 used to inoculate 10 mL of LBm broth (with antibiotic for mutants) in 50 mL Falcon tubes and incubated
400 for various amounts of time allowing the different strains to reach an $\text{OD}_{600}=1.0$. For inoculation
401 purposes, cultures were adjusted to $5 \times 10^7 \text{ cfu}\cdot\text{mL}^{-1}$.

402 *Oryza sativa subsp. japonica* cv. Nipponbare and *Oryza sativa subsp. indica* cv. IR64 were cultured as
403 described in King et al. (2019). Briefly, seeds were sterilized using successive 70% ethanol and 3.6%
404 sodium hypochlorite treatments and germinated seedlings were transferred onto sterile perlite in an
405 airtight hydroponic system. 5-days old seedlings were inoculated with 1 mL of bacterial solution at $5 \times$
406 $10^7 \text{ cfu}\cdot\text{mL}^{-1}$ and grown up to 14 days at 28°C (16h light, 8h dark). For competition assays, five plants
407 were inoculated with a mix at $1.10^7 \text{ cfu}\cdot\text{mL}^{-1}$ containing 0.5 mL of SptR *Bv* or *Pk* strains (previously
408 grown separately in broth LBm + Spt $50 \mu\text{g}\cdot\text{mL}^{-1}$ and washed to remove antibiotic) and 0.5 mL of either
409 one of the insertional mutants (ΩcobW , ΩpdeR or Ωedd) of the corresponding strain (grown in LBm +
410 Cm $100 \mu\text{g}\cdot\text{mL}^{-1}$ and washed to remove antibiotic).

411 Estimation of root colonizing bacterial population

412 For the estimation of root colonizing populations, the plants were prepared as described above and
413 inoculated with WT strains at $1 \times 10^7 \text{ cfu}\cdot\text{mL}^{-1}$. Three systems (15 plants) were prepared for each condition
414 and grown for 1, 7 or 14 days (28°C ; 16 h light; 8 h dark). At each time point, five plants were harvested

415 for each condition. The entire root systems were sampled and separately transferred to screw cap tubes
416 containing 1 mL of sterile water and a sterile ceramic bead. Roots were weighted and then pulverized
417 using a Fastprep-24 (MPbio) at 6 m.s⁻¹ for 40 seconds. A serial dilution of the resulting solution was
418 spotted out on square LBm plates and incubated at 28°C for 48 hours. The number of colony forming units
419 (cfu) were counted and adjusted to the weight of the root systems. The mean colonization values were
420 compared between cultivars and between bacteria genotypes at each time point and for each bacterium
421 separately using a Wilcoxon test. Results were considered significantly different at p<0.05. For the
422 competition assay between WT and mutant strains, plants were harvested at 7 dpi and processed as
423 described above, except the bacterial dilutions were spotted on LBm plates containing either streptomycin
424 (30 µg.ml⁻¹) or Cm (200 µg.ml⁻¹) to select respectively the WT or the mutant strains.

425 Tn-seq library preparation

426 A Himar1 mariner transposon carried by the pSAM_EC vector in *Escherichia coli* strain MFDpir (56, 57)
427 was introduced into *Paraburkholderia kururiensis* M130 Rif^R and *Burkholderia vietnamiensis* LMG
428 10929 Rif^R through conjugation. Both donor and recipient were grown until OD₆₀₀=1.0 in liquid low-salt
429 LB (LBm) and the medium of MFDpir was further supplemented with diaminopimelic acid (DAP; 300
430 µg.mL⁻¹). Cells were spun down and washed once with LBm and concentrated to a final OD₆₀₀=50. Donor
431 and recipient strains were mixed 1:1 and 50 µL suspensions were spotted on square Petri dishes
432 containing LBm and DAP. Growth rates were different for both recipient species and conjugation time
433 was adapted accordingly. The mating mix was incubated for 6 h and 24 h at 28°C for LMG 10929 and
434 M130, respectively. The spots were then resuspended in 2 mL LBm per plate. The mating mix was
435 further spread on LBm Petri dishes with rifampicin (30 µg.mL⁻¹) and kanamycin (50 µg.mL⁻¹) and
436 incubated at 28°C. This positively selects for recipient strains having integrated the transposon and
437 negatively selects for the DAP auxotroph *E. coli* donor. After growth, the colonies were resuspended in 1
438 mL LBm per plate. The library was separated into 1 mL aliquots, and stored in 20% glycerol at -80°C.
439 Additionally, a serial dilution of the mating mix followed by spreading on LBm with rifampicin (30
440 µg.mL⁻¹) and kanamycin (50 µg.mL⁻¹) was used to estimate the abundance of mutants in the library
441 through cfu counting.

442 Tn-seq experimental setup

443 Tn-seq libraries were thawed on ice and diluted in sterile water to OD₆₀₀=1 and then inoculated in 50 mL
444 liquid LBm with rifampicin (30 µg.mL⁻¹) and kanamycin (50 µg.mL⁻¹) at a final OD₆₀₀=0.1 and grown at
445 28°C 150 rpm until OD₆₀₀=1.0. One part was conserved for plant inoculation while the other was
446 centrifuged at top speed for 10 minutes and flash frozen for subsequent DNA extraction to serve as
447 control condition. Plants were grown in hydroponic systems as described above. After 5 days of growth,

448 each plant was inoculated with 1 mL Tn-seq library bacterial suspension at 5×10^7 cfu.mL⁻¹. The
449 experimentation was performed in triplicates with each replicate consisting of five plants. At 7 dpi, the
450 roots were harvested and placed in TE buffer.

451 DNA extraction and sequencing methods

452 The bacterial genomic DNA isolation using CTAB protocol (JGI) was used for the extraction of bacterial
453 DNA from the control and experimental conditions. For the latter, whole roots were used in the first
454 stages of the protocol. Whole roots were immersed in the extraction solution and vortexed for 5 min at
455 each step of the protocol to facilitate bacterial separation. Root grinding was avoided to prevent excessive
456 DNA contamination from the plant material. Roots were removed from the extraction buffer after the
457 lysis steps, before CTAB is added.

458 Ten µg of total DNA were digested with 1 µL of MmeI restriction enzyme (New England Biolabs) in 250
459 µL total volume with 10 µL of S-adenosyl methionine (SAM) 1.5 mM and 25 µL CutSmart buffer during
460 1 h at 37°C. Then, 1 µL of FastAP Thermosensitive Alkaline Phosphatase 1 U.µL⁻¹ (ThermoScientific)
461 was added and incubated 1h at 37°C. All enzymes were inactivated through a 5 min incubation at 75°C.
462 Digested DNA was column purified using a NucleoSpin Gel and PCR Clean-up kit (Macherey-Nagel). 2
463 µg of purified DNA was ligated to adapters containing specific barcode sequences. Adaptors were
464 obtained by annealing the primers 5'-TTCCCTACACGACGCTCTTCCGATCTXXXXNN-3' and 5'-
465 YYYYYAGATCGGAAGAGCGTCGTGTAGGGAAAGAGT-3' where NN are random nucleotides for
466 annealing to the dinucleotide overhang generated by MmeI and XXXXX and YYYYY are
467 complementary, barcode-specific sequences. Ligations were performed using 1 µL of 5 mM adapters, 1.5
468 µL of T4 DNA ligase 1 Weiss U.µL⁻¹ (ThermoScientific) and 2.5 µL 10x ligase buffer in a total volume
469 of 25 µL and incubated overnight at 16°C. 1 µL of ligation product was amplified using PCR with a GO
470 Taq DNA polymerase (Promega) and Illumina primers (P7 5'-
471 CAAGCAGAAGACGGCATAACGAGATAGACCGGGACTTATCATCCAACCTGT-3'; P5 5'-
472 AATGATACGGCGACCACCGAGATCTACACTCTTCCCTACACGACGCTCTTCCGATCT-3'). 22
473 PCR cycles were run (30 s at 92°C; 30 s at 60°C; 1 min at 72°C) with an initial step at 92°C for 2 min and
474 a final step at 72°C for 10 min. PCR products were subjected to a final gel purification using the
475 NucleoSpin Gel and PCR Clean-up kit (Macherey-Nagel).

476 Sequencing was performed at the high-throughput sequencing platform at I2BC (CNRS, Gif-sur-Yvette,
477 France) using Nextseq500 (Illumina) technology and 80 sequencing cycles (single read). Preliminary data
478 analysis (demultiplexing, trimming and mapping) was performed by the sequencing platform. Sequences
479 were trimmed from barcodes and mariner transposon sequences and deposited in European Nucleotide
480 Archive under Bioproject number PRJEB42565.

481 Tn-seq data analysis

482 Gene essentiality for optimal growth in a liquid LBm broth under agitation was assessed through a hidden
483 Markov model (HMM)-based analysis using the EL-ARTIST pipeline (window size of 7 TA
484 sites; *P* value of 0.03) (58). The pipeline predicts whether a TA site is located in an essential or non-
485 essential region. In some cases, the locus is predicted to be located in a region that contains both domains
486 that are required and dispensable for growth. After the first refinement, the HMM will recalculate the
487 transition probabilities based on the new data and repeat the run until the algorithm reaches convergence
488 (see **Supplemental dataset S6** for TA sites transposon insertion cover in each gene and condition).

489 For the identification of conditionally essential genetic regions, the sorted aligned sequences were first
490 normalized using TRANSIT 3.1.0 (59) with the Trimmed Total Reads method at default settings and an
491 additional LOESS correction. The normalized datasets were then analyzed using the TnseqDiff function
492 of the R package Tnseq (60). Adjusted *p*-values were calculated using the Benjamini & Hochberg
493 correction. Genetic loci were considered colonization depleted when their occurrence in the experimental
494 condition was at least 1.5-fold lower than in the control condition at a confidence level of $\text{padj} < 0.05$ and
495 inversely for colonization enriched genes.

496 Comparative genomics procedures

497 Core genome compositions were calculated using the Phyloprofile exploration tool implemented in the
498 MicroScope microbial genome annotation and analysis platform (61). Homology constraints were set at
499 $\text{minLrap} \geq 0.8$ and $\text{identity} \geq 50\%$. We used the COGNiTOR pre-computed COG category classification
500 available on MicroScope to infer COG category affiliations of *Bv* and *Pk* genes.

501 Construction of insertional mutants

502 For the production of the *cobW*, *pdeR* or *edd* insertional mutants of *B. vietnamiensis* LMG10929 and *P.*
503 *kururiensis* M130, a gene fragment of either *cobW*, *pdeR* and *edd* of each strain was amplified using
504 specific primers (**Supplementary Table S1**). Genes fragments were inserted into the target vector
505 pSHAFT2 (Shastri et al., 2017) carrying a chloramphenicol (Cm) resistance gene using an XbaI/XhoI
506 double digestion. Insert and plasmid were mixed at a 5:1 ratio with regard to their respective molecular
507 size, ligated using a T4 DNA ligase (Promega) and cloned into heat-shock competent *E. coli* DH5 α cells.
508 Positive cells were multiplied and their plasmid extracted using the Wizard Minipreps DNA purification
509 system (Promega). Plasmids were electroporated into WT *B. vietnamiensis* LMG10929 and *P. kururiensis*
510 M130 strains. Transformed cells were transferred in LBm for 3h at 30°C and then spotted on LBm plates
511 containing Cm (200 $\mu\text{g} \cdot \text{ml}^{-1}$) for selection of mutants. Plasmid insertion in each targeted gene was

512 checked by PCR. Mutants and WT strains showed identical growth rates in broth LBm or Hoagland
513 medium supplemented with glucose (not shown).

514

515 Acknowledgments

516 The authors acknowledge the IRD itrop HPC (South Green Platform; <https://bioinfo.ird.fr> -
517 <http://www.southgreen.fr>) at IRD Montpellier for providing HPC resources that have contributed to
518 the research results reported here. We acknowledge the High-throughput sequencing facility of I2BC
519 for its sequencing and bioinformatics expertise. The LABGeM (CEA/Genoscope & CNRS
520 UMR8030), the France Génomique and French Bioinformatics Institute national infrastructures are
521 acknowledged for support within the MicroScope annotation platform.

522 Funding

523 The authors gratefully acknowledge financial support from the CGIAR research program (CRP) RICE as
524 well as from the French national research agency (ANR) funding the BURKADAPT project (ANR-19-
525 CE20-0007). AW, JL and LG were supported by PhD fellowships from the French Ministry of Higher
526 Education, Research and Innovation.

527 References

- 528 1. Levy A, Gonzalez IS, Mittelviehhaus M, Clingenpeel S, Paredes SH, Miao J, Wang K, Devescovi G, Lebeis
529 S, Jin Z, McDonald M, Klein AP, Meghan E. 2018. Genomic features of bacterial adaptation to plants. *Nat*
530 *Genet* 50:138–150.
- 531 2. Compant S, Nowak J, Coenye T, Clement C, Ait Barka E. 2008. Diversity and occurrence of *Burkholderia*
532 spp. in the natural environment. *FEMS Microbiol Rev* 32:607–626.
- 533 3. Suárez-moreno ZR, Caballero-mellado J, Coutinho BG. 2012. Common features of environmental and
534 potentially beneficial plant-associated *Burkholderia*. *Microb Ecol* 63:249–266.
- 535 4. Dias GM, de Sousa Pires A, Grilo VS, Castro MR, de Figueiredo Vilela L, Neves BC. 2019. Comparative
536 genomics of *Paraburkholderia kururiensis* and its potential in bioremediation, biofertilization, and
537 biocontrol of plant pathogens. *Microbiologyopen* 8:e00801.
- 538 5. Wallner A, King E, Ngonkeu ELM, Moulin L, Béna G. 2019. Genomic analyses of *Burkholderia*
539 *cenocepacia* reveal multiple species with differential host-adaptation to plants and humans. *BMC Genomics*
540 20:803.
- 541 6. King E, Wallner A, Rimbault I, Barrachina C, Klonowska A, Moulin L, Czernic P. 2019. Monitoring of rice
542 transcriptional responses to contrasted colonizing patterns of phytobeneficial *Burkholderia* s.l. reveals a
543 temporal shift in JA systemic response. *Front Plant Sci* 10:1141.

- 544 7. Tr n Van V, Berge O, Ng  K  S, Balandreau J, Heulin T. 2000. Repeated beneficial effects of rice
545 inoculation with a strain of *Burkholderia vietnamiensis* on early and late yield components in low fertility
546 sulphate acid soils of Vietnam. *Plant Soil* 218/2:273–284.
- 547 8. Mattos KA, P dua VLM, Romeiro A, Hallack LF, Neves BC, Ulisses TMU, Barros CF, Todeschini AR,
548 Previato JO, Mendon a-Previato L. 2008. Endophytic colonization of rice (*Oryza sativa* L.) by the
549 diazotrophic bacterium *Burkholderia kururiensis* and its ability to enhance plant growth. *An Acad Bras*
550 *Cienc* 80:477–493.
- 551 9. Baldani V, Oliveira E, Balota E, Baldani J, Kirchhof G, Dobereiner J. 1997. *Burkholderia brasiliensis* sp.
552 nov., uma nova esp cie de bact ria diazotr fica endof tica. *An Acad Bras Cienc* 69.
- 553 10. Bournaud C, de Faria SM, dos Santos JMF, Tisseyre P, Silva M, Chaintreuil C, Gross E, James EK, Prin Y,
554 Moulin L. 2013. *Burkholderia* species are the most common and preferred nodulating symbionts of the
555 Piptadenia Group (Tribe Mimoseae). *PLoS One* 8:e63478.
- 556 11. Angus AA, Agapakis CM, Fong S, Yerrapragada S, Estrada-de los Santos P, Yang P, Song N, Kano S,
557 Caballero-Mellado J, de Faria SM, Dakora FD, Weinstock G, Hirsch AM. 2014. Plant-associated symbiotic
558 *Burkholderia* species lack hallmark strategies required in mammalian pathogenesis. *PLoS One* 9:e83779.
- 559 12. Estrada-de los Santos P, Palmer M, Ch vez-Ram rez B, Beukes C, Steenkamp E, Briscoe L, Khan N, Maluk
560 M, Lafos M, Humm E, Arrabit M, Crook M, Gross E, Simon M, dos Reis Junior F, Whitman W, Shapiro N,
561 Poole P, Hirsch A, Venter S, James E. 2018. Whole genome analyses suggests that *Burkholderia* sensu lato
562 contains two additional novel genera (*Mycetohabitans* gen. nov., and *Trinickia* gen. nov.): Implications for
563 the evolution of diazotrophy and nodulation in the Burkholderiaceae. *Genes (Basel)* 9:389.
- 564 13. Eberl L, Vandamme P. 2016. Members of the genus *Burkholderia*: good and bad guys. *F1000Research*
565 5:1007.
- 566 14. Coutinho BG, Licastro D, Mendon a-Previato L, C mara M, Venturi V. 2015. Plant-Influenced gene
567 expression in the rice endophyte *Burkholderia kururiensis* M130. *Mol Plant-Microbe Interact* 28:10–21.
- 568 15. Klonowska A, Melkonian R, Mich  L, Tisseyre P, Moulin L. 2018. Transcriptomic profiling of
569 *Burkholderia phymatum* STM815, *Cupriavidus taiwanensis* LMG19424 and *Rhizobium mesoamericanum*
570 STM3625 in response to *Mimosa pudica* root exudates illuminates the molecular basis of their nodulation
571 competitiveness and symbiotic evolutionary history. *BMC Genomics* 19:1–22.
- 572 16. Edwards J, Johnson C, Santos-Medell n C, Lurie E, Podishetty NK, Bhatnagar S, Eisen JA, Sundaresan V,
573 Jeffery LD. 2015. Structure, variation, and assembly of the root-associated microbiomes of rice. *Proc Natl*
574 *Acad Sci U S A* 112:E911–E920.
- 575 17. Shenton M, Iwamoto C, Kurata N, Ikeo K. 2016. Effect of wild and cultivated rice genotypes on rhizosphere
576 bacterial community composition. *Rice* 9:42.
- 577 18. Zhang J, Liu YX, Zhang N, Hu B, Jin T, Xu H, Qin Y, Yan P, Zhang X, Guo X, Hui J, Cao S, Wang X,
578 Wang C, Wang H, Qu B, Fan G, Yuan L, Garrido-Oter R, Chu C, Bai Y. 2019. NRT1.1B is associated with
579 root microbiota composition and nitrogen use in field-grown rice. *Nat Biotechnol* 37:676–684.
- 580 19. van Opijnen T, Bodi KL, Camilli A. 2009. Tn-seq: High-throughput parallel sequencing for fitness and

- 581 genetic interaction studies in microorganisms. *Nat Methods* 6:767–772.
- 582 20. Cole BJ, Feltcher ME, Waters RJ, Wetmore KM, Mucyn TS, Ryan EM, Wang G, Ul-Hasan S, McDonald
583 M, Yoshikuni Y, Malmstrom RR, Deutschbauer AM, Dangl JL, Visel A. 2017. Genome-wide identification
584 of bacterial plant colonization genes. *PLOS Biol* 15:e2002860.
- 585 21. Royet K, Parisot N, Rodrigue A, Gueguen E, Condemine G. 2019. Identification by Tn-seq of *Dickeya*
586 *dadantii* genes required for survival in chicory plants. *Mol Plant Pathol* 20:287–306.
- 587 22. Wheatley RM, Ford BL, Li L, Aroney STN, Knights HE, Ledermann R, East AK, Ramachandran VK, Poole
588 PS. 2020. Lifestyle adaptations of *Rhizobium* from rhizosphere to symbiosis. *Proc Natl Acad Sci U S A*
589 117:23823–23834.
- 590 23. Liu Z, Beskrovnaya P, Melnyk RA, Hossain SS, Khorasani S, O’Sullivan LR, Wiesmann CL, Bush J,
591 Richard JD, Haney CH. 2018. A genome-wide screen identifies genes in rhizosphere-associated
592 *Pseudomonas* required to evade plant defenses. *MBio* 9:e00433-18.
- 593 24. Sivakumar R, Ranjani J, Vishnu US, Jayashree S, Lozano GL, Miles J, Broderick NA, Guan C,
594 Gunasekaran P, Handelsman J, Rajendhran J. 2019. Evaluation of inseq to identify genes essential for
595 *Pseudomonas aeruginosa* PGPR2 corn root colonization. *G3 Genes, Genomes, Genet* 9:651–661.
- 596 25. Gutierrez MG, Yoder-Himes DR, Warawa JM. 2015. Comprehensive identification of virulence factors
597 required for respiratory melioidosis using Tn-seq mutagenesis. *Front Cell Infect Microbiol* 5:78.
- 598 26. Moule MG, Spink N, Willcocks S, Lim J, Guerra-Assunção JA, Cia F, Champion OL, Senior NJ, Atkins
599 HS, Clark T, Bancroft GJ, Cuccui J, Wren BW. 2016. Characterization of new virulence factors involved in
600 the intracellular growth and survival of *Burkholderia pseudomallei*. *Infect Immun* 84:701–710.
- 601 27. Wong Y-C, Abd El Ghany M, Naeem R, Lee K-W, Tan Y-C, Pain A, Nathan S. 2016. Candidate essential
602 genes in *Burkholderia cenocepacia* J2315 identified by genome-wide TraDIS. *Front Microbiol* 7:1288.
- 603 28. Gislason AS, Turner K, Domaratzki M, Cardona ST. 2017. Comparative analysis of the *Burkholderia*
604 *cenocepacia* K56-2 essential genome reveals cell envelope functions that are uniquely required for survival
605 in species of the genus *Burkholderia*. *Microb Genomics* 3:e000140.
- 606 29. Grazziotin AL, Vidal NM, Venancio TM. 2015. Uncovering major genomic features of essential genes in
607 Bacteria and a methanogenic Archaea. *FEBS J* 282:3395–3411.
- 608 30. Gil R, Silva FJ, Peretó J, Moya A. 2004. Determination of the core of a minimal bacterial gene Set.
609 *Microbiol Mol Biol Rev* 68:518–537.
- 610 31. Helmann TC, Deutschbauer AM, Lindow SE. 2019. Genome-wide identification of *Pseudomonas syringae*
611 genes required for fitness during colonization of the leaf surface and apoplast. *Proc Natl Acad Sci U S A*
612 116:18900–18910.
- 613 32. Suzuki K, Okazaki K, Tawaraya K, Osaki M, Shinano T. 2009. Gas chromatography–mass spectrometry
614 associated global analysis of rice root exudates under aseptical conditions. *Soil Sci Plant Nutr* 55:505–513.
- 615 33. Chavarría M, Nikel PI, Pérez-Pantoja D, De Lorenzo V. 2013. The Entner-Doudoroff pathway empowers
616 *Pseudomonas putida* KT2440 with a high tolerance to oxidative stress. *Environ Microbiol* 15:1772–1785.
- 617 34. Cumming RC, Andon NL, Haynes PA, Park M, Fischer WH, Schubert D. 2004. Protein disulfide bond

- 618 formation in the cytoplasm during oxidative stress. *J Biol Chem* 279:21749–21758.
- 619 35. Streit WR, Entcheva P. 2003. Biotin in microbes, the genes involved in its biosynthesis, its biochemical role
620 and perspectives for biotechnological production. *Appl Microbiol Biotechnol* 61:21-31.
- 621 36. Salaemae W, Booker GW, Polyak SW. 2016. The role of biotin in bacterial physiology and virulence: a
622 novel antibiotic target for *Mycobacterium tuberculosis*. *Microbiol Spectr* 4.
- 623 37. Selkrig J, Belousoff MJ, Headey SJ, Heinz E, Shiota T, Shen HH, Beckham SA, Bamert RS, Phan MD,
624 Schembri MA, Wilce MCJ, Scanlon MJ, Strugnell RA, Lithgow T. 2015. Conserved features in TamA
625 enable interaction with TamB to drive the activity of the translocation and assembly module. *Sci Rep* 5:1–
626 12.
- 627 38. Richter AM, Fazli M, Schmid N, Shilling R, Suppiger A, Givskov M, Eberl L, Tolker-Nielsen T. 2019. Key
628 players and individualists of Cyclic-di-GMP signaling in *Burkholderia cenocepacia*. *Front Microbiol*
629 10:3286.
- 630 39. Peekhaus N, Conway T. 1998. What's for dinner?: Entner-Doudoroff metabolism in *Escherichia coli*. *J*
631 *Bacteriol* 180:3495–3502.
- 632 40. Crouzet J, Levy-Schil S, Cameron B, Cauchois L, Rigault S, Rouyez MC, Blanche F, Debussche L, Thibaut
633 D. 1991. Nucleotide sequence and genetic analysis of a 13.1-kilobase-pair *Pseudomonas denitrificans* DNA
634 fragment containing five cob genes and identification of structural genes encoding Cob(I)alamin
635 adenosyltransferase, cobyrinic acid synthase, and bifunctional cobinamide kinase-cobinamide phosphate
636 guanylyltransferase. *J Bacteriol* 173:6074–6087.
- 637 41. Govindarajan M, Balandreau J, Muthukumarasamy R, Revathi G, Lakshminarasimhan C. 2006. Improved
638 yield of micropropagated sugarcane following inoculation by endophytic *Burkholderia vietnamiensis*. *Plant*
639 *Soil* 280:239–252.
- 640 42. Gillis M, Van Van T, Bardin R, Goor M, Hebbar P, Willems A, Segers P, Kersters K, Heulin T, Fernandez
641 MP. 1995. Polyphasic taxonomy in the genus *Burkholderia* leading to an emended description of the genus
642 and proposition of *Burkholderia vietnamiensis* sp. nov. for N₂-fixing isolates from rice in Vietnam. *Int. J.*
643 *Syst. Bacteriol.*, 45:274-289.
- 644 43. Mahenthiralingam E, Bischof J, Byrne SK, Radomski C, Davies JE, Av-Gay Y, Vandamme P. 2000. DNA-
645 Based diagnostic approaches for identification of *Burkholderia cepacia* complex, *Burkholderia*
646 *vietnamiensis*, *Burkholderia multivorans*, *Burkholderia stabilis*, and *Burkholderia cepacia* genomovars I and
647 III. *J Clin Microbiol* 38:3165–3173.
- 648 44. Tang S-Y, Hara S, Melling L, Goh K-J, Hashidoko Y. 2010. *Burkholderia vietnamiensis* isolated from root
649 tissues of nipa palm (*Nypa fruticans*) in Sarawak, Malaysia, proved to be its major endophytic nitrogen-
650 fixing bacterium. *Biosci Biotechnol Biochem* 74:1972–1975.
- 651 45. Raux E, Schubert HL, Warren MJ. 2000. Biosynthesis of cobalamin (vitamin B12): a bacterial conundrum.
652 *Cell Mol Life Sci C* 2000 5713 57:1880–1893.
- 653 46. Lertpiriyapong K, Gamazon ER, Feng Y, Park DS, Pang J, Botka G, Graffam ME, Ge Z, Fox JG. 2012.
654 *Campylobacter jejuni* type VI secretion system: Roles in adaptation to deoxycholic acid, host cell adherence,

- 655 invasion, and in vivo colonization. PLoS One 7: e42842.
- 656 47. Yang F, Xue D, Tian F, Hutchins W, Yang CH, He C. 2019. Identification of c-di-GMP signaling
657 components in xanthomonas oryzae and their orthologs in xanthomonads involved in regulation of bacterial
658 virulence expression. Front Microbiol 10:1402.
- 659 48. Yang F, Tian F, Sun L, Chen H, Wu M, Yang C-H, He C. 2012. A novel two-omponent system PdeK/PdeR
660 regulates c-di-GMP turnover and virulence of *Xanthomonas oryzae* pv. *oryzae*. Mol Plant Microbe
661 Interact 25:1361-9.
- 662 49. Knights HE, Jorin B, Haskett TL, Poole PS. 2021. Deciphering bacterial mechanisms of root colonization.
663 Environ Microbiol Rep. 13:428-444.
- 664 50. Do Amaral FP, Tuleski TR, Pankiewicz VCS, Melnyk RA, Arkin AP, Griffiths J, Tadra-Sfeir MZ, Maltempi
665 de Souza E, Deutschbauer A, Monteiro RA, Stacey G. 2020. Diverse bacterial genes modulate plant root
666 association by beneficial bacteria. MBio 11:1–15.
- 667 51. Kim HJ, Nam HS, Anderson AJ, Yang KY, Cho BH, Kim YC. 2007. Mutation in the edd gene encoding the
668 6-phosphogluconate dehydratase of *Pseudomonas chlororaphis* O6 impairs root colonization and is
669 correlated with reduced induction of systemic resistance. Lett Appl Microbiol 44:56–61.
- 670 52. Cheng X, Etalo DW, van de Mortel JE, Dekkers E, Nguyen L, Medema MH, Raaijmakers JM. 2017.
671 Genome-wide analysis of bacterial determinants of plant growth promotion and induced systemic resistance
672 by *Pseudomonas fluorescens*. Environ Microbiol 19:4638–4656.
- 673 53. Chanch T, Ohira K. 1982. Comparison of the uptake and assimilation of ammonium and nitrate in indica
674 and japonica rice plants using the tracer ¹⁵N method. Soil Sci Plant Nutr 28:79–90.
- 675 54. Cianfanelli FR, Monlezun L, Coulthurst SJ. 2016. Aim, load, fire: the type VI secretion system, a bacterial
676 nanoweapon. Trends Microbiol 24:51-62.
- 677 55. Vergunst AC, Meijer AH, Renshaw SA, O’Callaghan D. 2010. Burkholderia cenocepacia creates an
678 intramacrophage replication niche in zebrafish embryos, followed by bacterial dissemination and
679 establishment of systemic infection. Infect Immun 78:1495–508.
- 680 56. Ferrières L, Hémerly G, Nham T, Guérout AM, Mazel D, Beloin C, Ghigo JM. 2010. Silent mischief:
681 Bacteriophage Mu insertions contaminate products of *Escherichia coli* random mutagenesis performed using
682 suicidal transposon delivery plasmids mobilized by broad-host-range RP4 conjugative machinery. J
683 Bacteriol 192:6418–6427.
- 684 57. Wiles TJ, Norton JP, Russell CW, Dalley BK, Fischer KF, Mulvey MA. 2013. Combining quantitative
685 genetic footprinting and trait enrichment analysis to identify fitness determinants of a bacterial pathogen.
686 PLoS Genet 9:e1003716.
- 687 58. Pritchard JR, Chao MC, Abel S, Davis BM, Baranowski C, Zhang YJ, Rubin EJ, Waldor MK. 2014.
688 ARTIST: High-resolution genome-wide assessment of fitness using transposon-insertion sequencing. PLoS
689 Genet 10:e1004782.
- 690 59. DeJesus MA, Ambadipudi C, Baker R, Sasseti C, Ioerger TR. 2015. TRANSIT - A software tool for
691 Himar1 TnSeq analysis. PLoS Comput Biol 11.

- 692 60. Zhao L, Anderson MT, Wu W, Harry HL, Bachman MA. 2017. TnseqDiff: Identification of conditionally
693 essential genes in transposon sequencing studies. *BMC Bioinformatics* 18:326.
- 694 61. Médigue C, Calteau A, Cruveiller S, Gachet M, Gautreau G, Josso A, Lajus A, Langlois J, Pereira H, Planel
695 R, Roche D, Rollin J, Rouy Z, Vallenet D. 2019. MicroScope-an integrated resource for community
696 expertise of gene functions and comparative analysis of microbial genomic and metabolic data. *Brief*
697 *Bioinform* 20:1071-1084.
- 698

699 Tables

700 Table 1. Distribution of genes significantly impacting bacterial fitness on IR64 and Nipponbare rice.
701 Colonization events on IR64 and Nipponbare rice cultivars were analyzed separately.

702 Table 2. Selection of functions and number of corresponding genes impacting rice colonization
703 efficiencies when mutated. The detailed list of genes is provided in Table S3.

704 Table 3. Selection of functions and number of corresponding colonization-depleted genes on IR64 and
705 Nipponbare rice.

706 Table 4. Bacterial strains and plasmids used in this study.

707

708 Figures

709 **Figure 1. Distribution of genes significantly impacting bacterial fitness in a rich liquid medium.** (A)
710 Number of genes predicted to be essential, domain-essential (only insertions in part of the gene lead to
711 abundance decrease) or non-essential by the EL-ARTIST pipeline. (B) Principal component analysis of
712 essential and non-essential genes of *Bv* and *Pk* in a rich liquid medium. The genes were classified
713 according to their COG categories and the PCA was built in regard to the abundance of these categories.

714 **Figure 2. Rice colonization efficiencies for *Bv* and *Pk*.** Nipponbare and IR64 rice seedlings were
715 inoculated at 5 days post germination with 1.10^7 bacterial cells, total root systems were harvested at
716 different time points and the associated bacterial population was estimated ($p < 0.05$, Wilcoxon test). The
717 boxplots were generated using 5 replicates for each condition. Outliers are represented as black dots.

718 **Figure 3. Distribution of genes significantly impacting bacterial fitness during rice colonization.**
719 Colonization events on IR64 and Nipponbare rice cultivars were pooled for this analysis. (A) Transposon
720 insertions in genes leading to a 1.5-fold or higher decrease in abundance on roots at $\text{padj} < 0.05$ were
721 identified to be colonization- depleted. Those leading to a 1.5-fold or higher abundance were identified to

722 be colonization-enriched. (B) Distribution of genes significantly depleted or enriched during rice
723 colonization along COG categories in ratio of total impacted genes.

724 **Figure 4. Distribution of colonization-depleted genes along the core- and specific-genomes of *Bv* and**
725 ***Pk*.** There are a total of 2072 genes that present significant homologies in the genomes of *Bv* and *Pk* and
726 thus considered part of the core-genome (Materials and methods). A fraction of this core-genome is
727 colonization-depleted in both species. Additionally, a substantial proportion of the genes that are found to
728 be colonization depleted in a species or the other also belong to the core-genome.

729 **Figure 5. Distribution of colonization-depleted genes on IR64 and Nipponbare rice cultivars.**
730 Distribution along COG categories of: (A) core genes of *Bv* and *Pk* that are colonization-depleted on both
731 rice cultivars, (B) genes that are specifically colonization-depleted in *Bv* and (C) genes that are
732 specifically colonization-depleted in *Pk*.

733 **Figure 6. Colonization capacity of *Bv* and *Pk* insertion mutants in competition assays.** Three genes,
734 *cobW*, *edd* and *pdeR*, predicted by the Tn-seq approach to be involved in rice root colonization, were
735 selected for targeted disruption in *Bv* and *Pk*. Mutants and WT strains were inoculated simultaneously on
736 Nipponbare rice roots and enumerated at 7 dpi. Significance levels of pairwise comparisons were
737 estimated using a Wilcoxon test ($p < 0.05$). For each mutant, the log₂ FC value observed in the Tn-seq
738 approach on Nipponbare is displayed. Positive and negative correlations with the mutagenesis approach
739 are expressed with green and red squares respectively.

740 **Figure 7. Summary diagram of colonization-impacted bacterial functions involved in efficient rice**
741 **root colonization.** Pathways, functions, and genes described throughout this work are synthesized on this
742 integrative representation. Elements specific of *Bv* are placed in the top part of the schematic bacteria
743 (pink) and *Pk* specific elements at the bottom (green). Elements that are commonly found in both bacteria
744 are placed at the interface (white). Function and molecular systems which were detected to be
745 colonization-depleted are written in red font whereas colonization-enriched genes are written in blue font.

746

Table 1. Distribution of genes significantly impacting bacterial fitness on IR64 and Nipponbare rice.

Strain	Colonization impact	Common	IR64	Nipponbare	Total
<i>B. vietnamiensis</i> LMG10929	Enriched	507	83	130	720
	Depleted	635	225	134	994
	Total	1142	308	264	1714
<i>P. kururiensis</i> M130	Enriched	48	57	36	141
	Depleted	374	98	50	522
	Total	422	155	86	663

Table 2. Selection of functions and number of corresponding genes impacting rice colonization efficiencies when mutated.

Function	<i>B. vietnamiensis</i>		<i>P. kururiensis</i>	
	Depleted	Enriched	Depleted	Enriched
Motility	35	-	35	-
Chemotaxis	4	5	9	1
Cobalamin synthesis (B12)	1	8	15	-
Thiamin synthesis (B1)	4	-	5	-
Biotin synthesis (B7)	5	-	-	-
Potassium uptake	5	-	5	-
Ribose metabolism	-	5	-	-
Entner Doudoroff pathway	1	-	2	-
Aromatic compound metabolism	5	-	-	-
c-di-GMP cycling	1	7	-	1
Putrescine metabolism	3	-	1	-
Putrescine uptake	-	5	-	-
Cellulose synthase	-	5	-	-
Cell wall integrity	5	-	10	-
Peptidoglycan synthesis	3	-	-	-
Lytic transglycolases	4	-	1	-
Tol-Pal system	5	-	3	-
Hopanoid synthesis	7	-	1	-
O-antigen synthesis	2	-	-	-
Osmoprotection	2	-	1	-
DNA repair	9	-	9	-
Flp/Tad pilus	-	12	-	-
Arginine biosynthesis	5	-	4	-
Histidine biosynthesis	9	-	9	-
Isoleucine biosynthesis	6	-	4	-
Leucine biosynthesis	3	-	4	-
Lysine biosynthesis	-	-	1	-
Methionine biosynthesis	3	-	5	-
Proline biosynthesis	2	-	2	-
Serine biosynthesis	2	-	2	-
Tryptophan biosynthesis	6	-	6	-
Purine metabolism	9	1	1	-
Pyrimidine metabolism	5	-	6	-
Queuosine synthesis	6	-	1	-
Quorum sensing	-	-	-	2
T2SS	9	-	8	-
Autotransporter assistance	-	-	2	-
Prophage	-	-	-	22

Table 3. Selection of functions and number of corresponding colonization-depleted genes on IR64 and Nipponbare rice.

Function	<i>B. vietnamiensis</i>		<i>P. kururiensis</i>	
	IR64	Nipponbare	IR64	Nipponbare
T6SS	10	-	-	-
Glycerol metabolism	3	-	-	4
Phosphate import	3	-	-	-
Ammonium transport (AmtB)	-	1	-	1
Oxoprolinase	-	3	-	-
O-antigen biosynthesis	1	-	3	-

Table 4. Bacterial strains and plasmids used in this study.

Strains/plasmids	Relevant characteristics and plasmid constructions	Reference/Source
<i>E. coli</i>		
MFDpir	Donor strain containing the pSAM-Ec vector and auxotroph for diaminopimelate, $\Delta dapA::(erm-pir)$	(Ferrières et al., 2010)
<i>B. vietnamiensis</i>		
LMG10929	Wild-type strain	(Gillis et al., 1995)
Rif ^R	Spontaneous rifampicin resistance clone of LMG10929	This study
Spt ^R	Spontaneous spectinomycin resistance clone of LMG10929	This study
DsRed	LMG1029 + pIN29	(King et al., 2019)
$\Omega cobW$	Insertion mutant of pSHAFT2 in AK36_2246	This study
$\Omega pdeR$	Insertion mutant of pSHAFT2 in AK36_4666	This study
Ωedd	Insertion mutant of pSHAFT2 in AK36_178	This study
<i>P. kururiensis</i>		
M130	Wild-type strain	(Baldani et al., 1997)
Rif ^R	Spontaneous rifampicin resistance clone of M130	This study
Spt ^R	Spontaneous spectinomycin resistance clone of M130	This study
DsRed	M130 + pIN29	(King et al., 2019)
$\Omega cobW$	Insertion mutant of pSHAFT2 in ANSKv1_10390	This study
$\Omega pdeR$	Insertion mutant of pSHAFT2 in ANSKv1_51226	This study
Ωedd	Insertion mutant of pSHAFT2 in ANSKv1_70910	This study
Plasmids		
pIN29	Carries genes conferring DsRed fluorescence and chloramphenicol resistance. DsRed is under control of the highly active pTAC promoter	(Vergunst et al., 2010)
pSAM-Ec	Contains Himar1C9 transposase and a mariner transposon with a kanamycin resistance cassette	(Wiles et al., 2013)
pSHAFT2	Vector used for insertional mutagenesis containing a chloramphenicol resistance gene (non-replicative plasmid)	(Shastri et al., 2017)

Fig.1

A

ARTIST prediction	<i>B. vietnamiensis</i> LMG10929	<i>P. kururiensis</i> M130
Essential	661	638
Domain-essential	96	97
Non-essential	5845	5638
Unclear	112	77




Fig.2





Fig.3

A



B




Fig.4




Fig.5






Fig.6



Colonization
 Depleted
 Enriched



○ Ile
 * Trp, Pro, Met,
 Leu, His, Arg
 + Ser, Lys
 □ B1, B12 (Pk)
 B1, B12 (Bv)