



DNA metabarcoding of passive trap collection media for forest insect biomonitoring

Lucas Sire, Paul Schmidt Yáñez, Annie Bézier, Béatrice Courtial, Susan Mbedi, Sarah Sparmann, Rodolphe Rougerie, Christophe Bouget, Michael T Monaghan, Elisabeth A. Herniou, et al.

► To cite this version:

Lucas Sire, Paul Schmidt Yáñez, Annie Bézier, Béatrice Courtial, Susan Mbedi, et al.. DNA metabarcoding of passive trap collection media for forest insect biomonitoring. 6e Colloque de Génomique Environnementale. "Environmental and Agronomical Genomics Symposium", Sep 2021, Tours, France. , p.49, 2021, Environmental and Agronomical Genomics 2021 symposium. Book of Abstracts. hal-03750402

HAL Id: hal-03750402

<https://hal.inrae.fr/hal-03750402>

Submitted on 12 Aug 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.




FRANCE
GÉNOMIQUE



GDR
GÉNOMIQUE
ENVIRONNEMENTALE

Environmental And Agronomical Genomics Symposium

27th -29th October 2021
Tours (France)



Book of Abstracts



Institut de Recherche
sur la Biologie de l'Insecte



<https://eags-2021.sciencesconf.org/>



@fr_genomics
#FG_EAGS2021

DNA metabarcoding of passive trap collection media for forest insect biomonitoring

Lucas Sire^{*1}, Paul Schmidt Yáñez², Annie Bézier¹, Béatrice Courtial³, Susan Mbedi⁴, Sarah Sparmann⁴, Rodolphe Rougerie⁵, Christophe Bouget⁶, Michael T. Monaghan^{2,7}, Elisabeth Herniou¹, and Carlos Lopez-Vaamonde^{1,3}

¹Institut de recherche sur la biologie de l'insecte UMR7261 – Université de Tours, Centre National de la Recherche Scientifique : UMR7261 – Av Monge 37200 TOURS, France

²Leibniz Institute of Freshwater Ecology and Inland Fisheries – Berlin-FriedrichshagenNeuglobsowBerlin-AdlershofBerlin Center for Genomics in Biodiversity Research (BeGenDiv), Germany

³INRAE UR0633 Zoologie Forestière – INRAE, UR063 Zoologie Forestière - Orléans, France – Orléans, France, France

⁴Berlin Center for Genomics in Biodiversity Research (BeGenDiv) – Berlin, Germany, Germany

⁵Institut de Systématique, Evolution, Biodiversité – Université des Antilles, Museum National d'Histoire Naturelle, Ecole Pratique des Hautes Etudes, Université Paris sciences et lettres, Centre National de la Recherche Scientifique : UMR7205, Sorbonne Université – 57 rue Cuvier - CP 50 - 75005 Paris, France

⁶Ecosystèmes forestiers – Institut National de Recherche pour l'Agriculture, l'Alimentation et l'Environnement : UR1455 – Domaine des Barres, F-45290 Nogent-sur-Vernisson, France

⁷Freie Universität Berlin – Kaiserswerther Str. 16-18, 14195 Berlin, Germany

Abstract

Insect decline has been increasingly reported in the past years due to global change. Large-scale biomonitoring has thus become necessary to better understand the dynamics of insect communities and to preserve their essential role in ecosystem functioning. In that sense, coupling high-throughput sequencing and DNA metabarcoding has exponentially increased our potentiality to monitor insect communities over wider geographic regions and time scales. However, biomonitoring of entomofauna using molecular tools often results in destructive DNA extraction through voucher grinding, impeding primordial morphological back-up. Here, we filter unprocessed collection medium to assess insect communities through environmental DNA metabarcoding. We demonstrate that recovered communities are different yet complementary and that insect response to environmental changes remains similar to homogenate bulk metabarcoding. We also show that insect orders-by their contrasting sclerotization ratio-, and collection medium type, are unequal in yielding metabarcoding results. Overall, we believe it as an efficient alternative for biomonitoring insect response to ecological changes while preserving insect vouchers for identification and description, especially in tropical regions were singlettons or undescribed species can be very common in trap samples.

^{*}Speaker