



HAL
open science

Contribution of omics to biopreservation: Toward food microbiome engineering

Frédéric Borges, Romain Briandet, Cécile Callon, Marie-Christine Champomier-Vergès, Souad Christieans, Sarah Chuzeville, Catherine Denis, Nathalie Desmasures, Marie-Hélène Desmonts, Carole Feurer, et al.

► To cite this version:

Frédéric Borges, Romain Briandet, Cécile Callon, Marie-Christine Champomier-Vergès, Souad Christieans, et al.. Contribution of omics to biopreservation: Toward food microbiome engineering. *Frontiers in Microbiology*, 2022, 13, pp.1-16. 10.3389/fmicb.2022.951182 . hal-03751737

HAL Id: hal-03751737

<https://hal.inrae.fr/hal-03751737>

Submitted on 15 Aug 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License



OPEN ACCESS

EDITED BY

Jerome Combrisson,
Mars (United States), United States

REVIEWED BY

Minakshi Prasad,
Lala Lajpat Rai University of Veterinary
and Animal Sciences, India
Christian U. Riedel,
University of Ulm, Germany
Silvina Graciela Fadda,
CONICET Centro de Referencia para
Lactobacilos (CERELA), Argentina

*CORRESPONDENCE

Frédéric Borges
frederic.borges@univ-lorraine.fr

SPECIALTY SECTION

This article was submitted to
Food Microbiology,
a section of the journal
Frontiers in Microbiology

RECEIVED 23 May 2022

ACCEPTED 14 July 2022

PUBLISHED 02 August 2022

CITATION

Borges F, Briandet R, Callon C,
Champomier-Vergès M-C,
Christieans S, Chuzeville S, Denis C,
Desmases N, Desmonts M-H,
Feurer C, Leroi F, Leroy S, Mounier J,
Passerini D, Pilet M-F,
Schlüsselhuber M, Stahl V, Strub C,
Talon R and Zagorec M (2022)
Contribution of omics
to biopreservation: Toward food
microbiome engineering.
Front. Microbiol. 13:951182.
doi: 10.3389/fmicb.2022.951182

COPYRIGHT

© 2022 Borges, Briandet, Callon,
Champomier-Vergès, Christieans,
Chuzeville, Denis, Desmases,
Desmonts, Feurer, Leroi, Leroy,
Mounier, Passerini, Pilet,
Schlüsselhuber, Stahl, Strub, Talon and
Zagorec. This is an open-access article
distributed under the terms of the
[Creative Commons Attribution License
\(CC BY\)](https://creativecommons.org/licenses/by/4.0/). The use, distribution or
reproduction in other forums is
permitted, provided the original
author(s) and the copyright owner(s)
are credited and that the original
publication in this journal is cited, in
accordance with accepted academic
practice. No use, distribution or
reproduction is permitted which does
not comply with these terms.

Contribution of omics to biopreservation: Toward food microbiome engineering

Frédéric Borges^{1*}, Romain Briandet², Cécile Callon³,
Marie-Christine Champomier-Vergès², Souad Christieans⁴,
Sarah Chuzeville⁵, Catherine Denis⁶, Nathalie Desmases⁷,
Marie-Hélène Desmonts⁸, Carole Feurer⁹, Françoise Leroi¹⁰,
Sabine Leroy¹¹, Jérôme Mounier¹², Delphine Passerini¹⁰,
Marie-France Pilet¹³, Margot Schlüsselhuber⁷, Valérie Stahl⁸,
Caroline Strub¹⁴, Régine Talon¹¹ and Monique Zagorec¹³

¹Université de Lorraine, LIBio, Nancy, France, ²Université Paris-Saclay, INRAE, AgroParisTech, Micalis Institute, Jouy-en-Josas, France, ³Université Clermont Auvergne, INRAE, VetAgro Sup, UMR 545 Fromage, Aurillac, France, ⁴ADIV, Clermont-Ferrand, France, ⁵ACTALIA, Pôle d'Expertise Analytique, Unité Microbiologie Laitière, La Roche sur Foron, France, ⁶ACTALIA, Sécurité des Aliments, Saint Lô, France, ⁷Normandie Univ, UNICAEN, UNIROUEN, ABTE, Caen, France, ⁸Aerial, Illkirch, France, ⁹IFIP, Institut de la Filière Porcine, Le Rheu, France, ¹⁰Ifremer, MASAE, Laboratoire EM3B, Nantes, France, ¹¹Université Clermont Auvergne, INRAE, MEDIS, Clermont-Ferrand, France, ¹²Univ Brest, Laboratoire Universitaire de Biodiversité et Ecologie Microbienne, Plouzané, France, ¹³Oniris, INRAE, SECALIM, Nantes, France, ¹⁴Qualisud, Univ Montpellier, Avignon Université, CIRAD, Institut Agro, IRD, Université de La Réunion, Montpellier, France

Biopreservation is a sustainable approach to improve food safety and maintain or extend food shelf life by using beneficial microorganisms or their metabolites. Over the past 20 years, omics techniques have revolutionised food microbiology including biopreservation. A range of methods including genomics, transcriptomics, proteomics, metabolomics and meta-omics derivatives have highlighted the potential of biopreservation to improve the microbial safety of various foods. This review shows how these approaches have contributed to the selection of biopreservation agents, to a better understanding of the mechanisms of action and of their efficiency and impact within the food ecosystem. It also presents the potential of combining omics with complementary approaches to take into account better the complexity of food microbiomes at multiple scales, from the cell to the community levels, and their spatial, physicochemical and microbiological heterogeneity. The latest advances in biopreservation through omics have emphasised the importance of considering food as a complex and dynamic microbiome that requires integrated engineering strategies to increase the rate of innovation production in order to meet the safety, environmental and economic challenges of the agri-food sector.

KEYWORDS

shelf life, food, safety, pathogen, spoilage, microbiome, fermentation, biopreservation

Introduction

Foods of animal or plant origins are complex ecosystems, rich in nutrients, with physicochemical characteristics enabling microbial growth during processing and storage. These ecosystems are colonised by microbial communities that can include pathogenic or spoilage microorganisms but also beneficial ones. The consumption of food contaminated with pathogens is an important cause of morbidity and mortality worldwide. Every year, approximately 600 million people – 1 in 10 people – get sick from foodborne pathogens, 420,000 of whom die. Human damage caused by foodborne pathogens results in colossal economic losses amounting to USD 110 billion due to lost productivity and health expenses (World Health Organization, 2015). Spoilage organisms are responsible for colour, odour, texture, taste or packaging defects leading to inedible products. Food microbial contaminants (spoilage as well as pathogenic microorganisms) contribute to global food loss and waste. According to the Food and Agriculture Organization (FAO), food waste occurs during the retail and consumption stages while food loss occurs after harvest or slaughter until retail (FAO, 2019). The causes of food waste and loss are numerous but a significant part of food destruction linked to microbial contamination is due to non-compliance with pathogen-related regulations or spoilage. At the European Union level, 20% of total available food is lost or wasted, fruits and vegetables being the most impacted category (43.5% of the food group), ahead of meat and fish products (26.3%; Caldeira et al., 2019). Worldwide, one-third of food produced for human consumption, about 1.3 billion tonnes per year, is estimated to be lost or wasted along the food supply chain (HPLE, 2014), while about 12% of the world population suffers from hunger (Agriculture and Economic Development Analysis Division, 2013). Reducing food wastage is thus crucial not only for ethical reasons but also for economic reasons. Food loss and waste are responsible for direct costs of about USD one trillion every year, but hidden costs extend much further. Indeed, global costs (including environmental, social, and economic costs) are evaluated by the FAO to amount to USD 2.6 trillion per year (FAO, 2014).

In addition, as food waste is correlated with high greenhouse gas emissions, reducing the undesirable impact of microorganisms has become a major objective in a situation of climate emergency. In this context, microorganisms also offer potential levers of action and can represent real opportunities for resolving food safety issues (Cavicchioli et al., 2019). In the global food system, approaches based on the barrier properties of biological systems appear attractive because of their efficiency and sustainability. These approaches are grouped under various names including biosanitation, biocontrol, bioprotection, and biopreservation. In this review, we will refer to food biopreservation, which is based on the hurdle technology that consists in using microorganisms (often lactic acid bacteria)

as protective cultures and/or their metabolites to optimise the microbiological quality and shelf life of food by ensuring safety or reducing food waste, as defined by Stiles (Stiles, 1996). The use of protective cultures is often considered as an alternative to chemical additives or as a replacement for certain ingredients. Therefore, biopreservation should also help to meet the strong expectations of consumers who want “healthier” and more “natural” foods, and contribute to nutritional recommendations aimed at reducing salts, sugars, and additives in foods.

The concept of biopreservation was inspired by food fermentation ancestrally used to preserve food, except that fermentation involves substantial transformation of the food matrix, which is usually not the intention when engineering biopreservation systems. Consistently, the intentional addition of microorganisms or their metabolites for specific preservation purposes was largely investigated on fermented food, i.e., dairy products (cheeses, yoghurt), bakery products, or fermented sausages. Nevertheless, biopreservation was successfully extended to non-fermented food such as seafood, raw meat and non-fermented plant products. For example, biopreservation of seafood as fresh fish fillets, smoked fish or cooked shrimps aimed at controlling spoilers or pathogens such as *Listeria monocytogenes*, *Staphylococcus aureus*, *Vibrio* and histamine-producing bacteria (for a recent review, see Rathod et al., 2022). Biopreservation of raw meat (lamb, pork, beef, or poultry) and processed meat (sausage, cooked ham) has also focused on the control of pathogenic and spoilage bacteria as well as extension of shelf life (Jones et al., 2010; Zagorec and Champomier-Vergès, 2017; Wang et al., 2019). Biopreservation of non-fermented plant products is essentially dedicated to fighting against spoilage microorganisms including yeasts, moulds, spore-forming bacteria (*Bacillus subtilis*, *Bacillus licheniformis*), and pathogenic bacteria such as *L. monocytogenes* (Leyva Salas et al., 2017).

Most species used for food biopreservation by the food industry are lactic acid bacteria belonging essentially to the genera *Lactococcus*, *Lactobacillus lato sensu* and *Carnobacterium*. One strain of the Gram-negative species *Hafnia alvei* is also available on the market for anti-*Escherichia coli* purposes (Callon et al., 2016; Frétin et al., 2020). These bacteria are derived from food microbiota and are therefore particularly well adapted to food matrices. Moreover, as these species have been studied for several decades, their use as protective cultures is considered safe. *Latilactobacillus sakei*, *Latilactobacillus curvatus*, *Carnobacterium divergens*, *Carnobacterium maltaromaticum* and *Lactococcus lactis* are the main species providing protective cultures for meat/seafood products (Leroi et al., 2015; Comi et al., 2016; Ramarosan et al., 2018; Iacumin et al., 2020), while lactobacilli or even yeast strains can be used for vegetable food biopreservation (Siedler et al., 2019; Truchado et al., 2020; Windholtz et al., 2021). Although several biopreservation technologies are already available on the market, the rate and speed of innovation

in this area needs to increase considerably in order to meet global climate-related challenges. Indeed, the scientific literature dealing with biopreservation is considerably larger than the actual application of biopreservation in the food sector. Foods are complex systems because of their diversity, their various physical, chemical, and biological structures, and the numerous processes used to produce them. Moreover, food microbial community dynamics during shelf life depends on abiotic parameters, which are mostly linked to production processes or storage conditions, and biotic parameters where microbial interactions play a major role. The complexity and diversity of food microbial communities has been a major barrier to the widespread use of biopreservation. Therefore, the conception of efficient biopreservation technology requires the implementation of methodological approaches adapted to the high complexity of food ecosystems. Food microbiology has long been studied by classical culture-dependent methods. During the last decade, omics techniques including genomics, transcriptomics, proteomics, metabolomics, culturomics, and phenomics have revolutionised all areas of the life sciences, including food quality and safety assessment, because of their ability to decipher food systems as a whole (Cook and Nightingale, 2018). The application of omics provides a more realistic portrait of the complex interactions occurring in the food ecosystem (Gálvez et al., 2007), and it has significantly increased our understanding of the potential of microbiomes to increase the productivity and sustainability of food systems. These omics approaches have caused a paradigm shift from unsocial undesirable microorganisms colonising food to strongly interacting microorganisms establishing stable networks (Berg et al., 2020). Omics approaches have been applied to explore various aspects of biopreservation, such as selection and characterisation of protective cultures, investigation of the mechanisms involved in the protective effect, or the impact on food microbial communities. After a brief overview of food biopreservation, the different omics approaches used in studies dealing with biopreservation are reviewed below and analysed by illustrating to what extent they can answer questions related to the impact of biopreservation on the food microbial ecosystem.

Selection of biopreservation agents

Classical approaches to identify protective culture candidates are mainly based on the detection of inhibition zones in laboratory conditions. Recently, the use of phenomics and genomics has proved highly effective. Phenomics can be defined as the high-throughput study of phenotypes. In the case of biopreservation, the phenotype of interest is the inhibition of spoilage microorganisms or food-borne

pathogens. Phenomics has been successfully used to identify strains exhibiting remarkable anti-*L. monocytogenes* properties by using a high-throughput liquid handling system and a genetically engineered luminescent strain of *L. monocytogenes* to set up mixed culture competition assays in food matrices (Riedel et al., 2007; El Kheir et al., 2018). This method was first used to select anti-*Listeria* candidates from a collection of strains isolated from raw milk and a collection of *Lactococcus piscium*. The majority of the candidates obtained did not produce an inhibitory halo following a classical agar diffusion-based method, suggesting promising inhibitory mechanisms (El Kheir et al., 2018). Lately, this high-throughput competition assay was implemented to study the inhibition phenotype of a collection of *C. maltaromaticum* strains under multiple varying conditions. This method resulted in the selection of robust antagonistic *C. maltaromaticum* strains whose anti-*Listeria* properties are insensitive to fluctuations, i.e., inoculation level and time lag of *L. monocytogenes* and candidate inoculation (Borges and Revol-Junelles, 2019). It is expected that phenomics approaches extended to other target microorganisms (Besnard et al., 2021), as well as other phenotypes related to the sensory profile or use of nutrients (Wiernasz et al., 2017; Acin-Albiac et al., 2020), may in the future enable the identification of microorganisms with high biopreservation performances.

Genomics is an interesting approach for selecting protective cultures as genome mining may point out important features that can be involved in the preservation effect (Baltz, 2019), and also prove the absence of some unwanted functions such as antibiotic resistance or biogenic amine synthesis. Genome mining involves the analysis of functional gene annotation, resulting in particular from antiSMASH (Blin et al., 2017) and BAGEL (van Heel et al., 2018), which are designed to identify clusters of genes involved in the biosynthesis of antimicrobial compounds, combined with genome comparison to find correlation between the presence/absence of genes and protective properties. As an example, the genome sequence of *L. sakei* 23K, a meat adapted bacterium used as a starter for sausage fermentation, but also proposed as a biopreservative agent for raw meat products (Zagorec and Champomier-Vergès, 2017), revealed its strong ability to be competitive in meat products (Chaillou et al., 2005; Eijsink and Axelsson, 2005). Indeed, genome analysis highlighted the presence in the genome of elements putatively enabling the use of alternative carbon sources, such as ribose, inosine, and adenosine. Their efficacy was subsequently proven and helps to explain the fitness in meat of *L. sakei*, which thereby escapes competition for energy sources (Rimaux et al., 2011). Also, the requirement for haem and iron, two components present in meat, could be assessed by genomics through the gene repertoire and further evidenced by functional genomics (Duhutrel et al., 2010; Verplaetse et al., 2020).

Another example of genomics input is the production of antagonistic molecules by protective strains. This has long been studied through the production of bacteriocins. The

genome sequence analysis of *L. curvatus* CRL705, a strain known to produce two bacteriocins (lactocin 705 and AL705), revealed the presence of additional genes putatively involved in bacteriocin production (sakacin P, sakacin Q, sakacin X, and sakacin T; Hebert et al., 2012). Divercin V41 is a bacteriocin involved in the protective function of *C. divergens* V41 a lactic acid bacterium strain whose operon sequence was reported more than two decades ago (Metivier et al., 1998). The genome analysis of this strain revealed that an additional gene was present in the divercin V41 operon (Remenant et al., 2016), the function of which was shown to be important for bacteriocin production (Back et al., 2015). Comparative genomics of *Carnobacterium* highlighted potential new candidate strains for biopreservation, efficient against *L. monocytogenes* and harbouring original bacteriocin gene equipment. For example, five different bacteriocins and a 16 kDa new one were predicted in the *C. maltaromaticum* SF668 and *C. maltaromaticum* EBP3019 genomes, respectively (Begrem et al., 2020). Combining genome analysis and peptidomics of a *Companilactobacillus crustorum* strain enabled, from the peptides produced during the growth of this strain, the discovery of eight novel bacteriocins and two other antimicrobials with a broad spectrum of action against Gram-positive and Gram-negative pathogens (Yi et al., 2018).

L. piscium CNCM I-4031 (EU2241) is a protective strain for seafood products that improves the sensory quality of shrimp and cold-smoked salmon (Fall et al., 2012; Leroi et al., 2015). This strain is also particularly efficient against the pathogen *L. monocytogenes* by reducing growth and virulence (Saraoui et al., 2018). Combined phenotyping and genome analyses evidenced that the inhibitory effect is dependent on cell-to-cell contact instead of extracellular molecules such as bacteriocins, organic acids, or hydrogen peroxide (Saraoui et al., 2016; Marché et al., 2017). This unusual mechanism still remains to be elucidated; nevertheless medium- and high-throughput screening revealed that other strains of the same species could be selected as new protective cultures, notably related to their large antimicrobial capacities (Wiernasz et al., 2017; El Kheir et al., 2018).

Genomics can reveal other unexpected features, as exemplified by the genome sequence of *C. divergens* V41 which contains an intriguing long genomic island (~40 kb) encoding polyketide synthases/non-ribosomal peptide synthases (PKS/NRPS) or PKS/NRPS-like enzymes, putatively involved in the production of a secondary metabolite of unknown function (Remenant et al., 2016). Such molecules may have antimicrobial functions or be associated with oxidative stress resistance, and immunomodulatory or cytotoxicity activities. Comparative genomic analysis of *Carnobacterium* strains showed that this PKS/NRPS gene cluster is unique in *C. divergens* V41 (Begrem et al., 2020). Other PKS/NRPs antimicrobial compounds, such as milkisin produced by a *Pseudomonas* sp. strain, with potentially interesting properties for the biopreservation of

milk products have been described (Schlüsselhuber et al., 2018, 2020). Reuterin-producing *Limosilactobacillus reuteri* are known as bioprotective agents for dairy products (Ortiz-Rivera et al., 2017). Some rare strains harbour a PKS/NRPS genomic island encoding *rtc* genes involved in the synthesis, regulation, immunity, and secretion of an antimicrobial tetramic acid named reutericyclin (Gänzle et al., 2000; Lin et al., 2015).

While many published papers describe the screening and evaluation of bioprotective antifungal lactic acid bacteria strains *in vitro* and *in situ* (Delavenne et al., 2012; Leyva Salas et al., 2018), as well as the identification of metabolites involved in their antifungal activities, the application of genomics and functional genomics to the selection of antifungal microorganisms is still in its infancy. Studies providing insights into the metabolic pathways of antifungal metabolites such as phenyllactic acid (Wu et al., 2020), clearly point out that in-depth studies of antifungal lactic acid bacteria using comparative and functional genomics are needed. Such studies should help to elucidate yet uncharacterised biosynthetic pathways of known antifungal molecules and potentially reveal new ones and to establish whether other competition exclusion phenomena exist between lactic acid bacteria and spoilage fungi, with the objective to further understand their action mechanism and possibly provide helpful tools for strain selection.

Some species with bioprotective properties are also described as potential spoilage organisms, depending on the type of food or process (Brillet et al., 2004; Leisner et al., 2007; Andreevskaya et al., 2015; Leroi et al., 2015; Saraoui et al., 2016; Poirier et al., 2018). Thus comparative genomics between strains known to be responsible for spoilage or on the contrary known as protective cultures should be an interesting approach for the selection of strains of interest. In addition, as a complementary approach to phenotypic tests, genome analysis of potent protective strains can be a complementary tool for their safety assessment i.e., screening for the presence of genes related to biogenic amine production, antibiotic resistance genes, as well as their location with respect to mobile genetic elements, i.e., plasmids and bacteriophages. Such an approach was recently applied to a *Lactiplantibacillus plantarum* starter culture used in the manufacturing of *nahm* fermented pork (Chokesajjawatee et al., 2020), as well as to a potent *L. plantarum* protective culture (Barbosa et al., 2021).

Impact of biopreservation on targeted microorganisms

The role of protective cultures is mainly related to their fitness (nutritional competition, ability to resist harsh conditions encountered in food), enabling them to establish and dominate at the expense of other undesirable species, as well as to their ability to produce antimicrobial molecules,

TABLE 1 Main characteristics of biopreservation in food and mechanisms involved.

Protective mode of action	Resulting effect	Effect at cellular and/or molecular levels	Mode of use	References
Nutritional competition	<ul style="list-style-type: none"> - Jameson effect - Growth impairment because of lack of nutrients 	<ul style="list-style-type: none"> - Early entry into stationary phase of targeted microorganisms, and protective cultures - Growth cessation of targeted microorganisms 	Live microorganisms added to the food	Jameson, 1962; Guillier et al., 2008; Hibbing et al., 2010
Production of organic acids	<ul style="list-style-type: none"> - Extracellular pH drop - Diffusion across the microbial cytoplasmic membrane 	<ul style="list-style-type: none"> - Cytoplasmic pH decrease - Collapse of the proton gradient across the membrane - Disruption of cellular processes 		Warnecke and Gill, 2005
Production of hydrogen peroxide	<ul style="list-style-type: none"> - Oxidation of cellular components 	<ul style="list-style-type: none"> - Peroxidation and disruption of membrane layers - Oxidation of oxygen scavengers - Enzyme inhibition - Oxidation of nucleosides - Disruption of protein synthesis - Growth decrease at low concentrations - Cell death at high concentrations 		Finnegan et al., 2010
Respiration	Change in atmosphere composition (O ₂ decrease) leading to microaerophilic conditions	<ul style="list-style-type: none"> - Inhibition of strict aerobic bacteria 	Live microorganisms added to the food	Ben Said et al., 2019
Production of bacteriocins	Bactericidal or bacteriostatic activity against species taxonomically related to the producing strain	<ul style="list-style-type: none"> - Pore formation in the cytoplasmic membrane - Loss of structure and subsequent cell death 	Live microorganisms added to the food Bacteriocins purified from cultivated producer strains	Elsser-Gravesen and Elsser-Gravesen, 2013 Martinez et al., 2016


such as organic acids, hydrogen peroxide or specialised metabolites as bacteriocins, biosurfactants, or lipopeptides (Ben Said et al., 2019; Bourdichon et al., 2021; Table 1). The expected impact is induction of decay or at least growth inhibition of the undesirable microorganism. Classical methods such as qRT-PCR are *a priori* methods that involve a prior selection of the target genes to be studied. On the contrary, omics approaches such as RNAseq can be used without *a priori*, opening the possibility to identify original unsuspected mechanisms. These approaches have revealed that the mechanisms involved are more complex than previously thought, and involve microbial sensing capacity, gene regulation, and the potential for combining antimicrobial compounds for microbial inactivation (Figure 1).

Listeria monocytogenes

Two recent studies have explored the mechanism of inhibition of *L. monocytogenes* by antimicrobial lipopeptide (P34) or the peptide nisin (encapsulated or not) by proteomic analysis after incubation in a laboratory medium (Pinilla et al., 2021; Stincone et al., 2021). The lipopeptide P34 caused the downregulation of proteins involved in manganese transport and upregulation of proteins related to iron transport in *L. monocytogenes*. In addition, reduction of stress tolerance proteins related to the sigma B and VirR regulons

and modulation of phosphoenolpyruvate phosphotransferase systems for sugar transport were observed (Stincone et al., 2021). Exposure of *L. monocytogenes* to nisin induced the synthesis of proteins related to ATP-binding cassette transporter systems, transmembrane proteins, RNA-binding proteins, and diverse stress response proteins (Pinilla et al., 2021). Some of the proteins detected in the presence of free nisin were related to translocation of *L. monocytogenes* virulence factors, activation of the LiaR-mediated stress defence, and glycosylation of cell wall teichoic acid. The comparison of treatment by free and encapsulated nisin revealed that *L. monocytogenes* did not express some stress proteins when nisin was encapsulated, suggesting the production of nisin-resistance factors by exposure to encapsulated nisin. The authors suggested that liposomes allow controlled release of nisin in the medium, resulting in fewer interactions between nisin and bacteria compared with free nisin, which may impact the mechanism of action of nisin (Pinilla et al., 2021). The induction of resistance factors was also observed by exposing *L. monocytogenes* to sublethal doses of pediocin. Transcriptomic analysis revealed the expression modulation of the two-component system LisRK and the alternative sigma factors SigB and SigL, resulting in an increased resistance of *L. monocytogenes* to this bacteriocin (Laursen et al., 2015).

Recently, the ability of *C. maltaromaticum*, *Leuconostoc gelidum*, or *L. piscium* strains to inhibit *L. monocytogenes* was demonstrated in seafood (Saraoui et al., 2016, 2018;



Wiernasz et al., 2017). *L. piscium* inhibition required cell-to-cell contact with *L. monocytogenes*, affecting its cell surface, and decreasing its virulence (Saraoui et al., 2018). The metabolomic fingerprints suggested that this inhibition might not involve nutritional competition and remains to be explored (Saraoui et al., 2016).

Staphylococcus aureus

Lactococcus garvieae was shown to inhibit *S. aureus* growth in milk, in cheese, and *in vitro*, potentially through hydrogen peroxide (H_2O_2) production (Delbes-Paus et al., 2010). To better characterise this mechanism of inhibition *in vitro*, the transcriptomes of *L. garvieae* and *S. aureus* co-culture have been explored by RNA-seq and RT-qPCR (Delpuch et al., 2015, 2017). *L. garvieae* repressed the expression of *S. aureus* genes involved in stress response, including oxidative stress generated by H_2O_2 , and in cell division. It also modulated the expression of virulence-related genes (particularly *agrA*, *hld*, and enterotoxin-encoding genes; Delpuch et al., 2015). For *L. garvieae*, a high concentration of H_2O_2 was not associated with higher expression of the H_2O_2 synthesis genes *pox*, *sodA*, and *spxA1*, but rather with repression of H_2O_2 -degradation genes (*trxB1*, *ahpC*, *ahpF*, and *gpx*; Delpuch et al., 2017). The interaction between *L. lactis* and *S. aureus* has also been widely studied and transcriptomic analyses were performed with

microarrays and RT-qPCR. In a chemically defined medium held at a constant pH value of 6.6, the growth of the two bacteria in co-culture was not modified, but their transcriptome was modulated (Even et al., 2009; Nouaille et al., 2009). The expression of *S. aureus* virulence-related genes was impaired by *L. lactis*: the expression of genes encoding global regulators, including *agr* and consequently the *agr*-controlled enterotoxin genes and *sar*, was strongly reduced (Even et al., 2009). This downregulation of *agr* in *S. aureus* was associated with the reducing properties of *L. lactis* (Nouaille et al., 2014). *L. lactis* genes associated with amino acid metabolism, ion transport, oxygen response, menaquinone metabolism, cell surface, and phage expression were differentially expressed in co-culture compared to monoculture (Nouaille et al., 2009). In a complex medium such as the cheese matrix, the acidifying, proteolytic, and reducing activities of *L. lactis* were shown to affect carbohydrate and nitrogen metabolisms and the stress response of *S. aureus* (Cretenet et al., 2011). Enterotoxin gene expression was positively or negatively modulated by both *L. lactis* and the cheese matrix itself, depending on the enterotoxin type. Again, the *agr* operon was downregulated by the presence of *L. lactis*, in part because of a drop in pH (Cretenet et al., 2011). All these data highlight the intimate link between environment, metabolism, and virulence expression. A third binary interaction between *Enterococcus faecalis* and *S. aureus* was studied in milk and in cheese (Viçosa et al., 2018; Nogueira Viçosa et al., 2019). When

co-cultured, the growth of *S. aureus* was decreased and the classical enterotoxins were not produced. The expression of several enterotoxins and global regulator genes (including *agr*) was downregulated, while the expression of genes involved in metabolism was upregulated. Finally, the interaction of *S. aureus* with a mixed culture of *Enterococcus durans*, *E. faecalis* and *L. lactis* in milk confirmed that the production of enterotoxins was reduced in mixed culture and the expression of several genes involved in virulence was inhibited (Zdenkova et al., 2016). All these studies on the interaction of *S. aureus* with different lactic acid bacteria converge on a very important idea that the mechanism of action of biopreservation can result in the inhibition of virulence by inhibiting the production of enterotoxins through the decreased expression of genes involved in their synthesis.

Pathogenic *Escherichia coli*

L. curvatus, *L. plantarum*, and *Enterococcus mundtii* strains can inhibit the growth of *E. coli* O157:H7 when co-cultured in a meat model medium (Orihuel et al., 2018). The antagonistic effect of the most efficient *E. mundtii* strains against *E. coli* O157:H7 were characterised by a proteomic analysis (Orihuel et al., 2018). The expression of *E. mundtii* proteins involved in carbohydrate/amino acid metabolisms, energy production, transcription/translation, and cell division was modified in the presence of *E. coli*. Reciprocally, the presence of *E. mundtii* resulted in repression of *E. coli* synthesis of proteins related to metabolism and transport of amino acids and nucleotides, as well as overexpression of proteins involved in stress, energy production, and transcription (Orihuel et al., 2019). In addition, proteins associated with adhesion to extracellular matrix proteins of meat were modulated in *E. coli* in accordance with its decreased adhesion capacity when co-cultured with *E. mundtii*. *E. mundtii* did not influence the lytic cycle of the *E. coli* O157:H7 strain, indicating its potentially safe use as a bioprotective agent, since engagement in the lytic cycle results in the production of shiga toxin (Orihuel et al., 2019). The interaction of *E. coli* O157:H7 with ground beef microbiota was also studied (Galia et al., 2017). A beef piece was divided in two parts with the inner part considered sterile and the outer part as encompassing a natural meat microbiota. The microbial community structure was assessed by 16S rDNA amplicon sequencing, and the transcriptome of two inoculated strains (*E. coli* O157:H7 and *E. coli* O26:H11) was studied by RNAseq comparing samples of sterile and naturally contaminated meat. This study revealed that the two *E. coli* strains behave differently. On the one hand, an upregulation of genes involved in detoxification and stress response and a downregulation of *peR*, a gene negatively associated with virulence phenotype, were observed in *E. coli* O157:H7. On the other hand, the interaction of *E. coli* O26:H11 with ground beef microbiota revealed that

genes involved in division, peptidoglycan synthesis, DNA repair, metal acquisition, and carbohydrate and amino acid metabolism were downregulated (Galia et al., 2017).

Fungal food spoilers

In a recent review of biopreservation against moulds in dairy products (Shi and Maktabdar, 2022), the authors point out newly described antifungal mechanisms. Among these is a perfect example of how omics technologies shed new light on the understanding of the protective mechanisms of antifungal lactobacilli toward dairy product spoilage fungi (Siedler et al., 2019). This work revealed that manganese scavenging by *Lactocaseibacillus rhamnosus* and *Lactocaseibacillus paracasei* antifungal strains, previously known as a defence mechanism against oxidative stress, was a main inhibitory mechanism (i.e., competitive exclusion) against many yeast and mould species involved in dairy product spoilage. Indeed, following milk fermentation and supplementation with manganese, their bioactivity was completely lost. A transcriptomic approach based on RNA-seq further showed that one of the most highly expressed gene products in these strains encoded a manganese transporter (MntH1). The role of MntH1 in manganese scavenging was confirmed in a $\Delta mntH1$ *L. paracasei* strain in which no significant antifungal activity was detected, while bioactivity was restored in the $\Delta mntH1$ mutant complemented with a plasmid containing the *mntH1* gene under its own promoter.

Besides the above-mentioned discovery that competitive exclusion for manganese was an important antifungal mechanism, production of antifungal metabolites and pH decrease were believed to be the main mechanisms involved in the bioactivity of antifungal protective lactic acid bacteria. Through the use of metabolomic targeted and untargeted approaches with or without prior medium fractionation and bioactivity testing, more than 60 molecules have been thought to play a role in antifungal activity (see recent reviews by Leyva Salas et al., 2017 and Siedler et al., 2019). These metabolites include molecules produced through carbohydrate metabolism (e.g., organic acids such as lactic, acetic, formic, and succinic acids, volatile compounds such as diacetyl), proteolysis (e.g., bioactive peptides resulting from casein cleavage), amino acid metabolism (e.g., phenyllactic acid), lipolysis and free fatty acid metabolism (e.g., 3-hydroxydecanoic, caproic – i.e., hexanoic- and caprylic – i.e., octanoic – acids), but also complex compounds derived from bioconversions (e.g., benzoic acid) or peptide synthesis [e.g., cyclic dipeptides such as cyclo(L-Phe-L-Pro) and cyclo(L-Phe-trans-4-OH-L-Pro)] (Ström et al., 2002; Aunsbjerg et al., 2015; McNair et al., 2018; Leyva Salas et al., 2019; Garnier et al., 2020; Shi and Knöchel, 2021a,b). It should be underlined that with the exception of lactic and acetic acids which are produced in g/kg or g/L

amounts, these molecules are produced in quite low quantities, all of which are at concentrations far from their individual minimum inhibitory concentration (MIC), thus suggesting that they act in synergy or by additive effects. In a recent study, Leyva Salas et al. (2019) used a metabolomics approach coupled with supervised multivariate analysis to investigate 56 antifungal compounds as well as volatiles. It was found that 9 key compounds including acetic acid, 5 aromatic acids, and three volatiles were associated with antifungal activity against *Mucor racemosus* and *Penicillium commune*, although their concentrations were below their respective MICs. Further investigation on *Penicillium roqueforti* and *Mucor circinelloides* revealed that several combinations presented additive (e.g., diacetyl + 3-phenylpropanoic acid, diacetyl + acetic acid) or synergistic effects (diacetyl + octanoic acid, octanoic + 3-phenyl propanoic acids), clearly reinforcing the idea that additive and synergistic effects of antifungal molecules are involved in lactic acid bacteria bioactivity. To go further, future work could include the investigation of more complex mixtures of antifungal compounds at concentrations close to those encountered in biopreserved foods. Moreover, it is not clear how antifungal molecule synthesis and competitive exclusion interact together in different fungal species with various susceptibilities to protective strains.

Biopreservation at the microbiome level

The primary objective of biopreservation is to limit the presence of unwanted microorganisms in food. However, the addition of biopreservatives can have an overall impact on the food microbiome. Omics approaches have helped clarify the complex interactions occurring in the food ecosystem and their impact on the organoleptic properties of foods (Figure 1).

Metagenomics

Prior to the availability of high-throughput DNA sequencing (HTS) techniques, DNA fingerprinting techniques as PCR-denaturing gradient gel electrophoresis or PCR-temporal temperature gradient electrophoresis, targeting mainly the V3 region of 16S rDNA, were employed to determine whether bioprotective lactic acid bacteria strains were able to colonise the food matrix and dominate the microbiota (Hu et al., 2008; Saraoui et al., 2017; Zhang et al., 2018). However, discrepancies between these DNA fingerprinting approaches and cultural methods (Ercolini et al., 2010) have highlighted the need for methods of higher resolution. There are two commonly used HTS methods in microbiome research: amplicon sequencing and metagenomic sequencing (Liu et al., 2021). Amplicon sequencing is the most widely used,

including in the field of biopreservation. It demonstrated that the three protective cultures *L. rhamnosus* LRH05, *L. sakei* LSK04, and *C. maltaromaticum* CNB06, alone or in combination, were able to colonise cheese and became dominant after storage (Bassi et al., 2020). In shrimp, the results of amplicon sequencing were consistent with the successful colonisation of the food matrix by the biopreservative strains *L. plantarum* and *Lacticaseibacillus casei*, and a reduction in the relative abundance of *Shewanella*, which includes the spoilage species *Shewanella baltica* (Li et al., 2019). In addition, in beef burgers, the analysis of predicted metagenomes revealed that nisin-activated packaging resulted in a reduction in the abundance of specific metabolic pathways related to spoilage (Ferrocino et al., 2015). Table 2 summarises the different studies using omics approaches to assess the impact of biopreservation on food ecosystems and the main findings.

HTS approaches open up the possibility of going much further than simply answering the question of the implantation success of the protective microorganisms and their effect on the targets. Amplicon sequencing can be used to estimate the impact of biopreservation on food microbiota. When batches of cold-smoked salmon were inoculated with a bioprotective strain of *L. piscium*, the microbiota structure differed significantly between control and biopreserved products after 3 weeks of storage (Leroi et al., 2015). The impact of biopreservation strains can be stronger, as in the case of fermented sausages where 16S rRNA-based analysis revealed a markedly lower microbial diversity of the metabolically active microbial community (Giello et al., 2018). Thus, although special attention is paid to the selection of strains with a narrow antimicrobial spectrum in biopreservation (e.g., bacteriocin-producing bacteria), biopreservation can end up with significant changes in food microbiota structure. Although this side effect can be undesirable, especially in fermented foods, in some circumstances it is possible to take advantage of this broad impact on food microbiota to target spoilage microorganisms, which can encompass a large number of phylogenetic taxa. In dairy products such as low-salt fresh cheeses, spoilage bacteria are mainly psychrotrophic Gram-negative species, including several *Enterobacteriaceae* and *Pseudomonas* spp. (Ledenbach and Marshall, 2009; Spanu et al., 2018; Bassi et al., 2020). The three lactic acid bacterial strains *L. rhamnosus* LRH05, *L. sakei* LSK04, and *C. maltaromaticum* CNB06, added alone or in combination, were found after 5 weeks of storage to be effective in inhibiting the Gram-negative bacteria population of fresh Primo Sale cheese inoculated with a cocktail of 10 bacterial spoilage isolates (Bassi et al., 2020). In Italian fresh filled pasta cheese, the impact of protective cultures at the community level can be used to reduce the initial microbiota associated with raw materials and to confer a competitive advantage on safer or more acceptable bacterial species such as *Leuconostoc mesenteroides*, at the expense of more problematic species such as *Streptococcus uberis* and *Streptococcus parauberis* (Tabanelli et al., 2020).

TABLE 2 Overview of omics approaches used to assess the impact of biopreservation on food ecosystems and main findings.

Omics approach	Methodological details	Food	Main finding	References
PCR-Denaturing gradient gel electrophoresis/Temporal temperature gradient electrophoresis associated or not with band sequencing	V3 region of 16S rDNA	Cooked and peeled shrimp	<i>Carnobacterium divergens</i> V41 but not <i>Lactococcus piscium</i> CNCM I-4031 used to inoculate shrimp dominated and was associated with reduction of off-flavours	Saraoui et al., 2017
	V3 region of 16S rDNA	Vacuum-packaged beef meat	Bands associated with spoilage bacteria (<i>Pseudomonas</i> , Enterobacteriaceae, <i>Brochothrix thermosphacta</i>), but not with LAB, disappeared in samples inoculated with <i>Latilactobacillus sakei</i> and <i>Latilactobacillus curvatus</i> bioprotective strains	Zhang et al., 2018
	V3 region of 16S rDNA	Vacuum-packed cooked ham	Predominant spoilage LAB were not detected when the bioprotective <i>Latilactobacillus sakei</i> B-2 strain was used	Hu et al., 2008
	V6–V8 region of 16S rDNA	Beef cuts packaged in nisin-coated plastic bags	Similar diversity in control and nisin-treated samples although differences were observed with plate counts for <i>Brochothrix thermosphacta</i>	Ercolini et al., 2010
DNA sequencing	Pyrosequencing of V3–V4 region of 16S rDNA	Cold-smoked salmon	Different OTU ratios were observed between control and samples inoculated with <i>Lactococcus piscium</i> EU2241 (= CNCM I-4031). No correlation with sensory analysis	Leroi et al., 2015
	Illumina sequencing of V3–V4 region of 16S rDNA	Raw/peeled shrimp	<i>Shewanella baltica</i> significantly inhibited after co-inoculation with <i>Lactiplantibacillus plantarum</i> AB-1 and <i>Lactocaseibacillus casei</i> LC	Li et al., 2019
	Illumina sequencing of V3–V4 region of 16S rDNA	St Nectaire-type cheese	Implantation of an inhibitory consortium whose inhibitory activity toward <i>Escherichia coli</i> O26:H11 depended on indigenous microbiota composition	Frétin et al., 2020
	Illumina sequencing of V3–V4 region of 16S rDNA and of an internal 280 bp fragment of the <i>gyrB</i> gene	Diced cooked ham	Bioprotective activity and implantation of a nisin-producing strain of <i>Lactococcus lactis</i> depended on microbiota composition	Chaillou et al., 2022
16S rRNA sequencing	Illumina sequencing of V4 region of 16S rDNA	Fresh filled pasta	Cultures of <i>Lactiplantibacillus plantarum</i> and <i>Lactocaseibacillus paracasei</i> were not dominant but reduced the initial microbiota and gave a competitive advantage to other LAB species	Tabanelli et al., 2020
	Sequencing of V3–V4 region of 16S rRNA from cDNA	Fermented sausage	Large domination of <i>Lactobacillaceae</i> and reduction of bacterial diversity in samples inoculated with protective <i>Latilactobacillus curvatus</i> strain	Giello et al., 2018
	Sequencing of V3–V4 region of 16S rRNA from cDNA	Beef burgers in nisin-activated packaging	Lower abundance of some taxa in samples with nisin-activated packaging	Ferrocino et al., 2015
Volatilome analysis	Headspace SPME/GC-MS	Cooked and peeled tropical shrimp	Inhibition of <i>Brochothrix thermosphacta</i> by <i>Lactococcus piscium</i> CNCM I-4031 correlated with attenuation of off-odours and diminution of some volatile compounds	Fall et al., 2012
	Headspace SPME/GC-MS	Salmon gravlax	6 protective strains exhibited their own volatilome profiles. Quality improvement was not correlated with implantation of protective culture	Wiernasz et al., 2020
	Headspace SPME/GC-MS	Fresh filled pasta	<i>Lactocaseibacillus rhamnosus</i> and <i>Lactocaseibacillus paracasei</i> influenced the aroma profile with overall acceptability of the product	Tabanelli et al., 2020
	NMR spectroscopy	<i>in vitro</i>	Kinetic analysis of 11 major metabolites involved in the metabolism of <i>Lactocaseibacillus rhamnosus</i> and <i>Lactocaseibacillus plantarum</i>	Ebrahimi et al., 2016
	FTICR-MS	Red wines	No effect on the volatile compounds of a <i>Metschnikowia pulcherrima</i> bioprotective strain. Wines produced from bioprotected or sulphited must had different metabolic signatures	Simonin et al., 2020

While, on the one hand, biopreservation can have an impact on the microbiota, on the other hand, the microbiota can also affect the efficiency of biopreservation. The protective property of a nisin-producing *L. lactis* strain was tested in combination with high pressure under controlled conditions of microbiota composition in reduced-nitrite diced cooked ham (Chaillou et al., 2022). Sterile diced ham cubes were inoculated with two different microbiota collected from cooked hams, together with the protective *L. lactis* strain prior to vacuum packaging and high-pressure treatment. During storage, the two selected cooked ham microbiota were both characterised by a microbial community enriched in high potential spoilage bacterial species (especially *Proteobacteria*). Comparison of the bacterial community composition after 1 month of storage revealed that the protective effect of the *L. lactis*/high-pressure combination is highly dependent on the ham microbiota. Indeed, when ham samples were inoculated with *Pseudomonas* spp. and *Serratia* spp. rich-microbiota, *L. lactis* became dominant (>90% relative abundance). However, when ham samples were instead inoculated with microbiota dominated by *Psychrobacter* sp. and *Vibrio* sp., *L. lactis* was not competitive and *Brochothrix thermosphacta* became dominant (Chaillou et al., 2022). Thus, by interacting with the protective strain, food microbiota can act on the efficiency of biopreservation. By contrast, it is also possible that the microbiota can contribute to biopreservation in concert with the protective strains by acting directly on the unwanted microorganism through its intrinsic protective properties. In uncooked pressed cheese, the protective activity of a consortium comprising three strains belonging to the species *H. alvei*, *L. plantarum*, and *L. lactis* was dependent on the composition of the microbiota colonising the processed raw milk. The raw milk batches associated with the lowest growth of *E. coli* O26:H11 were characterised by greater relative abundance of lactic acid bacteria, the three *Gammaproteobacteria* *Acinetobacter*, *Serratia*, and *Hafnia*, as well as *Macrococcus*. On the other hand, the highest levels of *E. coli* O26:H11 were observed when the milk microbiota was significantly enriched with bacteria from the genera *Ramboutsia*, *Paeniclostridium*, and *Turicibacter* (Frélin et al., 2020).

Overall, these data were mainly produced thanks to amplicon sequencing. The major drawbacks of this approach are the biases in relative abundances resulting from PCR amplification and differences in the number of ribosomal operons between species (Edgar, 2017), the limited taxonomic resolution, mainly at the genus level even if some efforts have been made in developing specialised databases such as the DAIRYdb (Meola et al., 2019), and the lack of functional information. Amplicon sequencing targeting housekeeping genes such as *gyrB* is also promising for a better identification at the species or even intra-species level, with also less bias for relative abundance determination (Poirier et al., 2018). Investigating food communities by using shotgun metagenomics could help to improve accuracy, especially by

gathering information at the species or even the strain level, and could give a global view of the functions involved in the process of biopreservation at the microbial community scale. Furthermore, network analysis of metagenetic and metagenomic data could be used to identify patterns (i.e., co-occurrence and mutual exclusion) in food microbial communities, biopreserved or not. Such an approach would enable hypotheses to be drawn regarding biotic interactions occurring between microorganisms, which could then be tested experimentally, and could be applied to the selection of potential candidates for biopreservation or for a deeper understanding of the impact of selected bioprotective cultures at a community level.

Metabolomics

In addition to their impact on food microorganisms, biopreservation agents are likely to modify the properties of the food matrix and in particular its organoleptic characteristics. Metabolomics approaches allow us to go much further than sensory analyses in the study of the impact of biopreservation agents on the matrix. SPME/GC-MS showed that cooked peeled shrimp contains a reduced amount of unwanted aldehydes and alcohols associated with sensory spoilage when *L. piscium* is used as an inhibitor of *B. thermosphacta* (Fall et al., 2012). Biopreservatives can thus play a positive sensory role by inhibiting a target microorganism responsible for spoilage. However, biopreservation can also negatively impact the matrix and can be responsible for significant changes, highlighting the interest of the polyphasic omics approach to the selection of candidate strains. An exemplary study describes the use of Head Space-Solid Phase MicroExtraction/Gas Chromatography- Mass Spectrometry (HS-SPME/GC-MS) to reveal that the impact of biopreservation candidates varied dramatically depending on the strain considered, and that specific signatures could be associated with each strain. Coupled to microbial community structure investigation by amplicon sequencing, the authors were able to rationally select two strains with the highest protective effect and the lowest sensory activity in salmon gravlax (Wiernasz et al., 2020; Table 2). Nuclear Magnetic Resonance (NMR) also has great potential in the study of living organisms, owing to its non-destructive nature, i.e., it can be used for *in vivo* and *in vitro* measurements of biological processes, with no quenching of the metabolism required. NMR spectroscopy can be particularly helpful for the kinetic analysis of the metabolism of protective cultures, as in the development of *in vitro* NMR kinetic measurements of lactic acid bacteria (Ebrahimi et al., 2016; Table 2). Such a polyphasic approach was also used successfully in wine to assess biopreservation as an alternative to sulphites (Simonin et al., 2020). The wine metabolome being of high complexity, an ultra-high-resolution mass spectrometric

method -the Fourier transform ion cyclotron resonance mass spectrometry- was used to identify and annotate more than 7,000 molecules. Clustering analysis revealed that even if the biopreservation agent has a molecular impact on the product, it is significantly lower than the winery effect, showing that biopreservation has preserved the typicality of the products, which can be of high relevance for products with protected designation of origin (Simonin et al., 2020; Table 2).

Limits of global omics in structured food matrices

Genomic tools based on amplicon sequencing, although powerful for food microbial ecosystem description, may show some limits for biopreserved products. Bioprotective cultures inoculated at a high level are usually dominant, at least at the beginning of storage. As it is generally admitted that taxa representing less than 0.01% of the dominant ones are not detected with metagenetics, the richness of biopreserved food may be underestimated. In the gut, the use of shotgun metagenomics has revealed that the majority of species harbour multiple strains (Ellegaard and Engel, 2016). For instance, despite a low species complexity, the infant gut microbiota exhibits extensive intra-species diversity, with an average of 4.9 strains per subject (Luo et al., 2015). In food, low resolution is a limitation in estimating the impact of protective cultures on subdominant species, such as pathogens usually present at low levels ($<10^2$ CFU/g). This problem is less important for specific spoilage organisms that generally dominate at sensory rejection time. However, the interaction between bacteria can modify the sensory characteristics (Stohr et al., 2001; Joffraud et al., 2006; Silbände et al., 2018), and it may be that underestimated taxa play a role in the deterioration of food. In addition, most efforts have been focused on bacteria, and to a lesser extent on yeasts and moulds, although phages, viruses, or *Archaea* may also be present in food microbiota (Roh et al., 2010; Park et al., 2011).

Omics are powerful tools in analysing protective cultures and their interactions at the population level. However, most of these techniques hardly consider microbial population heterogeneity in food systems that can trigger important community functions. Diversification of cell types can originate from genetic variation, ageing, gene expression stochasticity, and environmental condition fluctuation (Bury-Moné and Sclavi, 2017). In structured food matrices such as ground meat or cheese, microorganism microenvironments are highly heterogeneous and vary over time depending upon local microbial activities (Ferrier et al., 2013; Jeanson et al., 2015). Above critical textural levels, individual cells are not able to sediment or swim inside the matrix and eventually grow as large 3D microcolonies (Darsonval et al., 2021; Saint Martin et al., 2022). Microcolony size and sphericity depend on several factors such as local rheological properties,

nutrient availability, competing microbiota, and associated interference interactions (Verheyen et al., 2018). Sharp gradients of nutrients and metabolites are generated inside and around the colonies, thus expanding the functional diversification of the local populations. In other structured biosystems such as bacterial biofilms (Lenz et al., 2008; Pérez-Osorio and Franklin, 2008) or gut (Consentino et al., 2021), laser capture microdissection has been put to use in applying omics to defined spatial regions or localised subpopulations of the sample. This could be of interest in deciphering the local behaviour of protective cultures and their targets in distinct and heterogeneous regions of food matrices. The few studies that have approached population functional heterogeneity in real food matrices took advantage of fluorescence microscopy associated with strains reporting the expression of genes of interest by fluorescent proteins (Fleurot et al., 2014; Hernández-Galán et al., 2017). However, food matrices are often opaque to fluorescence microscopy and the density of the population of interest can be too low for reliable quantitative measurements (e.g., bacterial pathogens $<10^2$ CFU/g). Synthetic microbial ecology approaches, where the complexity of the communities and the factors of influence are reduced to their minimum, but are increased in their controllability, can be used to examine interactions and ecological theories (Connell et al., 2012, 2014; Rothschild, 2016; Hynes et al., 2018). Such approaches can be combined with new creative experimental designs for the study of previously unexplored aspects of bacterial behaviour in spatially structured populations (Connell et al., 2013; Wessel et al., 2013; Bridier et al., 2017). In particular, 3D bioprinting of simplified structured matrices with patterned microbial ecosystems could help study population heterogeneities and interspecies interactions at a single-cell scale in structured matrices (Moon et al., 2016; Kyle, 2018; Gyimah et al., 2021; Krishna Kumar et al., 2021). Target microorganisms could be fluorescently tagged for their geolocalisation and the feeding of spatial models of interactions (with food components and with other microorganisms during growth in the printed matrix; Krishna Kumar et al., 2021). Recently, a transcriptome-imaging approach (par-seqFISH for *parallel sequential fluorescence in situ hybridisation*) was reported to capture gene expression and spatial context within microscale assemblies at a single-cell and molecule resolution and could be put to use in such synthetic ecology approaches (Dar et al., 2021). Biopreservation studies could also benefit from microfluidic approaches to assess small-scale interactions between microorganisms (Burmeister and Grünberger, 2020).

The consideration of population heterogeneity in structured food matrices is starting to be integrated into mathematical spatial modelling and predictive microbiology (Verheyen and Van Impe, 2021). Predictive microbiology is useful to quantify both the impact of biopreservation on the food matrix and to simulate the behaviour (survival, growth, inactivation) of the undesirable target microorganism (Leroy and De Vuyst, 2003;

Koutsoumanis et al., 2004; Couvert et al., 2010; Habimana et al., 2011; Møller et al., 2013). Another important point to consider is that the contamination of food products with pathogens such as *L. monocytogenes* and *E. coli* O157:H7 occurs accidentally and usually at low levels, thus requiring single-cell level approaches. Individual-based modelling combined with microenvironment (pH, aw variabilities) modelling of vacuum-packed cold-smoked salmon was more effective in describing variability in the growth of a few *L. monocytogenes* cells than the traditional population models (Ferrier et al., 2013; Augustin et al., 2015). Such stochastic approaches need to be improved by characterising a wider range of microenvironmental factors, such as the variability of the viscosity within food matrices between liquid and solid states, as well as considering biotic factors, namely food components and food microbial communities. However, they could provide complementary information about the behaviour of unwanted microorganisms at realistic contamination levels.

Conclusion

Omics tools have become essential in the field of biopreservation, both for selecting innovative agents and for studying their effectiveness, their mechanism of action, and their impact on the food ecosystem. These approaches have revealed the diversity and complexity of the molecular mechanisms responsible for the protective activity of biopreservation agents. They have provided fundamental knowledge about biopreservation issues in terms of community description (taxonomy), biotic interactions, and impact on the organoleptic quality of the product. They have also shown that the food microbiota plays a major role in biopreservation by acting positively or negatively (Figure 1). It is now clear that the food microbiota must, in the future, be fully integrated into the biopreservation system engineering process to bring the field into the dimension of food microbiome engineering, so that it can play its protective role against pathogens and spoilage microorganisms as well as serve its technological purpose. Moreover, beyond the functioning of the food microbiome, this engineering process must better integrate its interconnections with other microbiomes (soil, water, plant, animal and consumer) to avoid disrupting their functioning and even to

contribute to their balance. In this respect, efforts should be pursued to make more extensive use of multi-omics approaches and to combine them with other complementary approaches that take into account the heterogeneity of microorganisms at the cellular, population, and community levels, as well as the heterogeneity of the food matrix.

Author contributions

FB and MZ coordinated the work and consolidated the manuscript. All authors contributed to the design of the review, carried out the bibliographic data search, drafted the manuscript and approved the submitted version.

Funding

This work was financed by the RMT* Actia Florepro, a scientific and technical partnership in the field of biopreservation established and supported by the French Ministry responsible for Food, under the coordination of Actia. All authors of this review are members of the RMT Actia Florepro. *Réseau mixte technologique: Joint Technological Network.

Conflict of interest

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Publisher's note

All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

References

- Acin-Albiac, M., Filannino, P., Gobetti, M., and Di Cagno, R. (2020). Microbial high throughput phenomics: the potential of an irreplaceable omics. *Comput. Struct. Biotechnol. J.* 18, 2290–9. doi: 10.1016/j.csbj.2020.08.010
- Agriculture and Economic Development Analysis Division (2013). *The State of Food and Agriculture 2013: Food System for Better Nutrition*. Rome: FAO.
- Andreevskaya, M., Johansson, P., Laine, P., Smolander, O.-P., Sonck, M., Rahkila, R., et al. (2015). Genome sequence and transcriptome analysis of meat-spoilage-associated Lactic acid bacterium *Lactococcus piscium* MKFS47. *Appl. Environ. Microbiol.* 81, 3800–11. doi: 10.1128/AEM.00320-15
- Augustin, J.-C., Ferrier, R., Hezard, B., Lintz, A., and Stahl, V. (2015). Comparison of individual-based modeling and population approaches for

- prediction of foodborne pathogens growth. *Food Microbiol.* 45, 205–15. doi: 10.1016/j.fm.2014.04.006
- Aunbjerg, S. D., Honoré, A. H., Marcussen, J., Ebrahimi, P., Vogensen, F. K., Benfeldt, C., et al. (2015). Contribution of volatiles to the antifungal effect of *Lactobacillus paracasei* in defined medium and yogurt. *Int. J. Food Microbiol.* 194, 46–53. doi: 10.1016/j.ijfoodmicro.2014.11.007
- Back, A., Borges, F., Mangavel, C., Paris, C., Rondags, E., Kapel, R., et al. (2015). Recombinant pediocin in *Lactococcus lactis*: increased production by propeptide fusion and improved potency by co-production with PedC. *Microb. Biotechnol.* 9, 466–77. doi: 10.1111/1751-7915.12285
- Baltz, R. H. (2019). Natural product drug discovery in the genomic era: realities, conjectures, misconceptions, and opportunities. *J. Ind. Microbiol. Biotechnol.* 46, 281–99. doi: 10.1007/s10295-018-2115-4
- Barbosa, J., Albano, H., Silva, B., Almeida, M. H., Nogueira, T., and Teixeira, P. (2021). Characterization of a *Lactiplantibacillus plantarum* R23 isolated from Arugula by whole-genome sequencing and its bacteriocin production ability. *IJERPH* 18:5515. doi: 10.3390/ijerph18115515
- Bassi, D., Gazzola, S., Sattin, E., Dal Bello, F., Simionati, B., and Coconcelli, P. S. (2020). Lactic acid bacteria adjunct cultures exert a mitigation effect against spoilage microbiota in fresh cheese. *Microorganisms* 8:E1199. doi: 10.3390/microorganisms8081199
- Begrem, S., Ivaniuk, F., Gigout-Chevalier, F., Kolypczuk, L., Bonnetot, S., Leroi, F., et al. (2020). New insight into antimicrobial compounds from food and marine-sourced *Carnobacterium* species through phenotype and genome analyses. *Microorganisms* 8:1093. doi: 10.3390/microorganisms8071093
- Ben Said, L., Gaudreau, H., Dallaire, L., Tessier, M., and Fliss, I. (2019). Bioprotective culture: a new generation of food additives for the preservation of food quality and safety. *Ind. Biotechnol.* 15, 138–47. doi: 10.1089/ind.2019.29175.1b5
- Berg, G., Rybakova, D., Fischer, D., Cernava, T., Vergès, M.-C. C., Charles, T., et al. (2020). Microbiome definition re-visited: old concepts and new challenges. *Microbiome* 8:103. doi: 10.1186/s40168-020-00875-0
- Besnard, A., Desmasures, N., Voisin-Anastasia, A., Gréau, L., Lelièvre, V., Bré, J.-M., et al. (2021). *Aerococcus* sp. a promising genus as a source of anti-*Salmonella* bioprotective agents for the dairy industry revealed by a miniaturised screening method. *Int. Dairy J.* 116:104949. doi: 10.1016/j.idairyj.2020.104949
- Blin, K., Wolf, T., Chevrette, M. G., Lu, X., Schwalen, C. J., Kautsar, S. A., et al. (2017). antiSMASH 4.0-improvements in chemistry prediction and gene cluster boundary identification. *Nucleic Acids Res.* 45, W36–41. doi: 10.1093/nar/gkx319
- Borges, F., and Revol-Junelles, A.-M. (2019). *Nouvelles Souches de Carnobacterium Maltaromaticum et Leurs Utilisations*. French Patent No FR1911895. Courbevoie: Institut National de la Propriété Industrielle.
- Bourdichon, F., Arias, E., Bückle, A., Bello, F. D., Dubois, A., et al. (2021). The forgotten role of food cultures. *FEMS Microbiol. Lett.* 368:fnab085. doi: 10.1093/femsle/fnab085
- Bridier, A., Piard, J.-C., Pandin, C., Labarthe, S., Dubois-Brissonnet, F., and Briandet, R. (2017). Spatial Organization Plasticity as an adaptive driver of surface microbial communities. *Front. Microbiol.* 8:1364. doi: 10.3389/fmicb.2017.01364
- Brillet, A., Pilet, M.-F., Prévost, H., Bouttefroy, A., and Leroi, F. (2004). Biodiversity of *Listeria monocytogenes* sensitivity to bacteriocin-producing *Carnobacterium* strains and application in sterile cold-smoked salmon. *J. Appl. Microbiol.* 97, 1029–37. doi: 10.1111/j.1365-2672.2004.02383.x
- Burmeister, A., and Grünberger, A. (2020). Microfluidic cultivation and analysis tools for interaction studies of microbial co-cultures. *Curr. Opin. Biotechnol.* 62, 106–15. doi: 10.1016/j.copbio.2019.09.001
- Bury-Moné, S., and Scavi, B. (2017). Stochasticity of gene expression as a motor of epigenetics in bacteria: from individual to collective behaviors. *Res. Microbiol.* 168, 503–14. doi: 10.1016/j.resmic.2017.03.009
- Caldeira, C., De Laurentiis, V., Corrado, S., van Holsteijn, F., and Sala, S. (2019). Quantification of food waste per product group along the food supply chain in the European Union: a mass flow analysis. *Resour. Conserv. Recycl.* 149, 479–88. doi: 10.1016/j.resconrec.2019.06.011
- Callon, C., Arliguie, C., and Montel, M.-C. (2016). Control of Shigatoxin-producing *Escherichia coli* in cheese by dairy bacterial strains. *Food Microbiol.* 53, 63–70. doi: 10.1016/j.fm.2015.08.009
- Cavicchioli, R., Ripple, W. J., Timmis, K. N., Azam, F., Bakken, L. R., Baylis, M., et al. (2019). Scientists' warning to humanity: microorganisms and climate change. *Nat. Rev. Microbiol.* 17, 569–86. doi: 10.1038/s41579-019-0222-5
- Chaillou, S., Champomier-Vergès, M.-C., Cornet, M., Crutz-Le Coq, A.-M., Dudez, A.-M., Martin, V., et al. (2005). The complete genome sequence of the meat-borne lactic acid bacterium *Lactobacillus sakei* 23K. *Nat. Biotechnol.* 23, 1527–33. doi: 10.1038/nbt1160
- Chaillou, S., Ramarosan, M., Coeuret, G., Rossero, A., Anthoine, V., Champomier-Vergès, M., et al. (2022). Combination of high pressure treatment at 500 MPa and biopreservation with a *Lactococcus lactis* strain for lowering the bacterial growth during storage of diced cooked ham with reduced nitrite salt. *Microorganisms* 10:456. doi: 10.3390/microorganisms10020456
- Chokesajjawatee, N., Santiyanont, P., Chantarasakha, K., Kocharin, K., Thammarongtham, C., Lertpaiporn, S., et al. (2020). Safety assessment of a Nham starter culture *Lactobacillus plantarum* BCC9546 via whole-genome analysis. *Sci. Rep.* 10:10241. doi: 10.1038/s41598-020-66857-2
- Comi, G., Andyanto, D., Manzano, M., and Iacumin, L. (2016). *Lactococcus lactis* and *Lactobacillus sakei* as bio-protective culture to eliminate *Leuconostoc mesenteroides* spoilage and improve the shelf life and sensorial characteristics of commercial cooked bacon. *Food Microbiol.* 58, 16–22. doi: 10.1016/j.fm.2016.03.001
- Connell, J. L., Kim, J., Shear, J. B., Bard, A. J., and Whiteley, M. (2014). Real-time monitoring of quorum sensing in 3D-printed bacterial aggregates using scanning electrochemical microscopy. *Proc. Natl. Acad. Sci. U S A.* 111, 18255–60. doi: 10.1073/pnas.1421211111
- Connell, J. L., Ritschdorff, E. T., Whiteley, M., and Shear, J. B. (2013). 3D printing of microscopic bacterial communities. *Proc. Natl. Acad. Sci. U S A.* 110, 18380–5. doi: 10.1073/pnas.1309729110
- Connell, J. L., Whiteley, M., and Shear, J. B. (2012). Sociomicrobiology in engineered landscapes. *Nat. Chem. Biol.* 8, 10–3. doi: 10.1038/nchembio.749
- Consentino, L., Rejasse, A., Crapart, N., Bevilacqua, C., and Nielsen-LeRoux, C. (2021). Laser capture microdissection to study *Bacillus cereus* iron homeostasis gene expression during *Galleria mellonella* in vivo gut colonization. *Virulence* 12, 2104–21. doi: 10.1080/21505594.2021.1959790
- Cook, P. W., and Nightingale, K. K. (2018). Use of omics methods for the advancement of food quality and food safety. *Anim. Front.* 8, 33–41. doi: 10.1093/af/vfy024
- Couvert, O., Pinon, A., Bergis, H., Bourdichon, F., Carlin, F., Cornu, M., et al. (2010). Validation of a stochastic modelling approach for *Listeria monocytogenes* growth in refrigerated foods. *Int. J. Food Microbiol.* 144, 236–42. doi: 10.1016/j.ijfoodmicro.2010.09.024
- Cretenet, M., Nouaille, S., Thouin, J., Rault, L., Stenz, L., François, P., et al. (2011). *Staphylococcus aureus* virulence and metabolism are dramatically affected by *Lactococcus lactis* in cheese matrix: *S. aureus* interaction with *L. lactis* in cheese matrix. *Environ. Microbiol. Rep.* 3, 340–51. doi: 10.1111/j.1758-2229.2010.00230.x
- Dar, D., Dar, N., Cai, L., and Newman, D. K. (2021). Spatial transcriptomics of planktonic and sessile bacterial populations at single-cell resolution. *Science* 373:eabi4882. doi: 10.1126/science.abi4882
- Darsonval, M., Grégoire, M., Deschamps, J., and Briandet, R. (2021). “Confocal laser microscopy analysis of *Listeria monocytogenes* biofilms and spatially organized communities,” in *Listeria monocytogenes*. *Methods Mol Biol*, eds E. M. Fox, H. Bierne, and B. Stessl (New York, NY: Springer), 123–36. doi: 10.1007/978-1-0716-0982-8_10
- Delavenne, E., Déniel, F., Barbier, G., and Le Blay, G. (2012). Biodiversity of antifungal lactic acid bacteria isolated from raw milk samples from cow, ewe and goat over one-year period. *Int. J. Food Microbiol.* 155, 185–90. doi: 10.1016/j.ijfoodmicro.2012.02.003
- Delbes-Paus, C., Dorchie, G., Chaabna, Z., Callon, C., and Montel, M.-C. (2010). Contribution of hydrogen peroxide to the inhibition of *Staphylococcus aureus* by *Lactococcus garvieae* in interaction with raw milk microbial community. *Food Microbiol.* 27, 924–32. doi: 10.1016/j.fm.2010.05.031
- Delpech, P., Bornes, S., Alaterre, E., Bonnet, M., Gagne, G., Montel, M.-C., et al. (2015). *Staphylococcus aureus* transcriptomic response to inhibition by H2O2-producing *Lactococcus garvieae*. *Food Microbiol.* 51, 163–70. doi: 10.1016/j.fm.2015.05.014
- Delpech, P., Rifa, E., Ball, G., Nidelet, S., Dubois, E., Gagne, G., et al. (2017). New insights into the anti-pathogenic potential of *Lactococcus garvieae* against *Staphylococcus aureus* based on RNA sequencing Profiling. *Front. Microbiol.* 8:359. doi: 10.3389/fmicb.2017.00359
- Duhutrel, P., Bordat, C., Wu, T.-D., Zagorec, M., Guerquin-Kern, J.-L., and Champomier-Vergès, M.-C. (2010). Iron sources used by the nonpathogenic lactic acid bacterium *Lactobacillus sakei* as revealed by electron energy loss spectroscopy and secondary-ion mass spectrometry. *Appl. Environ. Microbiol.* 76, 560–5. doi: 10.1128/AEM.02205-09
- Ebrahimi, P., Larsen, F. H., Jensen, H. M., Vogensen, F. K., and Engelsen, S. B. (2016). Real-time metabolomic analysis of lactic acid bacteria as monitored by in vitro NMR and chemometrics. *Metabolomics* 12, 1–17. doi: 10.1007/s11306-016-0996-7
- Edgar, R. C. (2017). UNBIAS: an attempt to correct abundance bias in 16S sequencing, with limited success. *bioRxiv* 2017.124149. doi: 10.1101/124149

- Eijssink, V. G. H., and Axelsson, L. (2005). Bacterial lessons in sausage making. *Nat. Biotechnol.* 23, 1494–5. doi: 10.1038/nbt1205-1494
- El Kheir, S. M., Cherrat, L., Awussi, A. A., Ramia, N. E., Taha, S., Rahman, A., et al. (2018). High-throughput identification of candidate strains for biopreservation by using bioluminescent *Listeria monocytogenes*. *Front. Microbiol.* 9:1883. doi: 10.3389/fmicb.2018.01883
- Ellegaard, K. M., and Engel, P. (2016). Beyond 16S rRNA community profiling: intra-species diversity in the gut microbiota. *Front. Microbiol.* 7:1475. doi: 10.3389/fmicb.2016.01475
- Elsner-Gravesen, D., and Elsner-Gravesen, A. (2013). "Biopreservatives," in *Biotechnology of Food and Feed Additives Advances in Biochemical Engineering/Biotechnology*, eds H. Zorn and P. Cermak (Berlin: Springer), 29–49. doi: 10.1007/10_2013_234
- Ercolini, D., Ferrocino, I., La Storia, A., Mauriello, G., Gigli, S., Masi, P., et al. (2010). Development of spoilage microbiota in beef stored in nisin activated packaging. *Food Microbiol.* 27, 137–43. doi: 10.1016/j.fm.2009.09.006
- Even, S., Charlier, C., Nouaille, S., Ben Zakour, N. L., Cretenet, M., Cousin, F. J., et al. (2009). *Staphylococcus aureus* virulence expression is impaired by *Lactococcus lactis* in mixed cultures. *Appl. Environ. Microbiol.* 75, 4459–72. doi: 10.1128/AEM.02388-08
- Fall, P. A., Pilet, M. F., Leduc, F., Cardinal, M., Duflos, G., Guérin, C., et al. (2012). Sensory and physicochemical evolution of tropical cooked peeled shrimp inoculated by *Brochothrix thermosphacta* and *Lactococcus piscium* CNCM I-4031 during storage at 8°C. *Int. J. Food Microbiol.* 152, 82–90. doi: 10.1016/j.ijfoodmicro.2011.07.015
- FAO (2014). *Food Waste Footprint: Full Cost-Accounting: Final Report*. Rome: FAO.
- FAO (2019). *State of Food and Agriculture 2019. Moving Forward on Food Loss and Waste Reduction*. Rome: FAO.
- Ferrier, R., Hezard, B., Lintz, A., Stahl, V., and Augustin, J.-C. (2013). Combining individual-based modeling and food microenvironment descriptions to predict the growth of *Listeria monocytogenes* on smear soft cheese. *Appl. Environ. Microbiol.* 2013:13. doi: 10.1128/AEM.01311-13
- Ferrocino, I., Greppi, A., Storia, A. L., Rantsiou, K., Ercolini, D., and Coccolin, L. (2015). Impact of nisin-activated packaging on microbiota of beef burgers during storage. *Appl. Environ. Microbiol.* 2015:15. doi: 10.1128/AEM.03093-15
- Finnegan, M., Linley, E., Denyer, S. P., McDonnell, G., Simons, C., and Maillard, J. Y. (2010). Mode of action of hydrogen peroxide and other oxidizing agents: differences between liquid and gas forms. *J. Antimicrob. Chemother.* 65, 2108–15. doi: 10.1093/jac/dkq308
- Fleuret, I., Aigle, M., Fleuret, R., Darrigo, C., Hennekinne, J.-A., Gruss, A., et al. (2014). Following pathogen development and gene expression in a food ecosystem: the case of a *Staphylococcus aureus* isolate in cheese. *Appl. Environ. Microbiol.* 80, 5106–15. doi: 10.1128/AEM.01042-14
- Frétin, M., Chassard, C., Delbès, C., Lavigne, R., Rifa, E., Theil, S., et al. (2020). Robustness and efficacy of an inhibitory consortium against *E. coli* O26:H11 in raw milk cheeses. *Food Control* 115:107282. doi: 10.1016/j.foodcont.2020.107282
- Galia, W., Leriche, F., Cruveiller, S., Garnier, C., Navratil, V., Dubost, A., et al. (2017). Strand-specific transcriptomes of Enterohemorrhagic *Escherichia coli* in response to interactions with ground beef microbiota: interactions between microorganisms in raw meat. *BMC Genom.* 18:574. doi: 10.1186/s12864-017-3957-2
- Gálvez, A., López, R. L., and Ben Omar, N. (2007). Bacteriocin-based strategies for food biopreservation. *Int. J. Food Microbiol.* 120, 51–70. doi: 10.1016/j.ijfoodmicro.2007.06.001
- Garnier, L., Penland, M., Thierry, A., Maillard, M.-B., Jardin, J., Coton, M., et al. (2020). Antifungal activity of fermented dairy ingredients: identification of antifungal compounds. *Int. J. Food Microbiol.* 322:108574. doi: 10.1016/j.ijfoodmicro.2020.108574
- Giello, M., La Storia, A., De Filippis, F., Ercolini, D., and Villani, F. (2018). Impact of *Lactobacillus curvatus* 54M16 on microbiota composition and growth of *Listeria monocytogenes* in fermented sausages. *Food Microbiol.* 72, 1–15. doi: 10.1016/j.fm.2017.11.003
- Guillier, L., Stahl, V., Hezard, B., Notz, E., and Briandet, R. (2008). Modelling the competitive growth between *Listeria monocytogenes* and biofilm microflora of smear cheese wooden shelves. *Int. J. Food Microbiol.* 128, 51–7. doi: 10.1016/j.ijfoodmicro.2008.06.028
- Gyimah, N., Scheler, O., Rang, T., and Pardy, T. (2021). Can 3D printing bring droplet microfluidics to every lab?—A systematic review. *Micromachines* 12:339. doi: 10.3390/mi12030339
- Habimana, O., Guillier, L., Kulakauskas, S., and Briandet, R. (2011). Spatial competition with *Lactococcus lactis* in mixed-species continuous-flow biofilms inhibits *Listeria monocytogenes* growth. *Biofouling* 27, 1065–72. doi: 10.1080/08927014.2011.626124
- Hebert, E. M., Saavedra, L., Taranto, M. P., Mozzi, F., Magni, C., Nader, M. E. F., et al. (2012). Genome sequence of the bacteriocin-producing *Lactobacillus curvatus* strain CRL705. *J. Bacteriol.* 194, 538–9. doi: 10.1128/JB.06416-11
- Hernández-Galán, L., Cattenez, T., Le Feunteun, S., Canette, A., Briandet, R., Le-Guin, S., et al. (2017). Effect of dairy matrices on the survival of *Streptococcus thermophilus*, *Brevibacterium aurantiacum* and *Hafnia alvei* during digestion. *Food Res. Int.* 100, 477–88. doi: 10.1016/j.foodres.2017.07.044
- Hibbing, M. E., Fuqua, C., Parsek, M. R., and Peterson, S. B. (2010). Bacterial competition: surviving and thriving in the microbial jungle. *Nat. Rev. Microbiol.* 8, 15–25. doi: 10.1038/nrmicro2259
- Gänzle, M. G., Hötzel, A., Walter, J., Jung, G., and Hammes, W. P. (2000). Characterization of reutericyclin produced by *Lactobacillus reuteri* LTH2584. *Appl. Environ. Microbiol.* 66, 4325–33. doi: 10.1128/AEM.66.10.4325-4333.2000
- HPLÉ (2014). "Food losses and waste in the context of sustainable food systems," in *A report by the high level panel of experts on food security and nutrition. HLPE Report 8*, (Rome: FAO).
- Hu, P., Xu, X. L., Zhou, G. H., Han, Y. Q., Xu, B. C., and Liu, J. C. (2008). Study of the *Lactobacillus sakei* protective effect towards spoilage bacteria in vacuum packed cooked ham analyzed by PCR-DGGE. *Meat Sci.* 80, 462–9. doi: 10.1016/j.meatsci.2008.01.011
- Hynes, W. F., Chacón, J., Segrè, D., Marx, C. J., Cady, N. C., and Harcombe, W. R. (2018). Bioprinting microbial communities to examine interspecies interactions in time and space. *Biomed. Phys. Eng. Express* 4:55010. doi: 10.1088/2057-1976/aa544
- Iacumin, L., Cappellari, G., Colautti, A., and Comi, G. (2020). *Listeria monocytogenes* survey in cubed cooked ham packaged in modified atmosphere and bioprotective effect of selected lactic acid bacteria. *Microorganisms* 8:898. doi: 10.3390/microorganisms8060898
- Jameson, J. E. (1962). A discussion of the dynamics of salmonella enrichment. *J. Hyg.* 60, 193–207. doi: 10.1017/s0022172400039462
- Jeanson, S., Floury, J., Gagnaire, V., Lortal, S., and Thierry, A. (2015). Bacterial colonies in solid media and foods: a review on their growth and interactions with the micro-environment. *Front. Microbiol.* 6:1284. doi: 10.3389/fmicb.2015.01284
- Joffraud, J.-J., Cardinal, M., Cornet, J., Léon, S., Gigout, F., et al. (2006). Effect of bacterial interactions on the spoilage of cold-smoked salmon. *Int. J. Food Microbiol.* 112, 51–61. doi: 10.1016/j.ijfoodmicro.2006.05.014
- Jones, R. J., Wiklund, E., Zagorec, M., and Tagg, J. R. (2010). Evaluation of stored lamb bio-preserved using a three-strain cocktail of *Lactobacillus sakei*. *Meat Sci.* 86, 955–9. doi: 10.1016/j.meatsci.2010.07.023
- Koutsoumanis, K. P., Kendall, P. A., and Sofos, J. N. (2004). A comparative study on growth limits of *Listeria monocytogenes* as affected by temperature, pH and aw when grown in suspension or on a solid surface. *Food Microbiol.* 21, 415–22. doi: 10.1016/j.fm.2003.11.003
- Krishna Kumar, R., Meiller-Légrand, T. A., Alcinesio, A., Gonzalez, D., Mavridou, D. A. I., Meacock, O. J., et al. (2021). Droplet printing reveals the importance of micron-scale structure for bacterial ecology. *Nat. Commun.* 12:857. doi: 10.1038/s41467-021-20996-w
- Kyle, S. (2018). 3D printing of bacteria: the next frontier in bofabrication. *Trends Biotechnol.* 36, 340–1. doi: 10.1016/j.tibtech.2018.01.010
- Laursen, M. F., Bahl, M. I., Licht, T. R., Gram, L., and Knudsen, G. M. (2015). A single exposure to a sublethal pediocin concentration initiates a resistance-associated temporal cell envelope and general stress response in *Listeria monocytogenes*. *Environ. Microbiol.* 17, 1134–51. doi: 10.1111/1462-2920.12534
- Ledenbach, L. H., and Marshall, R. T. (2009). "Microbiological spoilage of dairy products," in *Compendium of the Microbiological Spoilage of Foods and Beverages*, eds W. H. Sperber and M. P. Doyle (New York, NY: Springer), 41–67. doi: 10.1007/978-1-4419-0826-1_2
- Leisner, J. J., Laursen, B. G., Prevost, H., Drider, D., and Dalgaard, P. (2007). *Carnobacterium*: positive and negative effects in the environment and in foods. *FEMS Microbiol. Rev.* 31, 592–613. doi: 10.1111/j.1574-6976.2007.00080.x
- Lenz, A. P., Williamson, K. S., Pitts, B., Stewart, P. S., and Franklin, M. J. (2008). Localized gene expression in *Pseudomonas aeruginosa* biofilms. *Appl. Environ. Microbiol.* 74, 4463–71. doi: 10.1128/AEM.00710-08
- Leroi, F., Cornet, J., Chevalier, F., Cardinal, M., Coeuret, G., Chaillou, S., et al. (2015). Selection of bioprotective cultures for preventing cold-smoked salmon spoilage. *Int. J. Food Microbiol.* 213, 79–87. doi: 10.1016/j.ijfoodmicro.2015.05.005

- Leroy, F., and De Vuyst, L. (2003). A combined model to predict the functionality of the bacteriocin-producing *Lactobacillus sakei* strain CTC 494. *Appl. Environ. Microbiol.* 69, 1093–9. doi: 10.1128/AEM.69.2.1093-1099.2003
- Leyva Salas, M., Mounier, J., Maillard, M.-B., Valence, F., Coton, E., and Thierry, A. (2019). Identification and quantification of natural compounds produced by antifungal bioprotective cultures in dairy products. *Food Chem.* 301:125260. doi: 10.1016/j.foodchem.2019.125260
- Leyva Salas, M., Mounier, J., Valence, F., Coton, M., Thierry, A., and Coton, E. (2017). Antifungal microbial agents for food biopreservation-A Review. *Microorganisms* 5:5030037. doi: 10.3390/microorganisms5030037
- Leyva Salas, M., Thierry, A., Lemaitre, M., Garric, G., Harel-Oger, M., Chatel, M., et al. (2018). Antifungal activity of lactic acid bacteria combinations in dairy mimicking models and their potential as bioprotective cultures in pilot scale applications. *Front. Microbiol.* 9:1787. doi: 10.3389/fmicb.2018.01787
- Li, J., Yang, X., Shi, G., Chang, J., Liu, Z., and Zeng, M. (2019). Cooperation of lactic acid bacteria regulated by the AI-2/LuxS system involve in the biopreservation of refrigerated shrimp. *Food Res. Int.* 120, 679–87. doi: 10.1016/j.foodres.2018.11.025
- Lin, X. B., Lohans, C. T., Duar, R., Zheng, J., Vederas, J. C., Walter, J., et al. (2015). Genetic determinants of reutericyclin biosynthesis in *Lactobacillus reuteri*. *Appl. Environ. Microbiol.* 2015:14. doi: 10.1128/AEM.03691-14
- Liu, Y.-X., Qin, Y., Chen, T., Lu, M., Qian, X., Guo, X., et al. (2021). A practical guide to amplicon and metagenomic analysis of microbiome data. *Protein Cell* 12, 315–30. doi: 10.1007/s13238-020-00724-8
- Luo, C., Knight, R., Siljander, H., Knip, M., Xavier, R. J., and Gevers, D. (2015). ConStrains identifies microbial strains in metagenomic datasets. *Nat. Biotechnol.* 33, 1045–52. doi: 10.1038/nbt.3319
- Marché, L., Saraoui, T., Remenant, B., Zagorec, M., Prévost, H., Delbarre-Ladrat, C., et al. (2017). Complete genome sequence of *Lactococcus piscium* CNCM I-4031, a bioprotective strain for seafood products. *Genome Announc.* 5:16. doi: 10.1128/genomeA.01510-16
- Martinez, R. C. R., Alvarenga, V. O., ávaro-Trindade, C. S., Sant'Ana, A., and de S. (2016). Assessment of the inhibitory effect of free and encapsulated commercial nisin (Nisaplin®), tested alone and in combination, on *Listeria monocytogenes* and *Bacillus cereus* in refrigerated milk. *LWT - Food Sci. Technol.* 68, 67–75. doi: 10.1016/j.lwt.2015.12.027
- McNair, L. K. F., Siedler, S., Vinther, J. M. O., Hansen, A. M., Neves, A. R., Garrigues, C., et al. (2018). Identification and characterization of a new antifungal peptide in fermented milk product containing bioprotective *Lactobacillus* cultures. *FEMS Yeast Res.* 18:foyo94. doi: 10.1093/femsyr/foyo94
- Meola, M., Rifa, E., Shani, N., Delbès, C., Berthoud, H., and Chassard, C. (2019). DAIRYdb: a manually curated reference database for improved taxonomy annotation of 16S rRNA gene sequences from dairy products. *BMC Genom.* 20:560. doi: 10.1186/s12864-019-5914-8
- Metivier, A., Pilet, M.-F., Dousset, X., Sorokine, O., Anglade, P., Zagorec, M., et al. (1998). Divercin V41, a new bacteriocin with two disulphide bonds produced by *Carnobacterium divergens* V41: primary structure and genomic organization. *Microbiology* 144, 2837–44. doi: 10.1099/00221287-144-10-2837
- Møller, C. O. A., Ilg, Y., Aabo, S., Christensen, B. B., Dalgaard, P., and Hansen, T. B. (2013). Effect of natural microbiota on growth of *Salmonella* spp. in fresh pork – A predictive microbiology approach. *Food Microbiol.* 34, 284–95. doi: 10.1016/j.fm.2012.10.010
- Moon, S., Fritz, I. L., Singer, Z. S., and Danino, T. (2016). Spatial control of bacteria using screen printing. *3D Print Addit. Manuf* 3, 194–203. doi: 10.1089/3dp.2016.0040
- Nogueira Viçosa, G., Vieira Botelho, C., Botta, C., Bertolino, M., Fernandes de Carvalho, A., Nero, L. A., et al. (2019). Impact of co-cultivation with *Enterococcus faecalis* over growth, enterotoxin production and gene expression of *Staphylococcus aureus* in broth and fresh cheeses. *Int. J. Food Microbiol.* 308:108291. doi: 10.1016/j.ijfoodmicro.2019.108291
- Nouaille, S., Even, S., Charlier, C., Le Loir, Y., Coccain-Bousquet, M., and Loubière, P. (2009). Transcriptomic response of *Lactococcus lactis* in mixed culture with *Staphylococcus aureus*. *Appl. Environ. Microbiol.* 75, 4473–82. doi: 10.1128/AEM.02653-08
- Nouaille, S., Rault, L., Jeanson, S., Loubière, P., Le Loir, Y., and Even, S. (2014). Contribution of *Lactococcus lactis* reducing properties to the downregulation of a major virulence regulator in *Staphylococcus aureus*, the *agr* system. *Appl. Environ. Microbiol.* 80, 7028–35. doi: 10.1128/AEM.02287-14
- Orihuel, A., Terán, L., Renaut, J., Planchon, S., Valacco, M. P., Masias, E., et al. (2019). Physiological and proteomic response of *Escherichia coli* O157:H7 to a bioprotective lactic acid bacterium in a meat environment. *Food Res. Int.* 125:108622. doi: 10.1016/j.foodres.2019.108622
- Orihuel, A., Terán, L., Renaut, J., Vignolo, G. M., De Almeida, A. M., Saavedra, M. L., et al. (2018). Differential proteomic analysis of lactic acid bacteria-*Escherichia coli* O157:H7 interaction and its contribution to bioprotection strategies in Meat. *Front. Microbiol.* 9:1083. doi: 10.3389/fmicb.2018.01083
- Ortiz-Rivera, Y., Sánchez-Vega, R., Gutiérrez-Méndez, N., León-Félix, J., Acosta-Muñiz, C., and Sepulveda, D. R. (2017). Production of reuterin in a fermented milk product by *Lactobacillus reuteri*: inhibition of pathogens, spoilage microorganisms, and lactic acid bacteria. *J. Dairy Sci.* 100, 4258–68. doi: 10.3168/jds.2016-11534
- Park, E. J., Kim, K. H., Abell, G. C. J., Kim, M. S., Roh, S. W., and Bae, J. W. (2011). Metagenomic analysis of the viral communities in fermented foods. *Appl. Environ. Microbiol.* 77, 1284–91. doi: 10.1128/AEM.01859-10
- Pérez-Osorio, A. C., and Franklin, M. J. (2008). Isolation of RNA and DNA from biofilm samples obtained by laser capture microdissection microscopy. *CSH Protoc.* 2008:rot5065. doi: 10.1101/pdb.prot5065
- Pinilla, C. M. B., Stincone, P., and Brandelli, A. (2021). Proteomic analysis reveals differential responses of *Listeria monocytogenes* to free and nanoencapsulated nisin. *Int. J. Food Microbiol.* 346:109170. doi: 10.1016/j.ijfoodmicro.2021.109170
- Poirier, S., Coeuret, G., Champomier-Vergès, M.-C., and Chaillou, S. (2018). Draft genome sequences of nine strains of *Brochothrix thermosphacta*, *Carnobacterium divergens*, *Lactobacillus algidus*, *Lactobacillus fuchuensis*, *Lactococcus piscium*, *Leuconostoc gelidium* subsp. *gasicomitatum*, *Pseudomonas lundensis*, and *Weissella viridescens*, a collection of psychrotrophic species involved in meat and seafood spoilage. *Genome Announc.* 6, e479–418. doi: 10.1128/genomeA.00479-18
- Ramaroson, M., Guillou, S., Rossero, A., Rezé, S., Anthoine, V., Moriceau, N., et al. (2018). Selection procedure of bioprotective cultures for their combined use with High Pressure Processing to control spore-forming bacteria in cooked ham. *Int. J. Food Microbiol.* 276, 28–38. doi: 10.1016/j.ijfoodmicro.2018.04.010
- Rathod, N. B., Nirmal, N. P., Pagarkar, A., Özogul, F., and Rocha, J. M. (2022). Antimicrobial impacts of microbial metabolites on the preservation of fish and fishery products: a review with current knowledge. *Microorganisms* 10:773. doi: 10.3390/microorganisms10040773
- Remenant, B., Borges, F., Cailliez-Grimal, C., Revol-Junelles, A.-M., Marché, L., Lajus, A., et al. (2016). Draft genome sequence of *Carnobacterium divergens* V41, a bacteriocin-producing strain. *Genome Announc.* 4, e1109–16. doi: 10.1128/genomeA.01109-16
- Riedel, C. U., Monk, I. R., Casey, P. G., Morrissey, D. O., Sullivan, G. C., Tangney, M., et al. (2007). Improved luciferase tagging system for *Listeria monocytogenes* allows real-time monitoring in vivo and in vitro. *Appl. Environ. Microbiol.* 73, 3091–4. doi: 10.1128/AEM.02940-06
- Rimaux, T., Vrancken, G., Vuylsteke, B., De Vuyst, L., and Leroy, F. (2011). The pentose moiety of adenosine and inosine is an important energy source for the fermented-meat starter culture *Lactobacillus sakei* CTC 494. *Appl. Environ. Microbiol.* 77, 6539–50. doi: 10.1128/AEM.00498-11
- Roh, S. W., Kim, K. H., Nam, Y. D., et al. (2010). Investigation of archaeal and bacterial diversity in fermented seafood using barcoded pyrosequencing. *ISME J.* 4, 1–16. doi: 10.1038/ismej.2009.83
- Rothschild, L. J. (2016). Synthetic biology meets bioprinting: enabling technologies for humans on Mars (and Earth). *Biochem. Soc. Trans.* 44, 1158–64. doi: 10.1042/BST20160067
- Saint Martin, C., Darsonval, M., Grégoire, M., Caccia, N., Midoux, L., Berland, S., et al. (2022). Spatial organisation of *Listeria monocytogenes* and *Escherichia coli* O157:H7 cultivated in gel matrices. *Food Microbiol.* 103:103965. doi: 10.1016/j.fm.2021.103965
- Saraoui, T., Cornet, J., Guillouet, E., Pilet, M. F., Chevalier, F., Joffraud, J.-J., et al. (2017). Improving simultaneously the quality and safety of cooked and peeled shrimp using a cocktail of bioprotective lactic acid bacteria. *Int. J. Food Microbiol.* 241, 69–77. doi: 10.1016/j.ijfoodmicro.2016.09.024
- Saraoui, T., Fall, P. A., Leroi, F., Antignac, J.-P., Chéreau, S., and Pilet, M. F. (2016). Inhibition mechanism of *Listeria monocytogenes* by a bioprotective bacteria *Lactococcus piscium* CNCM I-4031. *Food Microbiol.* 53, 70–8. doi: 10.1016/j.fm.2015.01.002
- Saraoui, T., Leroi, F., Chevalier, F., Cappelletti, J.-M., Passerini, D., and Pilet, M.-F. (2018). Bioprotective Effect of *Lactococcus piscium* CNCM I-4031 against *Listeria monocytogenes* growth and virulence. *Front. Microbiol.* 9:1564. doi: 10.3389/fmicb.2018.01564
- Schlusselhuber, M., Godard, J., Sebban, M., Bernay, B., Garon, D., Seguin, V., et al. (2018). Characterization of milkisin, a novel lipopeptide with antimicrobial properties produced by *Pseudomonas* sp. UCMA 17988 isolated from bovine raw milk. *Front. Microbiol.* 9:1030. doi: 10.3389/fmicb.2018.01030

- Schlüsselhuber, M., Godard, J., Sebban, M., Bernay, B., Garon, D., Seguin, V., et al. (2020). Corrigendum: characterization of milkisin, a novel lipopeptide with antimicrobial properties produced by *Pseudomonas* sp. UCMA 17988 isolated from bovine raw milk. *Front. Microbiol.* 11:1323. doi: 10.3389/fmicb.2020.01323
- Shi, C., and Knöchel, S. (2021a). Inhibitory effects of binary combinations of microbial metabolites on the growth of tolerant *Penicillium roqueforti* and *Mucor circinelloides*. *LWT* 149:112039. doi: 10.1016/j.lwt.2021.112039
- Shi, C., and Knöchel, S. (2021b). Susceptibility of dairy associated molds towards microbial metabolites with focus on the response to diacetyl. *Food Control* 121:107573. doi: 10.1016/j.foodcont.2020.107573
- Shi, C., and Maktabdar, M. (2022). Lactic acid bacteria as biopreservation against spoilage molds in dairy products – A review. *Front. Microbiol.* 12:819684. doi: 10.3389/fmicb.2021.819684
- Siedler, S., Balti, R., and Neves, A. R. (2019). Bioprotective mechanisms of lactic acid bacteria against fungal spoilage of food. *Curr. Opin. Biotechnol.* 56, 138–46. doi: 10.1016/j.copbio.2018.11.015
- Silbade, A., Cornet, J., Cardinal, M., Chevalier, F., Rochefort, K., Smith-Ravin, J., et al. (2018). Characterization of the spoilage potential of pure and mixed cultures of bacterial species isolated from tropical yellowfin tuna (*Thunnus albacares*). *J. Appl. Microbiol.* 124, 559–71. doi: 10.1111/jam.13663
- Simonin, S., Roullier-Gall, C., Ballester, J., Schmitt-Kopplin, P., Quintanilla-Casas, B., Vichi, S., et al. (2020). Bio-protection as an alternative to sulphites: impact on chemical and microbial characteristics of red wines. *Front. Microbiol.* 11:1308. doi: 10.3389/fmicb.2020.01308
- Spanu, C., Piras, F., Mocci, A. M., Nieddu, G., De Santis, E. P. L., and Scarano, C. (2018). Use of *Carnobacterium* spp protective culture in MAP packed Ricotta fresca cheese to control *Pseudomonas* spp. *Food Microbiol.* 74, 50–6. doi: 10.1016/j.fm.2018.02.020
- Stiles, M. E. (1996). Biopreservation by lactic acid bacteria. *Antonie van Leeuwenhoek* 70, 331–45. doi: 10.1007/BF00395940
- Stincone, P., Comerlato, C. B., and Brandelli, A. (2021). Proteomic analysis of *Listeria monocytogenes* exposed to free and nanostructured antimicrobial lipopeptides. *Mol. Omics* 17, 426–37. doi: 10.1039/D0MO00178C
- Stohr, V., Joffraud, J. J., Cardinal, M., and Leroi, F. (2001). Spoilage potential and sensory profile associated with bacteria isolated from cold-smoked salmon. *Food Res. Int.* 34, 797–806. doi: 10.1016/S0963-9969(01)00101-6
- Ström, K., Sjögren, J., Broberg, A., and Schnürer, J. (2002). *Lactobacillus plantarum* MiLAB 393 produces the antifungal cyclic dipeptides cyclo(L -Phe-L -Pro) and cyclo(L -Phe- trans -4-OH- L -Pro) and 3-phenyllactic acid. *Appl. Environ. Microbiol.* 68, 4322–7. doi: 10.1128/AEM.68.9.4322-4327.2002
- Tabanelli, G., Barbieri, F., Campedelli, I., Venturini, M. C., Gardini, F., and Montanari, C. (2020). Effects of bioprotective cultures on the microbial community during storage of Italian fresh filled pasta. *Food Control* 115:107304. doi: 10.1016/j.foodcont.2020.107304
- Truchado, P., Elsser-Gravesen, A., Gil, M. I., and Allende, A. (2020). Post-process treatments are effective strategies to reduce *Listeria monocytogenes* on the surface of leafy greens: a pilot study. *Int. J. Food Microbiol.* 313:108390. doi: 10.1016/j.ijfoodmicro.2019.108390
- van Heel, A. J., de Jong, A., Song, C., Viel, J. H., Kok, J., and Kuipers, O. P. (2018). BAGEL4: a user-friendly web server to thoroughly mine RiPPs and bacteriocins. *Nucleic Acids Res.* 46, W278–81. doi: 10.1093/nar/gky383
- Verheyen, D., and Van Impe, J. F. M. (2021). The inclusion of the food microstructural influence in predictive microbiology: state-of-the-art. *Foods* 10:2119. doi: 10.3390/foods10092119
- Verheyen, D., Érez-Rodríguez, F., Baka, M., Skåra, T., and Van Impe, J. F. (2018). Effect of food microstructure on growth dynamics of *Listeria monocytogenes* in fish-based model systems. *Int. J. Food Microbiol.* 283, 7–13. doi: 10.1016/j.ijfoodmicro.2018.05.032
- Verplaetse, E., André-Leroux, G., Duhutrel, P., Coeuret, G., Chaillou, S., Nielsen-Leroux, C., et al. (2020). Heme uptake in *Lactobacillus sakei* evidenced by a new energy coupling factor (ECF)-like transport system. *Appl. Environ. Microbiol.* 2020:19. doi: 10.1128/AEM.02847-19
- Viçosa, G. N., Botta, C., Ferrocino, I., Bertolino, M., Ventura, M., Nero, L. A., et al. (2018). *Staphylococcus aureus* undergoes major transcriptional reorganization during growth with *Enterococcus faecalis* in milk. *Food Microbiol.* 73, 17–28. doi: 10.1016/j.fm.2018.01.007
- Wang, X., Wang, S., and Zhao, H. (2019). Unravelling microbial community diversity and succession of Chinese Sichuan sausages during spontaneous fermentation by high-throughput sequencing. *J. Food Sci. Technol.* 56, 3254–63. doi: 10.1007/s13197-019-03781-y
- Warnecke, T., and Gill, R. T. (2005). Organic acid toxicity, tolerance, and production in *Escherichia coli* biorefining applications. *Microb. Cell Fact* 4:25. doi: 10.1186/1475-2859-4-25
- Wessel, A. K., Hmelo, L., Parsek, M. R., and Whiteley, M. (2013). Going local: technologies for exploring bacterial microenvironments. *Nat. Rev. Microbiol.* 11, 337–48. doi: 10.1038/nrmicro3010
- Wiernasz, N., Cornet, J., Cardinal, M., Pilet, M.-F., Passerini, D., and Leroi, F. (2017). Lactic acid bacteria selection for biopreservation as a part of hurdle technology approach applied on seafood. *Front. Mar. Sci.* 4:119. doi: 10.3389/fmars.2017.00119
- Wiernasz, N., Leroi, F., Chevalier, F., Cornet, J., Cardinal, M., Rohloff, J., et al. (2020). Salmon Gravlax biopreservation with lactic acid bacteria: a polyphasic approach to assessing the impact on organoleptic properties, microbial ecosystem and volatiles composition. *Front. Microbiol.* 10:3103. doi: 10.3389/fmicb.2019.03103
- Windholtz, S., Redon, P., Lacampagne, S., Farris, L., Lytra, G., Cameleyre, M., et al. (2021). Non-*Saccharomyces* yeasts as bioprotection in the composition of red wine and in the reduction of sulfur dioxide. *LWT* 149:111781. doi: 10.1016/j.lwt.2021.111781
- World Health Organization (2015). *WHO Estimates of the Global Burden of Foodborne Diseases: Foodborne Disease Burden Epidemiology Reference Group 2007-2015*. Geneva: World Health Organization.
- Wu, W., Deng, G., Liu, C., Gong, X., Ma, G., Yuan, Q., et al. (2020). Optimization and multiomic basis of phenyllactic acid overproduction by *Lactobacillus plantarum*. *J. Agric. Food Chem.* 68, 1741–9. doi: 10.1021/acs.jafc.9b07136
- Yi, L., Luo, L., and Lü, X. (2018). Efficient exploitation of multiple novel bacteriocins by combination of complete genome and peptidome. *Front. Microbiol.* 9:1567. doi: 10.3389/fmicb.2018.01567
- Zagorec, M., and Champomier-Vergès, M.-C. (2017). *Lactobacillus sakei*: a starter for sausage fermentation, a protective culture for meat products. *Microorganisms* 5:56. doi: 10.3390/microorganisms5030056
- Zdenkova, K., Alibayov, B., Karamonova, L., Purkrtova, S., Karpiskova, R., and Demnerova, K. (2016). Transcriptomic and metabolic responses of *Staphylococcus aureus* in mixed culture with *Lactobacillus plantarum*, *Streptococcus thermophilus* and *Enterococcus durans* in milk. *J. Ind. Microbiol. Biotechnol.* 43, 1237–47. doi: 10.1007/s10295-016-1794-y
- Zhang, Y., Zhu, L., Dong, P., Liang, R., Mao, Y., Qiu, S., et al. (2018). Bio-protective potential of lactic acid bacteria: effect of *Lactobacillus sakei* and *Lactobacillus curvatus* on changes of the microbial community in vacuum-packaged chilled beef. *Asian Austral. J. Anim. Sci.* 31, 585–94. doi: 10.5713/ajas.17.0540